



# Diphenyleneiodonium, an inhibitor of NOXes and DUOXes, is also an iodide-specific transporter<sup>☆</sup>

C. Massart<sup>a,\*</sup>, N. Giusti<sup>a</sup>, R. Beauwens<sup>b</sup>, J.E. Dumont<sup>a</sup>, F. Miot<sup>a</sup>, J. Van Sande<sup>a</sup>

<sup>a</sup>IRIBHM, School of Medicine, ULB, Université Libre de Bruxelles, Campus Erasme, Route de Lennik 808, Belgium

<sup>b</sup>Laboratory of Physiology, School of Medicine, ULB, Université Libre de Bruxelles, Campus Erasme, Route de Lennik 808, Belgium

## ARTICLE INFO

### Article history:

Received 27 September 2013

Received in revised form 7 November 2013

Accepted 28 November 2013

### Keywords:

Diphenyleneiodonium

O<sub>2</sub> radical generation

NOX

DUOX

Iodide efflux

Thyroid

## ABSTRACT

**NADPH oxidases (NOXes) and dual oxidases (DUOXes) generate O<sub>2</sub><sup>•−</sup> and H<sub>2</sub>O<sub>2</sub>. Diphenyleneiodonium (DPI) inhibits the activity of these enzymes and is often used as a specific inhibitor. It is shown here that DPI, at concentrations similar to those which inhibit the generation of O<sub>2</sub> derivatives, activated the efflux of radioiodide but not of its analog <sup>99m</sup>TcO<sub>4</sub><sup>−</sup> nor of the K<sup>+</sup> cation mimic <sup>86</sup>Rb<sup>+</sup> in thyroid cells, in the PCC13 rat thyroid cell line and in COS cell lines expressing the iodide transporter NIS. Effects obtained with DPI, especially in thyroid cells, should therefore be interpreted with caution.**

© 2013 The Authors. Published by Elsevier B.V. on behalf of Federation of European Biochemical Societies. All rights reserved.

## 1. Introduction

Seven specific O<sub>2</sub><sup>•−</sup> and H<sub>2</sub>O<sub>2</sub> generating enzymes have been demonstrated, the NADPH oxidases, NOXes and dual oxidases DUOXes. In many cell types, they generate both intracellular and extracellular O<sub>2</sub><sup>•−</sup> and H<sub>2</sub>O<sub>2</sub>. They share many properties, including enzymatic mechanisms, but differ in terms of function, tissue distribution and control [1–3].

Diphenyleneiodonium (DPI) inhibits the activity of these flavoenzymes. It is often considered as a specific inhibitor. Inhibition of any metabolic step by DPI is therefore often attributed to this effect [4–9]. Although caution has been advised against the use of inhibition by DPI as a specific biomarker of NOX and DUOX actions [4], such arguments are still commonly used [10].

In this article we demonstrate another action of DPI which raises new interesting questions. In the course of our study of iodide metabolism in thyroid cells we observed that it increased the release of <sup>125</sup>I iodide but not <sup>99m</sup>TcO<sub>4</sub><sup>−</sup>, thus behaving as a specific carrier of iodide.

<sup>☆</sup>This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Abbreviations:** DPI, diphenyleneiodonium; DUOXes, dual oxidases; HRP, horse radish peroxidase type II; HVA, homovanillic acid; KRH, Krebs–Ringer Hepes medium; MMI, methylmercaptoimidazole; NIS, sodium/iodide symporter; NOXes, NADPH oxidases.

\* Corresponding author. Tel.: +32 25554173.

E-mail address: [cmassart@ulb.ac.be](mailto:cmassart@ulb.ac.be) (C. Massart).

This shows an effect of DPI other than the inhibition of H<sub>2</sub>O<sub>2</sub> generation. It presents an interesting chemical property of this iodinated derivative and may suggest a therapeutic application.

## 2. Materials and methods

### 2.1. Reagents

<sup>125</sup>I as NaI (~17Ci(629GBq)/mg) and <sup>86</sup>Rb<sup>+</sup> (19.3 mCi(714.1 MBq)/mg) were purchased from Perkin Elmer Life and Analytical Science (Zaventem, Belgium), <sup>99m</sup>Tc as Na Pertechnetate (<sup>99m</sup>TcO<sub>4</sub><sup>−</sup>) was provided to us by the Medical School Hospital. It was prepared daily, carrier free, from a <sup>99m</sup>Tc generator (Ultra TechneKow, Mallinckrodt Nuclear Medicine). We used around 1 μCi tracer per dish.

Homovanillic acid (HVA), horse radish peroxidase type II (HRP), methylmercaptoimidazole (MMI) and diphenyleneiodonium chloride (DPI) were provided by Sigma–Aldrich BVBA (Diegem, Belgium). Sodium iodide and sodium perchlorate were purchased from Merck KGaA (Darmstadt, Germany).

### 2.2. Tissue

Pig thyroids were collected at a local slaughterhouse. The fresh tissue was sliced at room temperature with a Stadie–Riggs microtome (Arthur Thomas, Philadelphia, PA, USA). Slices (0.3 mm thick, 45–60 mg wet weight) were incubated in 2 ml incubation medium.

### 2.3. Cell culture

PCC13 cells, a rat thyroid cell line, were grown in Coon's modified Ham's F12 medium supplemented with 5% decomplexed fetal bovine serum, bovine TSH (1 mU/ml) transferrin (5 µg/ml), insulin (1 µg/ml), penicillin (100 IU/ml), streptomycin (100 µg/ml) and fungizone (2.5 µg/ml). COS–NIS cells are COS–7 cells transfected with the wild type human sodium/iodide symporter (NIS) cDNA. Stable cell lines were selected by resistance to geneticin [11]. They were cultured in DMEM supplemented with 10% fetal bovine serum, Na pyruvate (1 mM), penicillin (100 IU/ml), streptomycin (100 µg/ml) and fungizone (2.5 µg/ml).

### 2.4. Incubation

The slices and cells were incubated in Krebs–Ringer Hepes medium (NaCl 124 mM; KCl 5 mM; MgSO<sub>4</sub> 1.25 mM; CaCl<sub>2</sub> 1.45 mM; Hepes Buffer 25 mM) supplemented with glucose 8 mM and BSA (bovine serum albumin) 0.5 mg/ml [12].

### 2.5. Measurement of the H<sub>2</sub>O<sub>2</sub> production

Thyroid slices were preincubated 1 h in KRH medium and then transferred to fresh medium containing 0.1 mg/ml HRP, 440 µM HVA and the tested agonists for 90 min. When H<sub>2</sub>O<sub>2</sub> is produced by the tissue, the HVA is oxidized into a fluorescent dimer by the peroxidase. The fluorescence of the medium was measured at 315 and 425 nm, excitation and emission wavelengths respectively [13]. Generation of H<sub>2</sub>O<sub>2</sub> was expressed as ng H<sub>2</sub>O<sub>2</sub> produced by 100 mg of tissue, or ng H<sub>2</sub>O<sub>2</sub> per dish in the case of cells, during the length of the incubation.


### 2.6. Measurement of iodide uptake

The thyroid slices were preincubated 1 h in KRH supplemented with 0.1 mM MMI to block organification of iodide, then incubated for 1 h with fresh KRH medium supplemented with <sup>125</sup>I<sup>-</sup> (1 µCi/ml), 10<sup>-7</sup> M KI, 0.1 mM MMI and the tested agents, washed and counted. The uptake was evaluated by the ratio: counts per min <sup>125</sup>I<sup>-</sup>/100 mg tissue to counts per min <sup>125</sup>I<sup>-</sup>/100 µl medium [14]. Uptake of investigated cells in culture was divided by the uptake of cells incubated in the presence of NaClO<sub>4</sub> 1 mM to obtain concentration ratios (cell to medium C/M).


### 2.7. Measurement of tracer efflux from cells

300,000 cells were seeded in 35-mm-diameter dishes and grown 24 h in the case of COS–NIS cells or 48 h in the case of PCC13 cells, in their respective culture medium. The cells were then rinsed and incubated under slight agitation in KRH medium supplemented with KI 10<sup>-7</sup> M and 1 µCi/ml <sup>125</sup>I<sup>-</sup>, or 1 µCi/ml <sup>99m</sup>TcO<sub>4</sub><sup>-</sup>, or 1 µCi/ml <sup>86</sup>Rb<sup>+</sup>. As no iodination takes place in COS or PCC13 cells, MMI was not used routinely in these experiments. For <sup>86</sup>Rb<sup>+</sup> uptake, the K<sup>+</sup> concentration in the medium was 0.5 mM instead of 5 mM.

After 45 min uptake with <sup>125</sup>I<sup>-</sup> or <sup>99m</sup>TcO<sub>4</sub><sup>-</sup> and 2 h with <sup>86</sup>Rb<sup>+</sup>, the cells were rinsed twice with KRH and incubated with fresh medium, which was supplemented with KI (10<sup>-7</sup> M) and ClO<sub>4</sub><sup>-</sup> (1 mM) in the case of <sup>125</sup>I<sup>-</sup> or <sup>99m</sup>TcO<sub>4</sub><sup>-</sup> efflux, to avoid tracer recirculation. DPI (10 µM) is added at the start of this second incubation. Fifty microliter from 1 ml medium was withdrawn at the mentioned times and counted. At the end of incubation, the cells were rinsed twice with KRH medium kept on ice, then dissolved in 1 M NaOH and counted. The efflux was expressed in percent of the total radioactivity taken up by the cells (cells + medium + serial aliquots).



**Fig. 1.** Effect of DPI on iodide uptake level at equilibrium (Empty columns) and H<sub>2</sub>O<sub>2</sub> production (Filled columns) in pig thyroid slices. The incubation lasted 90 min. The results are the mean of 2 pig thyroids. The uptake control value, in absence of DPI, were 23.6 ± 4 and 18.5 ± 3 (mean of duplicates) in pig 1 and 2 respectively. The H<sub>2</sub>O<sub>2</sub> control value, in absence of DPI, were 421 ± 12 and 482 ± 18 (mean of duplicates) ng H<sub>2</sub>O<sub>2</sub>/100 mg wet weight tissue for pig 1 and 2 respectively.



**Fig. 2.** Comparison between the effect of the same range of DPI (Empty columns) and iodide (Filled columns) concentrations on iodide uptake value at equilibrium in pig thyroid slices. The incubation with DPI or KI lasted 1 h. The results presented are the mean of 2 pig thyroids. The uptake control value were 12.05 ± 1.4 and 7.1 ± 0.1 (means of duplicates) in pig 1 and 2 respectively. (Ratio of tissue vs medium radioactivity)


## 3. Results

DPI, in the same concentration range, inhibits both H<sub>2</sub>O<sub>2</sub> generation and radioiodide uptake in pig thyroid slices (Fig. 1).


The effect of DPI on iodide uptake level at equilibrium results from the balance between the continuous uptake by NIS and the release. The inhibition of iodide uptake by DPI takes place at a lower concentration (IC<sub>50</sub> 1–4 µM) than the effect of competing iodide (IC<sub>50</sub> 40 µM) (Fig. 2). Moreover the C–I bond of the iodine to the benzene ring of DPI is not labile [9]. The inhibition therefore is not due to iodide release from DPI.

For the release experiments (Fig. 3, 4, 5), DPI is added after the labeling of the cells and the release is measured immediately after; it is, in effect, quasi instantaneous. The effect of DPI does not correspond to a general membrane permeabilization as preloaded <sup>86</sup>Rb<sup>+</sup>, used as analog of K<sup>+</sup>, is not discharged from PCC13 nor COS–NIS cells (Fig. 3).


Moreover, DPI is a carrier more specific than NIS, the transporter



**Fig. 3.** Comparison of DPI effect on  $^{125}\text{I}^-$  and  $^{86}\text{Rb}^+$  efflux from COS-NIS preloaded cells. The efflux of the tracer from the cell to the medium is expressed as percentage of total uptake by the cells. The  $^{125}\text{I}^-$  uptake at equilibrium was 24.6 in this experiment. (Ratio of control vs  $\text{NaClO}_4$  treated cells). Empty circles: control. Filled square: DPI 10  $\mu\text{M}$ .



**Fig. 4.** Comparison of DPI effect on  $^{125}\text{I}^-$  and  $^{99\text{m}}\text{TcO}_4^-$  efflux from PCCl<sub>3</sub> and COS-NIS preloaded cells. DPI was added at the beginning of the efflux. The uptake values at equilibrium for  $^{125}\text{I}^-$  were  $24 \pm 1$  and  $24 \pm 0.05$  in PCCl<sub>3</sub> and COS-NIS cells respectively. The uptake values at equilibrium for  $^{99\text{m}}\text{TcO}_4^-$  were  $216 \pm 6$  and  $228 \pm 9$  in PCCl<sub>3</sub> and COS-NIS cells respectively (mean of duplicates). The results of  $^{125}\text{I}^-$  and  $^{99\text{m}}\text{TcO}_4^-$  efflux are expressed as percentage of total uptake by the cells. Empty circles: control. Filled square: DPI 10  $\mu\text{M}$ .



**Fig. 5.** Effect of temperature on DPI action on  $^{125}\text{I}^-$  efflux from PCCl3 preloaded cells. The cells were loaded with  $^{125}\text{I}^-$  at 37 °C for 45 min, then quickly rinsed with  $2 \times 1$  ml KRH at 37 °C or 22 °C or 15 °C. One milliliter of KRH at 37 °C ( $\square$ ) or 22 °C ( $\Delta$ ) or 15 °C ( $\circ$ ) was then added and efflux measured at those temperatures. The arrows bind the basal and DPI stimulated release curves at each temperature. Empty symbols: basal release. Filled symbols: DPI stimulated release.

of iodide ( $\text{Na}^+/\text{I}^-$  symporter) as it does not release  $^{99\text{m}}\text{TcO}_4^-$  from preloaded cells. Thus, contrary to the NIS [15], DPI does not recognize  $^{99\text{m}}\text{TcO}_4^-$  as radioiodide (Fig. 4).

Spontaneous iodide release is much decreased when the temperature is lowered from 37 to 22 °C and 15 °C (Fig. 5). However, the relative effect of DPI is as important at the three temperatures.

To check the reversible or permanent character of DPI effect on iodide uptake (through its effect on efflux) and on  $\text{H}_2\text{O}_2$  production we designed a protocol made of 3 sequential incubations called A, B and C. The measurement of iodide uptake (in the presence of  $^{125}\text{I}^-$  and  $10^{-7}$  M KI, for 45 min) and of  $\text{H}_2\text{O}_2$  production (in the presence of 440  $\mu\text{M}$  HVA and 100  $\mu\text{g}/\text{ml}$  Peroxidase II for 90 min), took place in the last incubation C. Incubation A and B lasted 45 min each for either parameter. No DPI was added in any of these incubations, for the control. To test if DPI effect was short or long lasting we added it at 3 different times: In A only, or in B only, or in C only. The results presented in Fig. 6 show that the effect of DPI on iodide uptake is strongest when added in C, but fade away if it is added in B only and more so if added in A only, which suggests that the DPI involved in iodide release is rapidly washed in the absence of DPI.


The effect of DPI on  $\text{H}_2\text{O}_2$  is much more persistent. Indeed, if added previously, even in the first incubation A only, it works as well. This suggests a more stable binding of the DPI involved in this inhibition.

#### 4. Discussion

We have shown that DPI, at concentrations of the same order as those that inhibit  $\text{H}_2\text{O}_2$  generation by thyroid cells, also causes an immediate specific release of iodide from thyroid and non-thyroid cells.

The effect of  $\text{NaClO}_4$  or  $\text{NaSCN}$  which only compete with the uptake of iodide and decrease by this mechanism the cell/medium equilibrium level [15] is relatively slow. The immediate effect of DPI on iodide efflux carried out in the presence of perchlorate is therefore not due to an inhibitory role on iodide uptake by NIS. It does not seem to require the specific iodide channel(s) involved in the efflux of iodide at the apical membrane of the thyrocyte [16,17], as it is reproduced in COS–NIS cells i.e. in non-thyroid cells.

Besides, an effect on NIS would similarly affect the efflux of



**Fig. 6.** Study of the reversibility of DPI effect on  $\text{H}_2\text{O}_2$  production or iodide uptake (C/M) in PCCl3 cells. The experimental protocol is made up of 3 sequential incubations called A, B or C.  $^{125}\text{I}^-$  for uptake (C/M) measurements or HVA and peroxidase II for  $\text{H}_2\text{O}_2$  measurements were added in incubation C only, which lasted 45 min for  $^{125}\text{I}^-$  uptake and 90 min for  $\text{H}_2\text{O}_2$  production. Incubation A and B lasted 45 min each. DPI 10  $\mu\text{M}$  was added either in A alone or in B alone, or in C alone. Empty columns: control. Filled columns: DPI 10  $\mu\text{M}$ .

$^{99\text{m}}\text{TcO}_4^-$  [15,18]. The absence of effect on  $^{99\text{m}}\text{TcO}_4^-$  or  $^{86}\text{Rb}^+$  release, excludes a general permeabilization of the cells by DPI.

Other unspecific effects of DPI have been observed but at higher concentrations than those effective on NOXes. For example, DPI (10  $\mu\text{M}$ ) also directly inhibits cholinesterase activity and to a lesser extent an internal  $\text{Ca}^{++}$  pump [19]. The effect of DPI on iodide efflux is not due to the inhibition of  $\text{H}_2\text{O}_2$  generation as it occurs in COS–NIS cells without DUOX and is, contrary to the effect on DUOX, relieved by incubation without the drug.

We have no explanation for this effect on iodide. Certainly because the covalent link in DPI between iodine and the other part of the DPI molecule is not labile, there is no dynamic exchange with medium iodide. The molecule is stable under stringent conditions [9].

Moreover, higher concentrations of iodide than DPI are needed to reach the same inhibition of C/M.

We can therefore only speculate on the mechanism of the DPI effect. A first explanation would be the rapid activation of an anion channel in the cell [20]. However DPI also acts on COS–NIS cells [11] which do not express any iodide specific channel, especially a channel distinguishing iodide and  $^{99\text{m}}\text{TcO}_4^-$ . A second explanation could be that DPI behaves as a membrane ionophore specific for iodide. Catalysis of an anion/hydroxyl ion exchange across mitochondrial membrane by DPI has been reported [21]. The fact that depressing membranes fluidity by decreasing the temperature of the cells slows down both basal and the increased efflux, does not distinguish the hypothesis of DPI acting as a shuttle for iodide from the concept that it acts as an intramembrane channel as is the case for valinomycin

and  $K^+$  [22,23]. An ionic type of carrier mechanism would be similar to the mitochondrial protein carrier mechanism which is believed to account for the uncoupling effect on mitochondrial oxidative phosphorylation by dinitrophenol. Both hypotheses of channel and shuttle, would suggest a specific affinity of DPI molecule for iodide.

Nevertheless the present data confirm that DPI cannot be considered as a specific inhibitor, of NOXes and DUOXes. They suggest that DPI or an analog with similar properties could therefore be considered as drugs for iodine or amiodarone induced hyperthyroidism as they would combine in one molecule the inhibition of thyroid hormone synthesis and iodide uptake (in fact retention) as a combination of antithyroperoxidase drugs and perchlorate.

## Acknowledgments

The authors thank Dr. J. Wolff (NIH, Bethesda) for critical reading of the manuscript, the Fondation Solvay and the Fondation Roi Baudouin “Fonds Yvonne Smits” and the Fonds de la Recherche Scientifique Médicale (FRSM) for their support and Mrs. D. Leemans for the preparation of the manuscript.

## References

- [1] Cross, A.R. and Segal, A.W. (2004) The NADPH oxidase of professional phagocytes – prototype of the NOX electron transport chain systems. *Biochim. Biophys. Acta* 1657, 1–22.
- [2] Katsuyama, M., Matsuno, K and Yabe-Nishimura, C. (2012) Physiological roles of NOX/NADPH oxidase, the superoxide-generating enzyme. *J. Clin. Biochem. Nutr.* 50, 9–22.
- [3] Lambeth, J.D. (2004) NOX enzymes and the biology of reactive oxygen. *Nat. Rev. Immunol.* 4, 181–189.
- [4] Aldieri, E., Riganti, C., Polimeni, M., Gazzano, E., Lussiana, C., Campia, I. et al. (2008) Classical inhibitors of NOX NAD(P)H oxidases are not specific. *Curr. Drug Metab.* 9, 686–696.
- [5] Bauernfeind, F., Bartok, E., Rieger, A., Franchi, L., Nunez, G. and Hornung, V. (2011) Cutting edge: reactive oxygen species inhibitors block priming, but not activation, of the NLRP3 inflammasome. *J. Immunol.* 187, 613–617.
- [6] Cross, A.R. and Jones, O.T. (1986) The effect of the inhibitor diphenylene iodonium on the superoxide-generating system of neutrophils. Specific labelling of a component polypeptide of the oxidase. *Biochem. J.* 237, 111–116.
- [7] Doussiere, J. and Vignais, P.V. (1991) Inhibition of  $O_2^-$  generating oxidase of neutrophils by iodonium biphenyl in a cell free system: effect of the redox state of the oxidase complex. *Biochem. Biophys. Res. Commun.* 175, 143–151.
- [8] O'Donnell, V.B., Smith, G.C. and Jones, O.T. (1994) Involvement of phenyl radicals in iodonium inhibition of flavoenzymes. *Mol. Pharmacol.* 46, 778–785.
- [9] O'Donnell, B.V., Tew, D.G., Jones, O.T. and England, P.J. (1993) Studies on the inhibitory mechanism of iodonium compounds with special reference to neutrophil NADPH oxidase. *Biochem. J.* 290(Pt 1), 41–49.
- [10] Doroshov, J.H., Gaur, S., Markel, S., Lu, J., van Balgooy, J., Synold, T.W. et al. (2013) Effects of iodonium-class flavin dehydrogenase inhibitors on growth, reactive oxygen production, cell cycle progression, NADPH oxidase 1 levels, and gene expression in human colon cancer cells and xenografts. *Free Radic. Biol. Med.* 57, 162–175.
- [11] Pohlenz, J., Duprez, L., Weiss, R.E., Vassart, G., Refetoff, S. and Costagliola, S. (2000) Failure of membrane targeting causes the functional defect of two mutant sodium iodide symporters. *J. Clin. Endocrinol. Metab.* 85, 2366–2369.
- [12] Corvilain, B., Van Sande, J., Laurent, E. and Dumont, J.E. (1991) The  $H_2O_2$ -generating system modulates protein iodination and the activity of the pentose phosphate pathway in dog thyroid. *Endocrinology* 128, 779–785.
- [13] Benard, B and Brault, J (1971) Production of peroxide in the thyroid. *Union Med. Can.* 100, 701–705.
- [14] Van Sande, J., Lamy, F., Lecocq, R., Mirkine, N., Rocmans, P., Cochaux, P. et al. (1988) Pathogenesis of autonomous thyroid nodules: in vitro study of iodine and adenosine 3',5'-monophosphate metabolism. *J. Clin. Endocrinol. Metab.* 66, 570–579.
- [15] Van Sande, J., Massart, C., Beauwens, R., Schoutens, A., Costagliola, S., Dumont, J.E. et al. (2003) Anion selectivity by the sodium iodide symporter. *Endocrinology* 144, 247–252.
- [16] Bizhanova, A. and Kopp, P. (2011) Controversies concerning the role of pendrin as an apical iodide transporter in thyroid follicular cells. *Cell Physiol. Biochem.* 28, 485–490.
- [17] Twyffels, L., Massart, C., Golstein, P.E., Raspe, E., Van Sande, J., Dumont, J.E. et al. (2011) Pendrin: the thyrocyte apical membrane iodide transporter? *Cell Physiol. Biochem.* 28, 491–496.
- [18] Dohan, O., Portulano, C., Basquin, C., Reyna-Neyra, A., Amzel, L.M. and Carrasco, N. (2007) The  $Na^+/I^-$  symporter (NIS) mediates electroneutral active transport of the environmental pollutant perchlorate. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20250–20255.
- [19] Tazzeo, T., Worek, F. and Janssen, L. (2009) The NADPH oxidase inhibitor diphenyleneiodonium is also a potent inhibitor of cholinesterases and the internal  $Ca^{2+}$  pump. *Br. J. Pharmacol.* 158, 790–796.
- [20] Gatley, S.J. and Martin, J.L. (1979) Some aspects of the pharmacology of diphenyleneiodonium, a bivalent iodine compound. *Xenobiotica* 9, 539–546.
- [21] Holland, P.C. and Sherratt, H.S. (1972) Biochemical effects of the hypoglycaemic compound diphenyleneiodonium. Catalysis of anion-hydroxyl ion exchange across the inner membrane of rat liver mitochondria and effects on oxygen uptake. *Biochem. J.* 129, 39–54.
- [22] Andreoli, T.E., Tieffenberg, M. and Tosteson, D.C. (1967) The effect of valinomycin on the ionic permeability of thin lipid membranes. *J. Gen. Physiol.* 50, 2527–2545.
- [23] Golstein, P., Abramow, M., Dumont, J.E. and Beauwens, R. (1992) The iodide channel of the thyroid: a plasma membrane vesicle study. *Am. J. Physiol.* 263, C590–C597.