

Contents

I	Introduction and Background	1
1	Introduction	2
1.1	General context	2
1.2	Outline of the thesis	4
2	The gauge/gravity correspondence	7
2.1	D-branes	7
2.1.1	Chan-Paton factors	9
2.2	The correspondence	10
2.2.1	The example of $\mathcal{N} = 4$ SYM	11
2.2.2	The example of D0-brane quantum mechanics	12
2.3	Take-home message	13
II	Simple models of Holography	14
3	Simple models of Holography	15
3.1	General properties	15
3.1.1	Chaotic behaviour of correlators	16
3.1.2	Models with probes	17
3.2	The IP and IOP models	19
3.2.1	Introduction	19
3.2.2	Definition of the IP model	20
3.2.3	Large M computation	20
3.2.4	Bose-Einstein condensation	22
3.2.5	Definition of the IOP model	25
3.2.6	Large M computation	25
3.2.7	Concluding remarks	27
3.3	Melonic models	27
3.3.1	The Sachdev-Ye-Kitaev model	28
3.3.2	The Gurau-Witten model	30
3.3.3	Quartic matrix-vector model	31

4	Diagrammatics and bilocal structure of melonic models	33
4.1	SYK diagrammatics	33
4.2	Large D matrix model diagrammatics	35
4.2.1	Models	35
4.2.2	Genus of a vertex	35
4.2.3	Standard scaling vs. new scaling	36
4.2.4	Proof	37
4.3	More on the new large D limit of matrix models	39
4.3.1	Multi-trace models	39
4.3.2	Models with reduced symmetry	39
4.3.3	Correlation functions	40
4.3.4	Model building	40
4.4	Bilocal structure	41
4.4.1	An instructive computation	41
5	Physical properties of melonic models	44
5.1	Two-point functions and emergent symmetry	44
5.2	Low temperature entropy	47
5.3	Four-point functions and chaos	47
5.4	Phase structure of complex melonic models	50
6	Quartic matrix-vector models	52
6.1	Definition and properties of the model	52
6.1.1	Leading diagrams for $\lambda_1 = 0$	53
6.1.2	Including the $I_{4,1}$ interaction term	55
6.1.3	Two-point functions	56
6.1.4	Writing the free energy in terms of the Matsubara coefficients G_k	57
6.2	Probe analysis	61
6.2.1	Generalities	62
6.2.2	A toy example	63
6.2.3	Probe analysis of the quartic matrix-vector model	65
6.2.4	Test of the fundamental relation for the on-shell probe action	71
III	Localization in $\mathcal{N} = 2$ SQCD	74
7	Models and observables	75
7.1	Introduction	75
7.2	$\mathcal{N} = 4$ SYM and $\mathcal{N} = 2$ SCQCD	76
7.3	Supersymmetric Wilson loops	80
7.3.1	The circular $1/2$ BPS Wilson loop in $\mathcal{N} = 2$ SYM	81

8	Localization	83
8.1	Overview	83
8.1.1	Super Yang-Mills on S^4	84
8.2	Correlators of chiral operators	89
8.2.1	From S^4 to \mathbb{R}_4	89
8.2.2	Localization results	91
8.2.3	Perturbative checks	93
8.3	1/2-BPS circular Wilson loops	96
8.3.1	Localization results in $\mathcal{N} = 4$ SYM and their interpretation	96
8.3.2	Localization results in $\mathcal{N} = 2$ SCQCD and their interpretation	99
8.4	Correlators between Wilson loop and chiral operators in $\mathcal{N} = 2$ conformal gauge theories	101
8.4.1	Wilson loop and its correlators with chiral operators . . .	101
8.4.2	The matrix model approach	102
8.4.3	Perturbative checks in field theory	102
A	Publication I:	
	More on the new large D limit of matrix models	103
B	Publication II:	
	Correlators between Wilson loop and chiral operators in $\mathcal{N} = 2$ conformal gauge theories	123
	Bibliography	162