1 Retrieval and characterization of carbon monoxide (CO)
2 vertical profiles in the Martian atmosphere from
3 observations of PFS/MEX

4 Jimmy Bouche^a, Sophie Bauduin^a, Marco Giuranna^b, Séverine Robert^c,
5 Shohei Aoki^c, Ann Carine Vandaele^c, Justin T. Erwin^c, Frank Daerden^c,
6 Paulina Wolkenberg^{b,d}, Pierre-François Coheur^a

7 ^a*Université libre de Bruxelles (ULB), Atmospheric spectroscopy, Service de Chimie*
8 *Quantique et Photophysique CP 160/09, Avenue F.D. Roosevelt 50, Bruxelles, Belgium.*

9 ^b*National Institute of Astrophysics INAF-IAPS, Rome, Italy.*

10 ^c*Royal Belgian Institute for Space Aeronomy, Brussels, Belgium.*

11 ^d*Space Research Centre of Polish Academy of Sciences, Warsaw, Poland.*

12 **Abstract**

13 The knowledge of the carbon monoxide (CO) abundance on Mars is es-
14 sential in order to assess the processes driving the carbon cycle on the planet.
15 Solar occultation measurements provide vertically-resolved measurements of
16 CO from a few kilometers to higher altitudes and can be complemented by
17 nadir measurements to enhance the spatial coverage of observations and the
18 monitoring of the near-surface layer. Up to now, in the shortwave, CO re-
19 trievals from nadir observations have, however, mostly been performed on
20 mean spectra and only total column abundances have been obtained. In this
21 work we explore the possibility of exploiting nadir measurements from the
22 Planetary Fourier Spectrometer (PFS) in the 1–0 band of CO (centered at
23 2143 cm⁻¹) to retrieve vertical profiles of that species on individual mea-
24 surement. The retrievals are performed for a set of 16 selected PFS spectra
25 with reasonable signal-to-noise ratios by applying the Optimal Estimation
26 Method (OEM) with appropriate constraints, built from model simulations
27 of the Martian atmosphere. The retrieved profiles are characterized in terms
28 of vertical sensitivity and errors. We demonstrate, in particular, that the
29 PFS nadir measurements carry information mostly on the CO column below
30 15 km, with a maximum sensitivity to the near-surface atmosphere. These
31 measurements allow to substantially reduce the prior uncertainty on the CO
32 abundance in this altitude range, with an estimated total retrieval error on

the column-averaged volume mixing ratio (VMR) around 10%. We show that the set of retrieved VMRs are in the range of values reported from other instruments. The retrieved VMRs also capture well the known spatial and seasonal CO variability, which is promising in the perspective of better exploiting the exceptional set of PFS observations on Mars.

Keywords:

PFS, Mars Express, carbon monoxide, vertical profiles, Optimal Estimation, Martian atmosphere, thermal infrared, nadir

1. Introduction

The first detection of carbon monoxide (CO) in the atmosphere of Mars has been made 50 years ago by [1] using ground-based observations. Since then, it has been discovered that CO plays a major role in the carbon photochemical cycle. CO is indeed formed by the photolysis of carbon dioxide (CO₂) and its recombination with oxygen (O) atoms is at the center of the so-called stability problem of CO₂ in the Martian atmosphere. This reaction is spin-forbidden and extremely slow compared to the recombination of O atoms into O₂. As a consequence, without an efficient recycling process, the total amount of CO₂ would have been converted into its photolysis products, and the abundance of CO and O₂ in the Martian atmosphere would have significantly exceeded the present amount. In reality, as proposed in the 1970's by [2] and [3], the recombination of CO into CO₂ is catalysed by odd hydrogen (H, OH and HO₂ and generally written HO_x) formed by the photodissociation of water vapour (H₂O). These earlier models were, however, not completely satisfactory; they used either an eddy diffusion coefficient unacceptably large for the middle atmosphere of Mars [2] or an amount of H₂O significantly exceeding the known values [3]. Although more recent models have solved some of these shortcomings [4, 5, 6, 7, 8], the global carbon cycle on Mars is nowadays still not completely understood. The contribution of heterogeneous chemistry [9, 10] in the different processes at play is part of the remaining uncertainties. The recently reported but debated detection of atmospheric methane, combined with its large apparent seasonal variability (e.g. [11, 12, 13] and references therein) has challenged further the understanding of the carbon cycle. To improve on this understanding, significantly more accurate observations of the spatial, vertical and temporal variability of the different species involved in the carbon cycle on Mars are obviously

68 needed.

69 This has already been partly the case for CO, for which the knowledge
70 of spatial and seasonal variations in the Martian atmosphere has improved
71 thanks to different remote observations, made from ground or from space in
72 the last 15 years. Contrary to what has been considered for a long time,
73 it was shown that CO was not uniformly mixed but has strong latitudinal
74 and seasonal variations, intimately related to the condensation-sublimation
75 cycle of CO₂ at the poles. The first evidence of the variability came from
76 ground-based observations [14], that showed a north-to-south increase of the
77 CO volume mixing ratio (VMR) has been reported during the Northern sum-
78 mer due to the condensation of CO₂ at the south polar cap. Since then, a
79 more complete picture of the latitudinal-seasonal changes of CO has been
80 brought thanks to different instruments, such as OMEGA (Observatoire
81 pour la Minéralogie, l'Eau, les Glaces et l'Activité) [15] and PFS (Plane-
82 tary Fourier Spectrometer) [16, 17, 18] on-board Mars Express (MEX), and
83 CRISM (Compact Reconnaissance Imaging Spectrometer for Mars) [19, 20]
84 on-board the Mars Reconnaissance Orbiter (MRO). Note also that atmo-
85 spheric CO has been measured in other spectral regions, such as in the
86 submillimeter and the microwave [21, 22, 23, 24]. These measurements in
87 the pure rotational lines of CO have been helpful for measuring the thermal
88 structure of the Martian atmosphere as well as wind velocities with suffi-
89 cient resolution. However, there has been no systematic retrieval of the CO
90 vertical profile, especially from nadir observations. The only study of this
91 kind attempted to exploit different spectral ranges or instruments to retrieve
92 information on the CO vertical distribution [25], but these synergistic ap-
93 proaches are difficult to implement. This is in stark contrast to Earth, where
94 the retrieval of vertical profile is becoming standard. For CO in particular,
95 it has been shown for thermal infrared (TIR) sounders that, in case of high
96 temperature contrasts between the ground and the air above it (hereafter
97 called thermal contrast), two partial columns of CO can be decorrelated for
98 the lower part of the atmosphere (surface layer and rest of the troposphere)
99 [26]. If possible on Mars, the retrieval of CO profiles, even if weakly resolved,
100 would provide useful constraints for the models regarding the carbon chem-
101 istry in the atmosphere close to the surface, a region generally not accessible
102 to limb/solar occultation measurements. This work examines therefore the
103 possibility of retrieving information on the CO vertical distribution from
104 Martian nadir TIR observations. This is performed on a small set of single
105 PFS nadir spectra, chosen to represent different conditions of CO on Mars,

106 in the spectral region of the CO fundamental vibrational band (centered at
 107 2143 cm^{-1}). The Optimal Estimation Method (OEM) [27] is used, which
 108 allows for a detailed characterization of the retrieved CO profiles in terms of
 109 error budget and vertical sensitivity. Section 2 presents the PFS instrument
 110 and section 3 is dedicated to a short description of the radiative transfer.
 111 Section 4 introduces the OEM. The selection of PFS observations and the
 112 retrieval settings are described in section 5 and the results are discussed in
 113 section 6, with a detailed interpretation of the error budget of the retrieved
 114 CO. The conclusions of the study are summarized in section 7.

115 2. PFS instrument

116 PFS [28] is a double pendulum Fourier transform interferometer on-board
 117 the Mars Express spacecraft. It circles Mars on a quasi-polar orbit with an
 118 inclination of 87° and a pericentre altitude of around 250 km. It covers a
 119 large spectral range in the infrared ($250\text{--}8200\text{ cm}^{-1}$) thanks to two different
 120 channels: the Long Wavelength Channel (LWC, $250\text{--}1700\text{ cm}^{-1}$) and the
 121 Short Wavelength Channel (SWC, $1700\text{--}8200\text{ cm}^{-1}$). Both channels have a
 122 sampling step of 1 cm^{-1} and a spectral resolution of $\sim 1.3\text{ cm}^{-1}$, when no
 123 apodization function is applied, and of $\sim 1.8\text{ cm}^{-1}$ when a Hamming function
 124 is applied to the measured interferograms (as in the case of the present work).
 125 The instrument has an instantaneous field-of-view (IFOV) of about 1.6° full
 126 width at half maximum (FWHM) for the SWC and of 2.8° FWHM for the
 127 LWC, which correspond to a spatial resolution on ground of 7 and 12 km
 128 respectively, when PFS is at 250 km from Mars. A full description of the
 129 PFS design, specification and operation can be found in [28]. Additional
 130 details about the spectral and radiometric calibration procedure for both
 131 channels may be found in [29, 30].

132 Since the beginning of its science operation, PFS has been collecting Mar-
 133 tian atmospheric spectra, currently covering more than 7 Martian years (MY
 134 26 to MY 34). In this work, we have selected only several PFS nadir spec-
 135 tra associated to different Martian conditions, especially in terms of seasonal
 136 and latitudinal variations of CO (see section 5.1 for details). Among the two
 137 channels, only the SWC spectral range covers the vibrational bands of CO,
 138 as the 1–0 and the 2–0 centered at 2143 cm^{-1} and at 4235 cm^{-1} respectively.
 139 For this study, only the stronger 1–0 band of CO has been considered in
 140 apodized SWC spectra.

141 3. Radiative transfer

142 The monochromatic infrared radiance at wavenumber $\tilde{\nu}$ measured by PFS
 143 at the top of the Martian atmosphere (TOA), $I^\uparrow(\tilde{\nu}, \theta, \text{TOA})$, can be described
 144 by the sum of two terms: 1) the transmission of the radiance emitted and
 145 reflected by the Martian surface, and 2) the contribution of scattering and
 146 emission by the atmosphere. This is expressed by the equation of the radia-
 147 tive transfer:

$$I^\uparrow(\tilde{\nu}, \theta, \text{TOA}) = I^\uparrow(\tilde{\nu}, \theta, 0) t(\tilde{\nu}, \theta, 0, \text{TOA}) + \int_0^{\text{TOA}} J(\tilde{\nu}, \mathbf{\Omega}, z) \frac{\partial t(\tilde{\nu}, \theta, z, \text{TOA})}{\partial z} dz, \quad (1)$$

148 where θ is the ground zenith angle, $t(\tilde{\nu}, \theta, 0, \text{TOA})$ is the transmittance of
 149 the Martian atmosphere, $t(\tilde{\nu}, \theta, z, \text{TOA})$ is the transmittance between alti-
 150 tude z and TOA, $J(\tilde{\nu}, \mathbf{\Omega}, z)$ is the atmospheric source term which depends
 151 on thermal emission and scattering and where $\mathbf{\Omega}$ represents the geometric
 152 angles. Note that in this work, only clear or almost clear atmospheres (see
 153 section 5.1) have been analysed. In this case, scattering can be neglected
 154 and the atmospheric source function can be approximated by a Plank func-
 155 tion $B(\tilde{\nu}, T)$. The radiance at the surface, $I^\uparrow(\tilde{\nu}, \theta, 0)$, takes into account the
 156 emission of the ground and the reflection of the solar incoming radiation as
 157 follows:

$$I^\uparrow(\tilde{\nu}, \theta, 0) = \epsilon(\tilde{\nu})B(T_s) + (1 - \epsilon(\tilde{\nu}))I_0^{\downarrow\uparrow}(\tilde{\nu}) + r(\tilde{\nu}, \theta, \phi, \theta^*, \phi^*)F_0^{\downarrow\star}(\tilde{\nu}), \quad (2)$$

158 where $B(T_s)$ is the Planck function at surface temperature T_s , $\epsilon(\tilde{\nu})$ is the
 159 wavenumber dependent emissivity of the Martian surface, $I_0^{\downarrow\uparrow}(\tilde{\nu})$ is the mean
 160 radiance associated to the downward flux of the atmosphere and $F_0^{\downarrow\star}(\tilde{\nu})$ is the
 161 solar flux reaching Mars' surface. $r(\tilde{\nu}, \theta, \phi, \theta^*, \phi^*)$ is the effective reflectivity
 162 of the surface. θ^* and ϕ^* are respectively the zenithal and azimuthal angles
 163 of the Sun, and ϕ is the azimuthal angle of the spacecraft.

164 In order to calculate the TOA radiance, equation 1 is discretized into a
 165 set of layers, in which the averaged temperature and pressure and the partial
 166 columns of molecular species are calculated. The upwelling radiance at the
 167 top of layer i is estimated as

$$I_i^\uparrow = B(\bar{T}_i) + (I_{i-1}^\uparrow - B(\bar{T}_i))\tilde{t}_i, \quad (3)$$

168 where $B(\bar{T}_i)$ is the emission of layer i approximated as a blackbody at the av-
 169 eraged temperature $\bar{T}_i$. The effective transmittance $\tilde{t}_i$ of layer i is calculated
 170 using

$$\tilde{t}_i = \exp \left[- \sum_j PC_{i,j} \sum_{j,k} \sigma_{j,k}(\tilde{\nu}, \bar{P}_i, \bar{T}_i) \right], \quad (4)$$

171 where j is associated to the molecular species and k to the relevant absorption
 172 lines at the chosen $\tilde{\nu}$. $PC_{i,j}$ is the partial column of species j in layer i and
 173 $\sigma_{j,k}(\tilde{\nu}, \bar{P}_i, \bar{T}_i)$ is the absorption cross section of that species ($\text{cm}^2/\text{molecules}$)
 174 calculated for the mean parameters of layer i .

175 In this work, the Atmosphit software has been used for radiative transfer
 176 calculations. It is a versatile line-by-line radiative transfer code initially
 177 developed for Earth atmosphere, but which can easily be adapted for Mars.
 178 Details about the Atmosphit software can be found in [31], [32] and [33]. For
 179 this work, the emissivity and the effective reflectivity introduced in equation
 180 2 are considered constant with wavenumber, and for the latter, a combined
 181 contribution of lambertian and specular reflections of the Sun radiation is
 182 assumed.

183 4. Retrieval method: Optimal Estimation

184 4.1. Background

185 The method used in this work to retrieve the vertical profiles of CO is
 186 the OEM [27]. The idea of this method is to find the CO profile that is the
 187 most consistent with both PFS observations and a certain prior knowledge
 188 of its distribution in the Martian atmosphere. This is done by minimizing
 189 the following cost function ϕ

$$\phi = (\mathbf{y} - \mathbf{F}(\mathbf{x}, \mathbf{b}))^T \mathbf{S}_\epsilon^{-1} (\mathbf{y} - \mathbf{F}(\mathbf{x}, \mathbf{b})) + (\mathbf{x} - \mathbf{x}_a)^T \mathbf{S}_a^{-1} (\mathbf{x} - \mathbf{x}_a), \quad (5)$$

190 where $\mathbf{y}$ is the measurement vector, consisting of the PFS spectral radiances,
 191 $\mathbf{x}$ is the atmospheric state vector, containing the variables to be retrieved,
 192 i.e. the CO profile but also the other parameters (see section 5.3), $\mathbf{b}$ includes
 193 all the other fixed atmospheric variables that influence PFS measurements,
 194 $\mathbf{x}_a$ is the *a priori* state vector (including the *a priori* profile of CO) and $\mathbf{F}$
 195 is the forward radiative transfer model. The covariance matrices, $\mathbf{S}_\epsilon$ and $\mathbf{S}_a$,
 196 are respectively the measurement and the *a priori* covariance matrices.

197 The minimization of ϕ is iteratively performed by the Atmosphit software
 198 using the numerical Gauss-Newton method for moderately non-linear cases
 199 [27]. At each iteration, a new state vector is evaluated using

$$\mathbf{x}_{i+1} = \mathbf{x}_a + (\mathbf{K}_i^T \mathbf{S}_\epsilon^{-1} \mathbf{K}_i + \mathbf{S}_a^{-1})^{-1} \mathbf{K}_i^T \mathbf{S}_\epsilon^{-1} [\mathbf{y} - \mathbf{F}(\mathbf{x}_i, \mathbf{b}) + \mathbf{K}_i(\mathbf{x}_i - \mathbf{x}_a)], \quad (6)$$

200 where $\mathbf{K}$ is the Jacobian matrix, whose rows correspond to the derivatives of
 201 the TOA radiance with respect to the retrieved variables. The convergence
 202 is reached when the absolute difference between the forward models of the
 203 two last iterations is lower than 20% of the noise. Note that Atmosphit does
 204 not retrieve the partial column in each layer directly but instead a scaling
 205 factor for each layer, which is applied to the *a priori* profile at the end of
 206 each iteration.

207 4.2. Characterization of the retrievals

208 4.2.1. Vertical sensitivity

209 The OEM offers a very adequate framework to characterize the retrieved
 210 CO profiles in terms of vertical sensitivity. This is done in this work mostly
 211 in terms of the averaging kernel matrix $\mathbf{A}$. It is defined as the sensitivity of
 212 the retrieved state (hereafter symbolized by $\hat{\mathbf{x}}$) to the true state ($\mathbf{x}$) and can
 213 be calculated as

$$\mathbf{A} = \frac{\partial \hat{\mathbf{x}}}{\partial \mathbf{x}} = (\mathbf{K}^T \mathbf{S}_\epsilon^{-1} \mathbf{K} + \mathbf{S}_a^{-1})^{-1} \mathbf{K}^T \mathbf{S}_\epsilon^{-1} \mathbf{K}. \quad (7)$$

214 The rows of $\mathbf{A}$ define an averaging kernel function (AVK) for each element of
 215 $\hat{\mathbf{x}}$, which peaks at the altitude from where most of the information is coming.
 216 In other words, the AVK provide an estimation of the vertical sensitivity of
 217 the retrieval and peak at the altitude of maximum sensitivity. The trace of
 218 the $\mathbf{A}$ matrix gives the degree of freedom for signal (DOFS), defined as the
 219 number of independent pieces of information in the retrieved profile.

220 4.2.2. Error analysis

221 As for the vertical sensitivity, a complete error budget analysis can be
 222 performed with the OEM, by separating different sources of errors from the
 223 total [27]. The following sources of errors have been evaluated:

- 224 1. The smoothing error: due to the limited vertical resolution of nadir
 225 observations, the retrieved state is a smoothed estimate of the true

226 state. The resulting error can be characterized by the covariance matrix
 227 $\mathbf{S}_s$, calculated from the $\mathbf{A}$ and $\mathbf{S}_a$ matrices as follows

$$\mathbf{S}_s = (\mathbf{A} - \mathbf{I})\mathbf{S}_a(\mathbf{A} - \mathbf{I})^T. \quad (8)$$

228 2. The measurement error, which directly propagates from the instrumen-
 229 tal noise. Its associated covariance matrix can be written as:

$$\mathbf{S}_m = \mathbf{G}\mathbf{S}_\epsilon\mathbf{G}^T, \quad (9)$$

230 where $\mathbf{G}$ is the gain matrix, whose rows are the derivatives of the
 231 retrieved state with respect to the measurement. Note that in this work,
 232 forward modelling errors are included in the $\mathbf{S}_\epsilon$ matrix (see section 5.3).
 233 The covariance matrix $\mathbf{S}_m$ therefore describes the total error from the
 234 measurement noise and the forward model.

235 3. The error from the fitted parameters: this type of error comes from
 236 the interferences existing between the retrieved CO profile and the rest
 237 of the state vector (other retrieved parameters), such as the surface
 238 temperature (see section 5.3) [34]. Its covariance matrix is estimated
 239 using the non-diagonal blocs ($\mathbf{A}_{ir}$) of the AVK matrix (see [34] for
 240 details):

$$\mathbf{S}_f = \mathbf{A}_{ir}\mathbf{S}_{ra}\mathbf{A}_{ir}^T, \quad (10)$$

241 where $\mathbf{i}$ is the part of the state vector under consideration (the CO pro-
 242 file), $\mathbf{r}$ represents the rest of the state vector (the surface temperature
 243 and the CO₂ profile, see section 5.3), and $\mathbf{S}_{ra}$ is the *a priori* covariance
 244 matrix of $\mathbf{r}$.

245 The sum of the covariance matrices associated to these different uncer-
 246 tainties provide the total retrieval error covariance matrix:

$$\mathbf{S}_T = \mathbf{S}_s + \mathbf{S}_m + \mathbf{S}_f. \quad (11)$$

247 The square root of the diagonal elements of $\mathbf{S}_T$ can be seen as the error
 248 on the retrieved CO at each altitude.

249 5. Spectra selection and retrieval set-up

250 5.1. Selection of spectra

251 CO vertical profiles have been retrieved from a small set of single nadir
 252 PFS observations. These observations have been chosen among all PFS mea-
 253 surements recorded between 10 January 2004 (MY 26) and 18 June 2017 (MY

34), following a 2 step procedure. The first step consists in finding spectra with almost clear sky conditions and sufficient signal-to-noise ratio (SNR) to detect CO lines in the 1–0 band. For this, only PFS observations associated to dust and ice column-integrated optical depths (τ_d and τ_i) lower than 0.15 have been considered. These two parameters were provided as part of the PFS level 2 (L2) products by the National Institute of Astrophysics INAF-IAPS [18, 35, 36]. Then, to guarantee sufficient SNR, only spectra meeting the following three criteria were kept:

- The radiance value at 2165.7 cm^{-1} (corresponding to a CO line) has to be larger than 3 times the value of the noise, which was estimated to be $0.025 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$ based on the standard deviation at 2165 cm^{-1} of an ensemble PFS spectra. This first criterion proves useful to discard PFS spectra with low radiance, but does not inform on the strength of the CO absorption.
- The strength of the CO line at 2165.7 cm^{-1} , calculated in the brightness temperature (BT) space, has to be larger than 2.2 K (the noise equivalent temperature difference is estimated to be 1.1 K at 280 K). This strength is calculated as the difference (ΔBT) between the BT at 2165.7 cm^{-1} and the average brightness temperature of two channels chosen for the baseline (2163.6 cm^{-1} and 2167.8 cm^{-1}).
- The surface temperature from the PFS L2 product has to be larger than 240 K to remove spectra with very low surface contribution.

A total of around 80000 spectra in the 1-0 band, corresponding to 8 % of all PFS observations are passing these selection criteria. Interestingly they cover a complete seasonal cycle and all latitudes, except polar winters. Note that with these first criteria, only daytime observations are selected; night-time observations are associated with lower SNR and have been discarded. Note finally that the possibility to apply less stringent selection criteria to extend the analysis on additional PFS observations will be tested in future studies.

After this first step, the selected PFS observations have been sorted in 5 latitude bands for the 4 seasons, as: the North Pole (60° to 90°), the mid-latitudes of the Northern hemisphere (30° to 60°), the Equator (-30° to 30°), the mid-latitudes of the Southern hemisphere (-60° to -30°) and the South Pole (-90° to -60°). The seasons are defined according to the solar longitude (L_s): Northern spring/Southern Autumn ($0^\circ - 90^\circ$), Northern

289 summer/Southern winter ($90^\circ - 180^\circ$), Northern autumn/Southern spring
 290 ($180^\circ - 270^\circ$) and Northern winter/Southern summer ($270^\circ - 360^\circ$). For
 291 each latitude band and each season, only one spectrum has been selected to
 292 restrict the number of cases. Note that there were no spectra passing the
 293 criteria for the autumn and winter at both poles, characterized by very low
 294 source signals, especially during the polar night. The 16 selected spectra for
 295 this prospective study are listed in Table 1 and shown in Figure 1. Most of
 296 them present a similar radiance of around $0.6\text{--}1.0 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$ in the
 297 CO band; the three with the larger radiances (yellow curves except for the
 298 mid-latitudes of the Northern hemisphere) are associated to scenes with high
 299 surface temperature. All 16 spectra show very clear CO lines as expected
 300 and have corresponding dust and ice integrated opacities significantly smaller
 301 than 0.15. The CO 1–0 band consists of the P and R branches, which are
 302 both well seen in the PFS spectra, respectively in the ranges $2020\text{--}2139 \text{ cm}^{-1}$
 303 and $2147\text{--}2240 \text{ cm}^{-1}$ (see Figure 1a). The spectral range on the longwave side
 304 of the $\text{CO}_2 \nu_3$ band (before 2200 cm^{-1}) contains some other CO_2 absorption
 305 features; the strongest are labelled in Figure 1a.

Table 1: Selected PFS spectra. I_{2165} stands for the value of the radiance at 2165.7 cm^{-1} . ΔBT is the brightness temperature difference between a CO line at 2165.7 cm^{-1} and two baseline channels (see text for details); τ_d and τ_i are the dust and ice integrated opacities [35, 18, 36]; L_s is the solar longitude, T_s is the surface temperature [18, 35, 36], MY is the Martian year, LT is the local time and the longitudes are East.

Lat. band	L_s ($^\circ$)	Latitude ($^\circ$)	Longitude ($^\circ$)	MY	LT (h)	T_s (K)	ΔBT (K)	I_{2165} ($\text{erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$)	τ_d	τ_i
N Pole ($60^\circ\text{--}90^\circ$)	68.07	63.56	53.35	32	15.44	251	3.00	0.548	0.091	0.052
	104.70	72.81	53.45	32	11.85	245	3.22	0.503	0.055	0.026
N mid lat. ($30^\circ\text{--}60^\circ$)	59.66	52.81	-40.13	32	15.97	254	3.69	0.612	0.067	0.013
	135.72	42.09	31.16	30	15.77	257	3.29	0.606	0.003	0.063
	181.22	31.62	8.12	31	9.26	252	4.43	0.750	0.067	0.024
	330.24	30.09	-86.76	31	10.02	245	3.16	0.561	0.033	0.054
Equator ($-30^\circ\text{--}30^\circ$)	65.89	-6.72	-5.32	32	15.25	256	3.61	0.642	0.055	0.096
	132.65	-1.92	58.51	28	10.16	259	3.29	0.790	0.043	0.021
	209.13	-5.69	132.43	30	10.34	271	3.35	1.408	0.066	0.014
	324.27	2.61	92.32	31	10.55	253	3.50	0.784	0.044	0.011
S mid lat. ($-60^\circ\text{--}-30^\circ$)	2.12	-40.05	-119.41	30	14.19	256	3.48	0.749	0.023	0.056
	155.82	-35.69	-155.81	31	11.02	248	2.32	0.530	0.026	0.064
	212.85	-47.88	151.47	30	9.98	264	2.54	0.934	0.125	0.006
	334.42	-45.38	118.75	31	10.26	253	3.60	0.766	0.081	0.049
S Pole ($-90^\circ\text{--}-60^\circ$)	262.90	-62.63	-134.47	31	15.86	275	2.36	1.403	0.129	0.009
	305.11	-76.37	26.78	32	10.41	252	3.77	0.832	0.034	0.019

Figure 1: Selected spectra for each band of latitude; see also Table 1. Panel (b) shows the spectra selected for both North and South poles.

5.2. Input to the radiative transfer

The radiative transfer equation is solved for an atmosphere extending from 0 km to 60 km. For each single PFS spectrum, the temperature profile is fixed to the one retrieved independently by INAF-IAPS from the corresponding spectrum recorded by the LWC (see [35, 36, 37] for details). In the spectral range of the 1–0 band of CO, there are lines of the ν_2 vibrational band of water vapour (H_2O ; band centered at 1595 cm^{-1}) and $\nu_1 + \nu_3$ combination band of ozone (O_3 ; centered at 2105 cm^{-1}) that exist. Although their contribution is close from being negligible, without observable spectral features in the spectral range, it has been taken into account in the radiative

316 transfer calculations by fixing the concentrations of the two species to clima-
 317 tological values. The most adequate value of surface emissivity has also to be
 318 considered, as errors on this parameter will propagate on the retrieved values
 319 of surface temperature and effective reflectivity, and hence ultimately on the
 320 retrieved CO vertical profile. The Planetary SURface Portal [38, 39] provides
 321 global maps of the emissivity of the surface at 2000 cm^{-1} on spatial grids of
 322 $0.025^\circ \times 0.025^\circ$ resolution; they were built from OMEGA/MEX observations
 323 taken over 3 Martian years (MY 27-30) [40]. The emissivity values for the
 324 grid box corresponding to the selected PFS spectra have been considered in
 325 the retrieval. The spectroscopic parameters have been taken from HITRAN
 326 2016 [41], including the pressure broadening due to CO_2 [42].

327 The PFS spectra analysed in this work are apodized by an Hamming func-
 328 tion. We have tested two different formulations of the ILS, which ultimately
 329 were shown to give very similar quality of fits. An example of the residuals
 330 (observed–calculated) obtained after fitting is shown in Figure 2. The first
 331 corresponds to the ILS formulation provided by INAF-IAPS [43] and which
 332 is the recommended one for the community; it is similar to the expected the-
 333 oretical apodized ILS for PFS, except slightly asymmetric, and leads to the
 334 best fit for half of the 16 spectra. The second one (hereafter called "HG")
 335 stems from the theoretical Hamming function, but slightly modified with
 336 an off-axis angle (misalignment in the interferometer), to introduce a small
 337 asymmetry in the function [44], and with second-order auto-apodization ef-
 338 fects. This second one leads to the best fits for the other half of the 16
 339 spectra analysed in this work. For consistency with previous works, all re-
 340 sults presented in the following are those obtained using the ILS provided
 341 by INAF-IAPS. The comparison between the two sets of fits is used later in
 342 section 6.2.3 to discuss the error possibly introduced by an imperfect ILS.

Figure 2: Example of residuals between the observed and fitted spectra calculated for the case using the HG function (blue) and the ILS provided by the INAF-IAPS (red). The RMS are shown for each case.

5.3. Retrievals settings

The retrieval of CO vertical profiles has been performed in two steps. In the first step, a large spectral range from 2040 to 2600 cm^{-1} was adjusted with the objective to obtain an accurate value for the effective surface reflectivity, which can then be used in the second step. The *a priori* value of the reflectivity has been set to 0.023 and its associated variability to 20%. The retrieved reflectivity is fed into the second and main step, which consists in the retrieval of the vertical profile of CO. It is performed on a shorter spectral range, extending from 2040 to 2230 cm^{-1} . Specifically, the CO vertical profiles are retrieved for 2 km thick layers from 0 to 24 km; the rest of the profile is kept fixed. The *a priori* profile $\mathbf{x}_a$ and covariance matrix $\mathbf{S}_a$ used to constrain the CO retrieval have been built from a global ensemble of modelled profiles from the Mars Climate Database (MCD, version 5.3) [7, 45] for one Martian year (climatology solar average scenario). These profiles were

357 simulated on a spatial grid of $3.75^\circ \times 5.625^\circ$ (latitude $\times$ longitude) for one
 358 day per Martian month (defined as a period of $30^\circ L_s$). This day corresponds
 359 to the centered L_s of the month and CO profiles were calculated every 2 Mar-
 360 tian hours (1 Martian hour is defined as $1/24^{th}$ of a sol), where daytime and
 361 nighttime are included. This results in an *a priori* covariance matrix with
 362 large but representative variability, that can be applied without changes for
 363 all observations on Mars. In total, 451584 CO profiles have been considered.
 364 The resulting $\mathbf{x}_a$ profile and the $\mathbf{S}_a$ matrix¹ are shown in Figure 3 for the
 365 2 km thick layers that have been retrieved over the altitude range from 0 to
 366 24 km. The *a priori* profile shows an almost uniform vertical distribution
 367 of CO at a volume mixing ratio (VMR) of about 700 ppm between 5 and
 368 15 km. Close to the surface, the VMR increases to about 750 ppm and from
 369 15 km of height, it starts to increase slightly due to the photodissociation
 370 of CO_2 at higher altitudes. The largest *a priori* variability is at the sur-
 371 face, where it reaches 57%. It then decreases with altitude and drops to a
 372 minimum of 20% in the 12–14 km layer. From this minimum, the *a priori*
 373 variability starts to increase again. Also noteworthy and of high relevance
 374 for this work is that the $\mathbf{S}_a$ presents non-negligible correlations between the
 375 different layers, meaning that the value of the profile retrieved at one given
 376 altitude will be influenced significantly from layers above and below. On a
 377 minor point, note that the $\mathbf{S}_a$ originally built from the model had unphysical
 378 correlation between the surface layer and the very high atmosphere, which
 379 were removed.

380 In the main step of the retrieval, in addition to the CO profile, the sur-
 381 face temperature and the profile of CO_2 have been retrieved simultaneously,
 382 which allows recalculating more accurately the CO VMR at the last itera-
 383 tion. The *a priori* uncertainty on the surface temperature has been set to
 384 20%. The *a priori* vertical profile of CO_2 and its associated variability have
 385 been determined using the model output, in the same way as for CO (see
 386 above). The CO_2 *a priori* profile is almost uniform, with a mole fraction of
 387 around 0.963, and the prior uncertainty is very small, largest close to the
 388 surface (2.2%) and then around 0.5% at higher altitudes. Despite the low
 389 variability allowed on the CO_2 profile, fitting the latter slightly improved the
 390 quality of the fits in most cases. Finally, the $\mathbf{S}_e$ has been considered diagonal

¹The $\mathbf{S}_a$ is expressed in multiplicative factor. $\sqrt{\sigma_a^2} \times 100\%$ represents the variability
 on the *a priori* profile.

391 with a value for σ_ϵ of $0.028 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$, larger than the noise value
 392 mentioned in section 5.1 to take into account forward modelling errors.

Figure 3: A priori profile (left) and covariance matrix (right) built using a representative ensemble of CO profiles from the MCD database (climatology solar average scenario). The matrix is expressed in multiplicative factor. The error bars on the profile correspond to the square root of the diagonal elements of the matrix, applied on the VMR.

393 6. Results

394 6.1. Spectral fittings and retrieved profiles

395 With this choice of retrieval parameters, the 16 spectra are fitted well,
 396 with the RMS of the residual varying between $0.019 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$
 397 and $0.046 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$. Among all spectra, 9 have a RMS lower
 398 than the level of σ_ϵ considered. Figure 4 illustrates the quality of the fits for
 399 the two extreme cases, with respectively the lowest (Figure 4.a) and largest
 400 RMS (Figure 4.b) obtained. For the lowest RMS case, the residual of the fit
 401 lies well within the level of $\pm \sigma_\epsilon$ we have considered. For the highest RMS
 402 case, some spectral features remain above the noise level. This is the case
 403 in the R branch of the CO band, where the fitted spectrum is too high in
 404 radiance, mainly due to the difficulty of reproducing accurately the baseline
 405 of the spectrum, which is affected by the surface emissivity, reflectivity and
 406 temperature but also the CO₂ continuum. It is worth noting that the surface

radiance is significantly higher in this case (around $2 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$ at 2050 cm^{-1} , almost twice larger) than for the lowest RMS case. The increase of the RMS with the radiance level seems to be a general behaviour among the 16 spectra analysed. Overall, we find that the average RMS is very similar to the $\pm \sigma_\epsilon$ value considered in $\mathbf{S}_\epsilon$, suggesting that this choice for the constrain on the measurement is appropriate and in some cases quite conservative.

Figure 4: Example fits among the 16 PFS spectra analysed. For the case (a) with the lowest RMS of the residual ($0.019 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$) and, (b) the largest RMS ($0.046 \text{ erg s}^{-1} \text{ sr}^{-1} \text{ cm}^{-2} \text{ cm}$).

Figure 5 shows the retrieved CO profiles for three representative cases. Above 5 km of height, all retrieved profiles present a similar shape as the *a priori* profile, although not exactly at the same VMR. In the last layer, the maximal difference between the retrieved CO VMRs is equal to 457 ppm, with a minimum and a maximum CO VMRs respectively of 514 and 971 ppm. Closer to the Martian surface, the variability in the retrieved CO VMRs around the *a priori* is larger, with a minimum and a maximum of respectively 313 ppm (spectrum recorded in the South Pole during Southern spring) and 1687 ppm (spectrum recorded in mid-latitudes of the Northern hemisphere during Northern autumn), leading to a maximal difference of 1374 ppm.

424 This departure from the *a priori* tends to show that the sensitivity of PFS
 425 observations is significant close to the surface, and this will be discussed in
 426 more details in the next section.

Figure 5: Examples of retrieved CO profiles. The VMRs correspond to the averaged VMRs of each retrieved layer (every 2 km). The *a priori* profile is shown in blue along with the associated *a priori* variability (square root of the diagonal element of the $\mathbf{S}_a$).

427 The total columns of CO calculated for all spectra vary between 8.83×10^{19}
 428 molecules/cm² and 2.76×10^{20} molecules/cm². These columns are associated
 429 with column-averaged VMRs between 464 ppm and 1115 ppm, which are
 430 shown in Figure 6 as a function of L_s and latitude. The observed variability is
 431 at first sight in agreement with the results of other studies (e.g. [20]). Indeed,
 432 previous measurements have reported that CO presents strong latitudinal
 433 and seasonal variations, especially in the polar regions, where it is mainly due
 434 to variations of the CO₂ surface pressure related to its alternating condensation
 435 and sublimation. In autumn/winter, CO₂ condenses on the polar ice cap,
 436 leading to a decrease of the CO₂ pressure and, as a result, to an increase of
 437 the CO VMR. The reverse happens during spring/summer, when the CO₂
 438 ice sublimates. From Figure 6, the column-averaged CO VMRs retrieved
 439 in the polar regions during spring/summer are generally lower than those

440 retrieved elsewhere, with averaged values around 650 ppm for the North
 441 Pole and 500 ppm for the South Pole. Note again that we did not find PFS
 442 observations with detectable CO lines for the poles during the winter. The
 443 averaged of the CO VMR calculated for all observations is around 750 ppm.
 444 These values are in the range of those retrieved from CRISM observations and
 445 interestingly capture similar broad patterns of spatial and seasonal variability
 446 of CO reported from CRISM [20]. A larger ensemble of PFS observations,
 447 recorded at similar locations and times as CRISM measurements, would be
 448 required to achieve a better and more insightful comparison between the two
 449 satellite datasets.

Figure 6: Column-averaged CO VMRs (color scale, ppm) calculated from the retrieved profiles as a function of latitude and L_s . Note that the PFS observations correspond to different Martian years.

450 6.2. Characterization of the retrieved profiles

451 6.2.1. Vertical sensitivity

452 The vertical sensitivity of the retrieval can be firstly described with the
 453 Jacobians, which provide an estimation of the sensitivity of the TOA radiance
 454 to a change of the CO concentration in a certain layer. The Jacobians for
 455 one out of the 16 spectra is shown as an example in Figure 7 (note that they
 456 are similar for all spectra). As seen from Figure 7, for the entire spectral

457 range, the Jacobians reach their largest value close to the surface, showing
 458 that this is where the nadir PFS observations have the maximal sensitivity
 459 to CO. The values of the Jacobians then decrease with altitude to almost 0
 460 above 15 km, where the sensitivity of PFS measurements to CO is vanishing.
 461 An important result that follows is that the sensitivity is altitude dependent
 462 and that this *a posteriori* justifies the idea of fitting a profile rather than
 463 a column. It is also worth pointing that the PFS sensitivity to CO is not
 464 constant over the whole spectral range used for the retrieval: the instrument
 465 is more sensitive to CO in the strongest R2 to R8 lines (2154–2176 cm^{-1}) for
 466 the R-branch and in the P3 to P8 lines (2111–2131 cm^{-1}) for the P-branch.

Figure 7: Example of Jacobians ($\text{erg s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1} \text{ cm}$ per scaling factor, color scale) calculated for the PFS observation at -5.69° latitude and 132.43° longitude; 209.13° L_s . The corresponding spectrum is shown in pink.

467 The maximal sensitivity that PFS has to CO close to the surface is con-
 468 firmed by the AVK functions, which are shown in Figure 8 for the lowest
 469 (a) and highest (b) DOFS cases. For all retrieved layers, the AVK functions
 470 peak at the surface, with values ranging between 0.36 and 0.44. The width
 471 of the AVK is large, with each function spanning the range from 0 to about
 472 12 km. This shows that all retrieved layers are correlated with each other
 473 and especially that the information in the upper altitudes is mostly driven by
 474 the retrieval at the near-surface. While nadir TIR sounders on Earth have

generally low sensitivity to the surface (e.g. IASI, [26]) this is obviously not the case here. There are two reasons for this: 1) the thermal contrast associated to the 16 selected observations varies between 15 K and 64 K, which is high compared to the usual values encountered on Earth, and even more importantly 2) the contribution of the reflected solar radiation to the upwelling surface radiance is more significant in the case of Mars than for Earth, which makes the dependency to thermal contrast less relevant. Finally, note that above 15 km, all the AVK functions drop close to 0; the CO retrieval is no longer sensitive to the CO abundance in these layers.

The averaging kernels show that the retrieved profile is not resolved vertically. In fact among the 16 PFS observations, the DOFS varies between 0.74 and 0.97. This shows that the only quantity retrieved independently is the total column of CO. The variation of the DOFS is mostly related to that of the signal-to-noise ratio of PFS observations, and hence mainly to surface temperature and reflectivity. The dependence on thermal contrast is second-order and does not affect the values of the DOFS much. There is, however, a positive correlation between the first diagonal element of the AVK matrix (sensitivity of the retrieved CO column in the first layer to the true column in that layer) and the thermal contrast. On the contrary, from the small set of spectra analysed we do not observe a dependence of the vertical sensitivity on the abundance of CO.

Figure 8: AVK functions associated to the retrieval of CO profile for the case with the lowest DOFS (a), and for the case with the highest DOFS (b). The DOFS is indicated on top along with latitude and solar longitude.

6.2.2. Error budget

The different sources of error on the retrieved CO profile have been evaluated from equations 8, 9, 10 and 11 along with the total error. Figure 9 shows the vertical distribution of these errors for two cases: the one with the lowest uncertainty (Figure 9.a) and the one with the highest uncertainty (Figure 9.b). For all 16 spectra, the largest error is found at the surface (where the prior uncertainty is the largest), and varies between 26% and 32%. The retrieval error then decreases with height. Above 5 km, it stays almost constant and varies between 10% and 13%. Overall, at all altitudes, the prior uncertainty is reduced by around 50%, showing the improvement in the knowledge of the CO vertical distribution brought by PFS measurements. The total retrieval error on the total column of CO is between 5 and 15%, which is promising in the perspective to study the CO spatial and temporal variability on Mars.

Figure 9: Vertical distribution of the total error on the retrieved CO profile and of the different sources of uncertainty. Two cases are shown: the one with the lowest total error (a) and the one with the highest error (b). The *a priori* variability is shown as the black line.

As seen from Figure 9, the two main sources of uncertainty are the smoothing and the measurement errors, which have very similar contributions to the total retrieval error. Above 6 km, the smoothing error is generally the largest and varies between 6% and 11%. The measurement error is lower in this altitude range and varies between 5 and 9%. At the surface, the

515 measurement error varies between 12% and 24% depending on the strength
 516 of the radiance, and the smoothing error is around 23%. In Figure 9, we can
 517 see that the contributions of the uncertainties on the CO₂ profile and surface
 518 temperature are below 1%. This small impact is related to the fact that these
 519 two parameters are retrieved along with the CO profile; the errors would in
 520 reality be larger if these parameters - especially surface temperature - were
 521 treated as fixed parameters with a given uncertainty.

522 6.2.3. Other factors affecting the retrievals

523 Here we discuss the impact on the profile retrievals and their character-
 524 ization of two factors: the choice of the $\mathbf{S}_a$ matrix to constrain the profile,
 525 and that of the ILS formulation (see also discussion in section 5.2). In order
 526 to test the *a priori* covariance matrix, we have repeated the CO retrievals
 527 for four observations out of the 16, with an *ad hoc* matrix, showing 50%
 528 variability at all levels (i.e. the diagonal elements are all equal to 0.25) and a
 529 correlation length of 11 km (exponential decay from the diagonal elements).
 530 The results, presented in Figure 10 (a), clearly show spurious oscillations of
 531 the retrieved VMR as compared to the retrievals with the model-derived $\mathbf{S}_a$,
 532 Figure 10 (b). This is a simple but solid illustration of the need to consider
 533 realistic constraints in retrieving vertical profiles from nadir observations.

Figure 10: The retrieved CO profiles on the equator for different season by testing two *a priori* covariance matrices.

534 As mentioned in section 5.2, two different ILS have been found to give

535 similar fits quality of PFS observations. The baseline chosen for this work is
 536 the one recommended by the PI team of PFS (hereafter called "PILS") and
 537 in this section, we evaluate the impact of this choice on the retrieved CO
 538 profile by also evaluating the results obtained with the modified Hamming
 539 function (HG). This is done by calculating, for each observation, the relative
 540 difference (RD) between the retrieved CO profiles using the two ILS models as
 541 $RD = \frac{(CO_{PILS} - CO_{HG}) \times 100}{CO_{PILS}}$. The average relative difference for all the retrieved
 542 layers is shown in Figure 11. It is clear from this figure that the choice of
 543 the ILS function has a small but non negligible effect on the retrieved CO
 544 profile. The CO VMR is systematically higher when using the PILS, with
 545 a maximum relative difference at the surface, where it reaches 9%. It then
 546 decreases to around 3% at higher altitudes. On the total column of CO,
 547 this translates to relative differences around 5% on average. It is useful to
 548 have these differences in mind for future work, as these can be taken as a
 549 systematic uncertainty on the retrieved CO profile and columns.

Figure 11: Averaged relative difference between the CO profiles retrieved with the ILS recommended by the PI team of PFS and the HG ILS adjusted in this work.

550 7. Conclusions

551 A first attempt to retrieve CO vertical profiles from nadir observations
552 on Mars has been performed for the first time. This has been achieved on
553 high spectral resolution measurements made by the PFS instrument on Mars
554 Express, in the spectral region of the 1–0 fundamental band of CO (centered
555 at 2143 cm^{-1}). Despite the fact that only a very small set of spectra with
556 sufficient signal-to-noise ratios has been considered in this exploratory study,
557 our results have clearly demonstrated the potential of using this spectral
558 range for monitoring CO spatially and temporally on single PFS observa-
559 tions, except in the polar winters. The CO profiles have been retrieved using
560 the Optimal Estimation Method, with *a priori* information (*a priori* profile
561 and variance-covariance matrix) built from a large ensemble of global model
562 profiles, which we have shown to provide appropriate constraints. The CO
563 profiles retrieved are very weakly resolved vertically, with the PFS spectra
564 carrying no more than one piece of information. The vertical sensitivity is,
565 however, altitude-dependent (sensitivity from 0 to about 20 km, maximal at
566 the surface) and this alone justifies the retrieval of the profile. The errors on
567 each level of the profile vary between 25% close to the surface to less than
568 10% above 3 km, due to the combined contribution of the measurement noise
569 and the smoothing error. This translates to an error on the retrieved column
570 or the column-averaged VMR of 5 to 15% depending on the scene. Other
571 sources of errors have been identified and we have characterized in particu-
572 lar the one introduced by the uncertainty on the PFS ILS. In all cases we
573 found that the PFS measurements substantially reduce the prior uncertainty
574 on the CO profile and the column, and, as a result, that the retrieved CO
575 abundances allow capturing the spatial and seasonal variability of CO on
576 Mars. We found a preliminary agreement, considering the different kind of
577 datasets that are compared, with CO abundances retrieved from the CRISM
578 spectrometer [20]. Generally, the possibility of retrieving CO with good ac-
579 curacy from PFS single observations in the 1–0 band is very promising and
580 stimulates larger-scale retrievals, which would provide strong constraints on
581 the cycle of carbon species in the lowest (below 15 km) Martian atmosphere.

582 8. Acknowledgements

583 This work is funded by the FNRS CRAMIC project under grant number
584 T.0171.16. PFS is funded by the Italian Space Agency under grant 2018-
585 2-HH.0. Sophie Bauduin is Chargée de Recherches with FRS-FNRS. The

586 authors would like to thank the LMD for providing the access to the Mars
587 Climate Database, needed to build the *a priori* vertical profile CO and the
588 associated covariance matrix.

589 9. References

- 590 [1] L. D. Kaplan, J. Connes, and P. Connes, “Carbon monoxide in the
591 Martian atmosphere,” *The Astrophysical Journal*, vol. 157, pp. L187–
592 L192, 1969.
- 593 [2] M. B. McElroy and T. M. Donahue, “Stability of the Martian Atmo-
594 sphere,” *Science*, vol. 177, pp. 986–988, Sept. 1972.
- 595 [3] T. D. Parkinson and D. M. Hunten, “Spectroscopy and Acronomy of
596 O₂ on Mars,” *Journal of Atmospheric Sciences*, vol. 29, pp. 1380–1390,
597 Oct. 1972.
- 598 [4] S. K. Atreya and Z. G. Gu, “Stability of the Martian atmosphere: Is
599 heterogeneous catalysis essential?,” *Journal of Geophysical Research*,
600 vol. 99, pp. 13133–13145, June 1994.
- 601 [5] S. K. Atreya and Z. G. Gu, “Photochemistry and stability of the atmo-
602 sphere of Mars,” *Advances in Space Research*, vol. 16, Aug. 1995.
- 603 [6] H. Nair, M. Allen, A. D. Anbar, Y. L. Yung, and R. T. Clancy, “A pho-
604 tochemical model of the martian atmosphere,” *Icarus*, vol. 111, pp. 124–
605 150, Sept. 1994.
- 606 [7] F. Forget, F. Hourdin, R. Fournier, C. Hourdin, O. Talagrand,
607 M. Collins, S. R. Lewis, P. L. Read, and J.-P. Huot, “Improved general
608 circulation models of the Martian atmosphere from the surface to above
609 80 km,” *Journal of Geophysical Research: Planets*, vol. 104, no. E10,
610 pp. 24155–24175, 1999.
- 611 [8] L. Neary and F. Daerden, “The GEM-Mars general circulation model
612 for Mars: Description and evaluation,” *Icarus*, vol. 300, pp. 458–476,
613 2018.
- 614 [9] S. K. Atreya and J. E. Blamont, “Stability of the Martian atmosphere -
615 Possible role of heterogeneous chemistry,” *Geophysical Research Letters*,
616 vol. 17, pp. 287–290, Mar. 1990.

- 617 [10] F. Lefèvre, J.-L. Bertaux, R. T. Clancy, T. Encrenaz, K. Fast, F. Forget,
618 S. Lebonnois, F. Montmessin, and S. Perrier, “Heterogeneous chemistry
619 in the atmosphere of Mars,” *Nature*, vol. 454, pp. 971–975, Aug. 2008.
- 620 [11] V. Formisano, S. Atreya, T. Encrenaz, N. Ignatiev, and M. Giuranna,
621 “Detection of Methane in the Atmosphere of Mars,” *Science*, vol. 306,
622 pp. 1758–1761, Dec. 2004.
- 623 [12] K. Zahnle, R. S. Freedman, and D. C. Catling, “Is there methane on
624 Mars?,” *Icarus*, vol. 212, pp. 493–503, Apr. 2011.
- 625 [13] C. R. Webster, P. R. Mahaffy, S. K. Atreya, J. E. Moores, G. J. Flesch,
626 C. Malespin, C. P. McKay, G. Martinez, C. L. Smith, J. Martin-Torres,
627 J. Gomez-Elvira, M.-P. Zorzano, M. H. Wong, M. G. Trainer, A. Steele,
628 D. Archer, B. Sutter, P. J. Coll, C. Freissinet, P.-Y. Meslin, R. V. Gough,
629 C. H. House, A. Pavlov, J. L. Eigenbrode, D. P. Glavin, J. C. Pear-
630 son, D. Keymeulen, L. E. Christensen, S. P. Schwenzer, R. Navarro-
631 Gonzalez, J. Pla-García, S. C. R. Rafkin, Á. Vicente-Retortillo, H. Ka-
632 hanpää, D. Viudez-Moreiras, M. D. Smith, A.-M. Harri, M. Genzer,
633 D. M. Hassler, M. Lemmon, J. Crisp, S. P. Sander, R. W. Zurek, and
634 A. R. Vasavada, “Background levels of methane in Mars’ atmosphere
635 show strong seasonal variations,” *Science*, vol. 360, pp. 1093–1096, June
636 2018.
- 637 [14] V. A. Krasnopolsky, “Spectroscopic mapping of Mars CO mixing ratio:
638 Detection of north-south asymmetry,” *Journal of Geophysical Research*
639 (*Planets*), vol. 108, p. 5010, Feb. 2003.
- 640 [15] T. Encrenaz, T. Fouchet, R. Melchiorri, P. Drossart, B. Gondet,
641 Y. Langevin, J.-P. Bibring, F. Forget, and B. Bézard, “Seasonal vari-
642 ations of the martian CO over Hellas as observed by OMEGA/Mars
643 Express,” *Astronomy & Astrophysics*, vol. 459, pp. 265–270, Nov. 2006.
- 644 [16] F. Billebaud, J. Brillet, E. Lellouch, T. Fouchet, T. Encrenaz, V. Cottini,
645 N. Ignatiev, V. Formisano, M. Giuranna, A. Maturilli, and F. Forget,
646 “Observations of CO in the atmosphere of Mars with PFS onboard Mars
647 Express,” *Planetary and Space Science*, vol. 57, pp. 1446–1457, Oct.
648 2009.

- [17] G. Sindoni, V. Formisano, and A. Geminale, “Observations of water vapour and carbon monoxide in the Martian atmosphere with the SWC of PFS/MEX,” *Planetary and Space Science*, vol. 59, no. 2-3, pp. 149–162, 2011.
- [18] M. Giuranna, P. Wolkenberg, D. Grassi, A. Aronica, S. Aoki, D. Scacabarozzi, B. Saggin, V. Formisano, and the PFS team, “New dataset of atmospheric parameters retrieved by PFS-MEx.” Abstract Scientific Workshop: From Mars Express to ExoMars, 2018.
- [19] M. D. Smith, M. J. Wolff, R. T. Clancy, and S. L. Murchie, “Compact Reconnaissance Imaging Spectrometer observations of water vapor and carbon monoxide,” *Journal of Geophysical Research: Planets*, vol. 114, no. E2, 2009.
- [20] M. D. Smith, F. Daerden, L. Neary, and A. Khayat, “The climatology of carbon monoxide and water vapor on Mars as observed by CRISM and modeled by the GEM-Mars general circulation model,” *Icarus*, vol. 301, pp. 117–131, 2018.
- [21] R. T. Clancy, D. O. Muhleman, and B. M. Jakosky, “Variability of carbon monoxide in the Mars atmosphere,” *Icarus*, vol. 55, no. 2, pp. 282–301, 1983.
- [22] E. Lellouch, J. J. Goldstein, S. W. Bougher, G. Paubert, and J. Rosenqvist, “First absolute wind measurements in the middle atmosphere of Mars,” *The Astrophysical Journal*, vol. 383, pp. 401–406, 1991.
- [23] A. S. Khayat, G. L. Villanueva, M. J. Mumma, and A. T. Tokunaga, “A search for SO₂, H₂S and SO above Tharsis and Syrtis volcanic districts on Mars using ground-based high-resolution submillimeter spectroscopy,” *Icarus*, vol. 253, pp. 130–141, 2015.
- [24] A. S. J. Khayat, G. L. Villanueva, M. J. Mumma, and A. T. Tokunaga, “A deep search for the release of volcanic gases on Mars using ground-based high-resolution infrared and submillimeter spectroscopy: sensitive upper limits for OCS and SO₂,” *Icarus*, vol. 296, pp. 1–14, 2017.
- [25] S. Robert, C. Camy-Peyret, F. Daerden, M. De Mazière, E. De Wachter, L. Neary, S. Vandenbussche, and A. Vandaele, “Two test-cases for synergistic detections in the Martian atmosphere: Carbon monoxide and

- methane,” *Journal of Quantitative Spectroscopy and Radiative Transfer*, vol. 189, pp. 86–104, 2017.
- [26] S. Bauduin, L. Clarisse, M. Theunissen, M. George, D. Hurtmans, C. Clerbaux, and P.-F. Coheur, “IASI’s sensitivity to near-surface carbon monoxide (CO): Theoretical analyses and retrievals on test cases,” *Journal of Quantitative Spectroscopy and Radiative Transfer*, vol. 189, pp. 428–440, 2017.
- [27] C. D. Rodgers, *Inverse methods for atmospheric sounding: theory and practice*, vol. 2. World scientific, 2000.
- [28] V. Formisano, F. Angrilli, G. Arnold, S. Atreya, G. Bianchini, D. Biondi, A. Blanco, M. Blecka, A. Coradini, L. Colangeli, *et al.*, “The planetary Fourier spectrometer (PFS) onboard the European Mars Express mission,” *Planetary and Space Science*, vol. 53, no. 10, pp. 963–974, 2005.
- [29] M. Giuranna, V. Formisano, D. Biondi, A. Ekonomov, S. Fonti, D. Grassi, H. Hirsch, I. Khatuntsev, N. Ignatiev, M. Michalska, *et al.*, “Calibration of the Planetary Fourier Spectrometer short wavelength channel,” *Planetary and Space Science*, vol. 53, no. 10, pp. 975–991, 2005.
- [30] M. Giuranna, V. Formisano, D. Biondi, A. Ekonomov, S. Fonti, D. Grassi, H. Hirsch, I. Khatuntsev, N. Ignatiev, M. Malgoska, *et al.*, “Calibration of the Planetary Fourier Spectrometer long wavelength channel,” *Planetary and Space Science*, vol. 53, no. 10, pp. 993–1007, 2005.
- [31] B. Barret, D. Hurtmans, M. R. Carleer, M. De Mazière, E. Mahieu, and P.-F. Coheur, “Line narrowing effect on the retrieval of HF and HCl vertical profiles from ground-based FTIR measurements,” *Journal of Quantitative Spectroscopy and Radiative Transfer*, vol. 95, no. 4, pp. 499–519, 2005.
- [32] P.-F. Coheur, B. Barret, S. Turquety, D. Hurtmans, J. Hadji-Lazaro, and C. Clerbaux, “Retrieval and characterization of ozone vertical profiles from a thermal infrared nadir sounder,” *Journal of Geophysical Research: Atmospheres*, vol. 110, no. D24, 2005.

- [33] C. Clerbaux, P.-F. Coheur, D. Hurtmans, B. Barret, M. Carleer, R. Colin, K. Semeniuk, J. McConnell, C. Boone, and P. Bernath, “Carbon monoxide distribution from the ACE-FTS solar occultation measurements,” *Geophysical research letters*, vol. 32, no. 16, 2005.
- [34] C. D. Rodgers and B. J. Connor, “Intercomparison of remote sounding instruments,” *Journal of Geophysical Research (Atmospheres)*, vol. 108, p. 4116, Feb. 2003.
- [35] P. Wolkenberg, M. Giuranna, D. Grassi, A. Aronica, S. Aoki, D. Scacabarozzi, and B. Saggin, “Characterization of dust activity on Mars from MY27 to MY32 by PFS-MEX observations,” *Icarus*, vol. 310, pp. 32–47, 2018.
- [36] M. Giuranna and et al., “The current weather and climate of Mars: 12 years of atmospheric monitoring by the Planetary Fourier Spectrometer on Mars Express.” *Icarus*, 2019 (submitted).
- [37] D. Grassi, N. I. Ignatiev, L. V. Zasova, A. Maturilli, V. Formisano, G. A. Bianchini, and M. Giuranna, “Methods for the analysis of data from the Planetary Fourier Spectrometer on the Mars Express Mission,” *Planetary and Space Science*, vol. 53, pp. 1017–1034, Aug. 2005.
- [38] F. Poulet, C. Quantin-Nataf, H. Ballans, K. Dassas, J. Audouard, J. Carter, B. Gondet, L. Lozach, J.-C. Malapert, C. Marmo, *et al.*, “PSUP: A Planetary SURface Portal,” *Planetary and Space Science*, 2017.
- [39] C. Quantin-Nataf, L. Lozac’h, P. Thollot, D. Loizeau, B. Bultel, J. Fernando, P. Allemand, F. Dubuffet, F. Poulet, A. Ody, *et al.*, “MarsSI: Martian surface data processing information system,” *Planetary and Space Science*, vol. 150, pp. 157–170, 2018.
- [40] J. Audouard, F. Poulet, M. Vincendon, J.-P. Bibring, F. Forget, Y. Langevin, and B. Gondet, “Mars surface thermal inertia and heterogeneities from OMEGA/MEX,” *Icarus*, vol. 233, pp. 194–213, 2014.
- [41] I. E. Gordon, L. S. Rothman, C. Hill, R. V. Kochanov, Y. Tan, P. F. Bernath, M. Birk, V. Boudon, A. Campargue, K. V. Chance, B. J. Drouin, J.-M. Flaud, R. R. Gamache, J. T. Hodges, D. Jacquemart,

- 746 V. I. Perevalov, A. Perrin, K. P. Shine, M.-A. H. Smith, J. Tennyson,
747 G. C. Toon, H. Tran, V. G. Tyuterev, A. Barbe, A. G. Császár, V. M.
748 Devi, T. Furtenbacher, J. J. Harrison, J.-M. Hartmann, A. Jolly, T. J.
749 Johnson, T. Karman, I. Kleiner, A. A. Kyuberis, J. Loos, O. M. Lyulin,
750 S. T. Massie, S. N. Mikhailenko, N. Moazzen-Ahmadi, H. S. P. Müller,
751 O. V. Naumenko, A. V. Nikitin, O. L. Polyansky, M. Rey, M. Rotger,
752 S. W. Sharpe, K. Sung, E. Starikova, S. A. Tashkun, J. V. Auwera,
753 G. Wagner, J. Wilzewski, P. Wcisło, S. Yu, and E. J. Zak, “The HI-
754 TRAN2016 molecular spectroscopic database,” *Journal of Quantitative*
755 *Spectroscopy and Radiative Transfer*, vol. 203, pp. 3–69, 2017.
- 756 [42] G. Li, I. E. Gordon, L. S. Rothman, Y. Tan, S.-M. Hu, S. Kassi, A. Cam-
757 pargue, and E. S. Medvedev, “Rovibrational Line Lists for Nine Isotopo-
758 logues of the CO Molecule in the X $^1\Sigma^+$ Ground Electronic State,” *The*
759 *Astrophysical Journal Supplement Series*, vol. 216, p. 15, Jan. 2015.
- 760 [43] M. Giuranna, S. Viscardy, F. Daerden, L. Neary, G. Etiope, D. Oehler,
761 V. Formisano, A. Aronica, P. Wolkenberg, S. Aoki, A. Cardesn-Moinelo,
762 J. Marn-Yaseli de la Parra, D. Merritt, and M. Amoroso, “Independent
763 confirmation of a methane spike on Mars and a source region east of
764 Gale Crater,” *Nature Geoscience*, 2019.
- 765 [44] J. Kauppinen and P. Saarinen, “Line-shape distortions in misaligned
766 cube corner interferometers,” *Applied Optics*, vol. 31, pp. 69–74, 1992.
- 767 [45] E. Millour, F. Forget, A. Spiga, M. Vals, V. Zakharov, L. Montabone,
768 F. Lefèvre, F. Montmessin, J.-Y. Chaufray, M. A. López-Valverde,
769 F. González-Galindo, S. R. Lewis, P. L. Read, M. C. Desjean, F. Cipri-
770 ani, and the MCD development team, “The Mars climate database
771 (version 5.3),” in *Scientific Workshop: From Mars Express to ExoMars*
772 *(ESA-ESAC)*, 2018.