

REFERENCES (INTRODUCTION AND DISCUSSION)

1. Zeanah CH. Adaptation following perinatal loss: a critical review. *J Am Acad Child Adolesc Psychiatry*. 1989 Jul;28(4):467–80.
2. McCoid JLM. Pregnancy interrupted: loss of a desired pregnancy after diagnosis of fetal anomaly. *J Psychosom Obstet Gynaecol*. 2007 Mar;28(1):37–48.
3. Obi SN, Onah HE, Okafor II. Depression among Nigerian women following pregnancy loss. *Int J Gynaecol Obstet*. 2009 Apr;105(1):60–2.
4. Boyd PA, Tondi F, Hicks NR, Chamberlain PF. Autopsy after termination of pregnancy for fetal anomaly: retrospective cohort study. *BMJ*. 2004 Jan;328(7432):137.
5. World Health Organisation. WHO | Neonatal and perinatal mortality 2000 [Internet]. WHO. 2006. Available from: https://www.who.int/maternal_child_adolescent/documents/9241563206/en/.
6. Hobbins JC, Grannum PA, Berkowitz RL, Silverman R, Mahoney MJ. Ultrasound in the diagnosis of congenital anomalies. *Am J Obstet Gynecol*. 1979 Jun;134(3):331–45.
7. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 101: Ultrasonography in pregnancy. *Obstet Gynecol*. 2009 Feb;113(2 Pt 1):451–61.
8. Salomon LJ, Alfirevic Z, Bilardo CM, Chalouhi GE, Ghi T, Kagan KO, Lau TK, Papageorghiou AT, Raine-Fenning NJ, Stirnemann J, Suresh S, Tabor A, Timor-Tritsch IE, Toi A, Yeo G. ISUOG practice guidelines: performance of first-trimester fetal ultrasound scan. *Ultrasound Obstet Gynecol*. 2013 Jan;41(1):102–13.

9. International Society of Ultrasound in Obstetrics and Gynecology, Carvalho JS, Allan LD, Chaoui R, Copel JA, DeVore GR, Hecher K, Lee W, Munoz H, Paladini D, Tuschek B, Yagel S. ISUOG Practice Guidelines (updated): sonographic screening examination of the fetal heart. *Ultrasound Obstet Gynecol.* 2013 Mar;41(3):348–59.
10. Donald I, Macvicar J, Brown TG. Investigation of abdominal masses by pulsed ultrasound. *Lancet.* 1958 Jun;271(7032):1188–95.
11. Yagel S, Cohen SM, Messing B, Valsky DV. Three-dimensional and four-dimensional ultrasound applications in fetal medicine. *Curr Opin Obstet Gynecol.* 2009 Apr;21(2):167–74.
12. Grandjean H, Larroque D, Levi S. The performance of routine ultrasonographic screening of pregnancies in the Eurofetus Study. *Am J Obstet Gynecol.* 1999 Aug;181(2):446–54.
13. Levi S. Ultrasound in prenatal diagnosis: polemics around routine ultrasound screening for second trimester fetal malformations. *Prenat Diagn.* 2002 Apr;22(4):285–95.
14. Clementi M, Stoll C. The Euroscan study. *Ultrasound Obstet Gynecol.* 2001 Oct;18(4):297–300.
15. Stoll C, Garne E, Clementi M, EUROSCAN Study Group. Evaluation of prenatal diagnosis of associated congenital heart diseases by fetal ultrasonographic examination in Europe. *Prenat Diagn.* 2001 Apr;21(4):243–52.
16. Carvalho JS. Fetal heart scanning in the first trimester. *Prenat Diagn.* 2004 Dec;24(13):1060–7.
17. Kagan KO, Etchegaray A, Zhou Y, Wright D, Nicolaides KH. Prospective validation of first-trimester combined screening for trisomy 21. *Ultrasound Obstet Gynecol.* 2009 Jul;34(1):14–8.

18. Chaoui R, Benoit B, Mitkowska-Wozniak H, Heling KS, Nicolaides KH. Assessment of intracranial translucency (IT) in the detection of spina bifida at the 11-13-week scan. *Ultrasound Obstet Gynecol.* 2009 Sep;34(3):249–52.
19. Nicolaides KH. Screening for fetal aneuploidies at 11 to 13 weeks. *Prenat Diagn.* 2011 Jan;31(1):7–15.
20. Syngelaki A, Chelemen T, Dagklis T, Allan L, Nicolaides KH. Challenges in the diagnosis of fetal non-chromosomal abnormalities at 11-13 weeks. *Prenat Diagn.* 2011 Jan;31(1):90–102.
21. Vintzileos AM, Ananth CV, Smulian JC, Beazoglou T, Knuppel RA. Routine second-trimester ultrasonography in the United States: A cost-benefit analysis. *Am J Obstet Gynecol.* 2000 Mar;182(3):655–60.
22. Vogt C, Blaas H-GK, Salvesen KÅ, Eik-Nes SH. Comparison between prenatal ultrasound and postmortem findings in fetuses and infants with developmental anomalies. *Ultrasound Obstet Gynecol.* 2012 Jun;39(6):666–72.
23. Smith FW, Adam AH, Phillips WDP. NMR imaging in pregnancy. *Lancet.* 1983 Jan;321(8314):61–2.
24. Bahado-Singh RO, Goncalves LF. Techniques, terminology, and indications for MRI in pregnancy. *Semin Perinatol.* 2013 Oct;37(5):334–9.
25. Whitby EH, Paley MNJ, Sprigg A, Rutter S, Davies NP, Wilkinson ID, Griffiths PD. Comparison of ultrasound and magnetic resonance imaging in 100 singleton pregnancies with suspected brain abnormalities. *BJOG.* 2004 Aug;111(8):784–92.
26. Levine D, Barnes PD, Robertson RR, Wong G, Mehta TS. Fast MR imaging of fetal central nervous system abnormalities. *Radiology.* 2003 Oct;229(1):51–61.
27. Griffiths PD, Porteous M, Mason G, Russell S, Morris J, Fanou EM, Reeves MJ. The use of in utero MRI to supplement ultrasound in the foetus at high risk of developmental brain or spine abnormality. *Br J Radiol.* 2012 Nov;85(1019):e1038-1045.

28. Levine D. Timing of MRI in pregnancy, repeat exams, access, and physician qualifications. *Semin Perinatol.* 2013 Oct;37(5):340–4.
29. Garel C, Chantrel E, Elmaleh M, Brisse H, Sebag G. Fetal MRI: normal gestational landmarks for cerebral biometry, gyration and myelination. *Childs Nerv Syst ChNS.* 2003 Aug;19(7–8):422–5.
30. Malinge G, Ben-Sira L, Lev D, Ben-Aroya Z, Kidron D, Lerman-Sagie T. Fetal brain imaging: a comparison between magnetic resonance imaging and dedicated neurosonography. *Ultrasound Obstet Gynecol.* 2004 Apr;23(4):333–40.
31. Baker PN, Johnson IR, Harvey PR, Gowland PA, Mansfield P. A three-year follow-up of children imaged in utero with echo-planar magnetic resonance. *Am J Obstet Gynecol.* 1994 Jan;170(1 Pt 1):32–3.
32. Clements H, Duncan KR, Fielding K, Gowland PA, Johnson IR, Baker PN. Infants exposed to MRI in utero have a normal paediatric assessment at 9 months of age. *Br J Radiol.* 2000 Feb;73(866):190–4.
33. Kanal E, Borgstede JP, Barkovich AJ, Bell C, Bradley WG, Etheridge S, Felmlee JP, Froelich JW, Hayden J, Kaminski EM, Lester JW Jr, Scoumis EA, Zaremba LA, Zinninger MD; American College of Radiology. American College of Radiology White Paper on MR Safety: 2004 update and revisions. *AJR.* 2004 May;182(5):1111–4.
34. Novak Z, Thurmond AS, Ross PL, Jones MK, Thornburg KL, Katzberg RW. Gadolinium-DTPA transplacental transfer and distribution in fetal tissue in rabbits. *Invest Radiol.* 1993 Sep;28(9):828–30.
35. Sundgren PC, Leander P. Is administration of gadolinium-based contrast media to pregnant women and small children justified? *J Magn Reson Imaging.* 2011 Oct;34(4):750–7.

36. Rofsky NM, Pizzarelli DJ, Weinreb JC, Ambrosino MM, Rosenberg C. Effect on fetal mouse development of exposure to MR imaging and gadopentetate dimeglumine. *J Magn Reson Imaging*. 1994 Dec;4(6):805–7.
37. Rofsky NM, Pizzarelli DJ, Duhaney MO, Falick AK, Prendergast N, Weinreb JC. Effect of magnetic resonance exposure combined with gadopentetate dimeglumine on chromosomes in animal specimens. *Acad Radiol*. 1995 Jun;2(6):492–6.
38. De Santis M, Straface G, Cavaliere AF, Carducci B, Caruso A. Gadolinium periconceptional exposure: pregnancy and neonatal outcome. *Acta Obstet Gynecol Scand*. 2007;86(1):99–101.
39. Bulas D, Egloff A. Benefits and risks of MRI in pregnancy. *Semin Perinatol*. 2013 Oct;37(5):301–4.
40. Votino C, Jani J, Damry N, Dessy H, Kang X, Cos T, Divano L, Foulon W, De Mey J, Cannie M. Magnetic resonance imaging in the normal fetal heart and in congenital heart disease. *Ultrasound Obstet Gynecol*. 2012 Mar;39(3):322–9.
41. Cheung SW, Shaw CA, Yu W, Li J, Ou Z, Patel A, Yatsenko SA, Cooper ML, Furman P, Stankiewicz P, Lupski JR, Chinault AC, Beaudet AL. Development and validation of a CGH microarray for clinical cytogenetic diagnosis. *Genet Med*. 2005 Aug;7(6):422–32.
42. Armengol L, Nevado J, Serra-Juhé C, Plaja A, Mediano C, García-Santiago FA, Garcia-Aragonés M, Villa O, Mansilla E, Preciado C, Fernández L, Angeles Mori M, Garcia-Pérez L, Lapunzina PD, Pérez-Jurado LA. Clinical utility of chromosomal microarray analysis in invasive prenatal diagnosis. *Hum Genet*. 2012 Mar;131(3):513–23.
43. Rooryck C, Toutain J, Cailley D, Bouron J, Horovitz J, Lacombe D, Arveiler B, Saura R. Prenatal diagnosis using array-CGH: a French experience. *Eur J Med Genet*. 2013 Jul;56(7):341–5.

44. Schmid M, Stary S, Springer S, Bettelheim D, Husslein P, Streubel B. Prenatal microarray analysis as second-tier diagnostic test: single-center prospective study. *Ultrasound Obstet Gynecol.* 2013 Mar;41(3):267–73.
45. Wapner RJ, Martin CL, Levy B, Ballif BC, Eng CM, Zachary JM, Savage M, Platt LD, Saltzman D, Grobman WA, Klugman S, Scholl T, Simpson JL, McCall K, Aggarwal VS, Bunke B, Nahum O, Patel A, Lamb AN, Thom EA, Beaudet AL, Ledbetter DH, Shaffer LG, Jackson L. Chromosomal Microarray versus Karyotyping for Prenatal Diagnosis. *N Engl J Med.* 2012 Dec;367(23):2175–84.
46. Hillman SC, McMullan DJ, Hall G, Togneri FS, James N, Maher EJ, Meller CH, Williams D, Wapner RJ, Maher ER, Kilby MD. Use of prenatal chromosomal microarray: prospective cohort study and systematic review and meta-analysis. *Ultrasound Obstet Gynecol.* 2013 Jun;41(6):610–20.
47. Callaway JLA, Shaffer LG, Chitty LS, Rosenfeld JA, Crolla JA. The clinical utility of microarray technologies applied to prenatal cytogenetics in the presence of a normal conventional karyotype: a review of the literature. *Prenat Diagn.* 2013 Dec;33(12):1119–23.
48. Novelli A, Grati FR, Ballarati L, Bernardini L, Bizzoco D, Camurri L, Casalone R, Cardarelli L, Cavalli P, Ciccone R, Clementi M, Dalprà L, Gentile M, Gelli G, Grammatico P, Malacarne M, Nardone AM, Pecile V, Simoni G, Zuffardi O, Giardino D. Microarray application in prenatal diagnosis: a position statement from the cytogenetics working group of the Italian Society of Human Genetics (SIGU), November 2011. *Ultrasound Obstet Gynecol.* 2012 Apr;39(4):384–8.
49. American College of Obstetricians and Gynecologists Committee on Genetics. Committee Opinion No. 581: the use of chromosomal microarray analysis in prenatal diagnosis. *Obstet Gynecol.* 2013 Dec;122(6):1374–7.
50. Bove KE. Practice guidelines for autopsy pathology: the perinatal and pediatric autopsy. Autopsy Committee of the College of American Pathologists. *Arch Pathol Lab Med.* 1997 Apr;121(4):368–76.

51. Burton JL, Underwood J. Clinical, educational, and epidemiological value of autopsy. *Lancet*. 2007 Apr;369(9571):1471–80.
52. Saller DN, Lesser KB, Harrel U, Rogers BB, Oyer CE. The clinical utility of the perinatal autopsy. *JAMA*. 1995 Feb;273(8):663–5.
53. Gordijn SJ, Erwich JJHM, Khong TY. Value of the perinatal autopsy: critique. *Pediatr Dev Pathol*. 2002 Oct;5(5):480–8.
54. Dickinson JE, Prime DK, Charles AK. The role of autopsy following pregnancy termination for fetal abnormality. *Aust N Z J Obstet Gynaecol*. 2007 Dec;47(6):445–9.
55. Sankar VH, Phadke SR. Clinical utility of fetal autopsy and comparison with prenatal ultrasound findings. *J Perinatol*. 2006 Apr;26(4):224–9.
56. Vujanić GM, Cartlidge PH, Stewart JH, Dawson AJ. Perinatal and infant postmortem examinations: how well are we doing? *J Clin Pathol*. 1995 Nov;48(11):998–1001.
57. Bjugn R, Berland J. Quality of fetal, perinatal and infant autopsy reports. An audit of all reports of postmortem examinations following fetal, perinatal and infant death in Rogaland County, Western Norway, 1997-1999. *APMIS*. 2002 Oct;110(10):746–52.
58. Vujanić GM, Cartlidge PH, Stewart JH. Improving the quality of perinatal and infant necropsy examinations: a follow up study. *J Clin Pathol*. 1998 Nov;51(11):850–3.
59. Thayyil S, Chitty LS, Robertson NJ, Taylor AM, Sebire NJ. Minimally invasive fetal postmortem examination using magnetic resonance imaging and computerised tomography: current evidence and practical issues. *Prenat Diagn*. 2010 Aug;30(8):713–8.
60. Heazell AEP, McLaughlin M-J, Schmidt EB, Cox P, Flenady V, Khong TY, Downe S. A difficult conversation? The views and experiences of parents and

- professionals on the consent process for perinatal postmortem after stillbirth. BJOG. 2012 Jul;119(8):987–97.
61. Rose C, Evans M, Tooley J. Falling rates of perinatal postmortem examination: are we to blame? Arch Dis Child Fetal Neonatal Ed. 2006 Nov;91(6):F465.
 62. Tan GC, Hayati AR, Khong TY. Low perinatal autopsy rate in Malaysia: time for a change. Pediatr Dev Pathol. 2010 Oct;13(5):362–8.
 63. Li D-Z, Yu M. Autopsy after termination of pregnancy for fetal anomaly in Mainland China. Prenat Diagn. 2007 Dec;27(12):1183–4.
 64. Khong TY, Tanner AR. Foetal and neonatal autopsy rates and use of tissue for research: the influence of “organ retention” controversy and new consent process. J Paediatr Child Health. 2006 Jun;42(6):366–9.
 65. Cannie M, Votino C, Moerman P, Vanheste R, Segers V, Van Berkel K, Hanssens M, Kang X, Cos T, Kir M, Balepa L, Divano L, Foulon W, De Mey J, Jani J. Acceptance, reliability and confidence of diagnosis of fetal and neonatal virtuopsy compared with conventional autopsy: a prospective study. Ultrasound Obstet Gynecol. 2012 Jun;39(6):659–65.
 66. Bishop KL, Dupuis C, Nanton P, Clarke K, Bolt C, Chin-See C. Perinatal autopsy rates at the University Hospital of the West Indies: 2002-2008. West Indian Med J. 2013 Jan;62(1):35–8.
 67. Breeze ACG, Statham H, Hackett GA, Jessop FA, Lees CC. Perinatal postmortems: what is important to parents and how do they decide? Birth. 2012 Mar;39(1):57–64.
 68. Ben-Sasi K, Chitty LS, Franck LS, Thayyil S, Judge-Kronis L, Taylor AM, Sebire NJ. Acceptability of a minimally invasive perinatal/paediatric autopsy: healthcare professionals' views and implications for practice. Prenat Diagn. 2013 Apr;33(4):307–12.

69. Bourlière-Najean B, Russel AS, Panuel M, Piercecchi-Marti MD, Sigaudy S, Fredouille C, Petit P, Philip N, Devred P. Value of fetal skeletal radiographs in the diagnosis of fetal death. *Eur Radiol.* 2003 May;13(5):1046–9.
70. Stempflé N, Huten Y, Fondacci C, Lang T, Hassan M, Nessmann C. Fetal bone age revisited: proposal of a new radiographic score. *Pediatr Radiol.* 1995;25(7):551–5.
71. Brookes JA, Hall-Craggs MA, Sams VR, Lees WR. Non-invasive perinatal necropsy by magnetic resonance imaging. *Lancet.* 1996 Oct;348(9035):1139–41.
72. Ros PR, Li KC, Vo P, Baer H, Staab EV. Preautopsy magnetic resonance imaging: initial experience. *Magn Reson Imaging.* 1990;8(3):303–8.
73. Roberts MD, Lange RC, McCarthy SM. Fetal anatomy with magnetic resonance imaging. *Magn Reson Imaging.* 1995 Jan 1;13(4):645–9.
74. Deng J, Brookes JA, Gardener JE, Rodeck CH, Lees WR. Three-dimensional magnetic resonance imaging of the postmortem fetal heart. *Fetal Diagn Ther.* 1996 Dec;11(6):417–21.
75. Woodward PJ, Sohaey R, Harris DP, Jackson GM, Klatt EC, Alexander AL, Kennedy A. Postmortem fetal MR imaging: comparison with findings at autopsy. *AJR.* 1997 Jan;168(1):41–6.
76. Langer B, Choquet P, Ravier S, Gasser B, Schlaeder G, Constantinesco A. Low-field dedicated magnetic resonance imaging: a potential tool for assisting perinatal autopsy. *Ultrasound Obstet Gynecol.* 1998 Oct;12(4):271–5.
77. Brookes JA, Deng J, Wilkinson ID, Lees WR. Three-dimensional imaging of the postmortem fetus by MRI: early experience. *Fetal Diagn Ther.* 1999 Jun;14(3):166–71.
78. Breeze ACG, Gallagher FA, Lomas DJ, Smith GCS, Lees CC, Cambridge Post-Mortem MRI Study Group. Postmortem fetal organ volumetry using magnetic

- resonance imaging and comparison to organ weights at conventional autopsy. *Ultrasound Obstet Gynecol.* 2008 Feb;31(2):187–93.
79. Thayyil S, Schievano S, Robertson NJ, Jones R, Chitty LS, Sebire NJ, Taylor AM; MaRIAS (Magnetic Resonance Imaging Autopsy Study) Collaborative group. A semi-automated method for non-invasive internal organ weight estimation by post-mortem magnetic resonance imaging in fetuses, newborns and children. *Eur J Radiol.* 2009 Nov;72(2):321–6.
80. Votino C, Verhoye M, Segers V, Cannie M, Bessieres B, Cos T, Lipombi D, Jani J. Fetal organ weight estimation by postmortem high-field magnetic resonance imaging before 20 weeks' gestation. *Ultrasound Obstet Gynecol.* 2012 Jun;39(6):673–8.
81. Schievano S, Sebire NJ, Robertson NJ, Taylor AM, Thayyil S. Reconstruction of fetal and infant anatomy using rapid prototyping of post-mortem MR images. *Insights Imaging.* 2010 Sep;1(4):281–6.
82. Huisman TAGM, Wisser J, Stallmach T, Krestin GP, Huch R, Kubik-Huch RA. MR autopsy in fetuses. *Fetal Diagn Ther.* 2002 Feb;17(1):58–64.
83. Alderliesten ME, Peringa J, van der Hulst VPM, Blaauwgeers HLG, van Lith JMM. Perinatal mortality: clinical value of postmortem magnetic resonance imaging compared with autopsy in routine obstetric practice. *BJOG.* 2003 Apr;110(4):378–82.
84. Griffiths PD, Variend D, Evans M, Jones A, Wilkinson ID, Paley MNJ, Whitby E. Postmortem MR imaging of the fetal and stillborn central nervous system. *AJNR.* 2003 Jan;24(1):22–7.
85. Breeze ACG, Cross JJ, Hackett GA, Jessop FA, Joubert I, Lomas DJ, Set PA, Whitehead AL, Lees CC. Use of a confidence scale in reporting postmortem fetal magnetic resonance imaging. *Ultrasound Obstet Gynecol.* 2006 Dec;28(7):918–24.

86. Widjaja E, Whitby EH, Cohen M, Paley MNJ, Griffiths PD. Post-mortem MRI of the foetal spine and spinal cord. *Clin Radiol.* 2006 Aug;61(8):679–85.
87. Hagmann CF, Robertson NJ, Sams VR, Brookes J a. S. Postmortem magnetic resonance imaging as an adjunct to perinatal autopsy for renal-tract abnormalities. *Arch Dis Child Fetal Neonatal Ed.* 2007 May;92(3):F215-218.
88. Cohen MC, Paley MN, Griffiths PD, Whitby EH. Less invasive autopsy: benefits and limitations of the use of magnetic resonance imaging in the perinatal postmortem. *Pediatr Dev Pathol.* 2008 Feb;11(1):1–9.
89. Carroll S, Porter H, Abdel-Fattah S, Kyle P, Sotthill P. Correlation of prenatal ultrasound diagnosis and pathologic findings in fetal brain abnormalities. *Ultrasound Obstet Gynecol.* 2000 Aug;16(2):149–53.
90. Thayyil S, Sebire NJ, Chitty LS, Wade A, Chong W, Olsen O, Gunny RS, Offiah AC, Owens CM, Saunders DE, Scott RJ, Jones R, Norman W, Addison S, Bainbridge A, Cady EB, Vita ED, Robertson NJ, Taylor AM; MARIAS collaborative group. Post-mortem MRI versus conventional autopsy in fetuses and children: a prospective validation study. *Lancet.* 2013 Jul;382(9888):223–33.
91. Thayyil S, Cleary JO, Sebire NJ, Scott RJ, Chong K, Gunny R, Owens CM, Olsen OE, Offiah AC, Parks HG, Chitty LS, Price AN, Yousry TA, Robertson NJ, Lythgoe MF, Taylor AM. Post-mortem examination of human fetuses: a comparison of whole-body high-field MRI at 9.4 T with conventional MRI and invasive autopsy. *Lancet.* 2009 Aug;374(9688):467–75.
92. Votino C, Jani J, Verhoye M, Bessieres B, Fierens Y, Segers V, Vorsselmans A, Kang X, Cos T, Foulon W, De Mey J, Cannie M. Postmortem examination of human fetal hearts at or below 20 weeks' gestation: a comparison of high-field MRI at 9.4 T with lower-field MRI magnets and stereomicroscopic autopsy. *Ultrasound Obstet Gynecol.* 2012 Oct;40(4):437–44.

93. Verhoye M, Votino C, Cannie MM, Segers V, Mabiglia C, Cos T, Lipombi D, Jani JC. Post-mortem high-field magnetic resonance imaging: effect of various factors. *J Matern-Fetal Neonatal Med.* 2013 Jul;26(11):1060–5.
94. Thayyil S, Chandrasekaran M, Chitty LS, Wade A, Skordis-Worrall J, Bennett-Britton I, Cohen M, Withby E, Sebire NJ, Robertson NJ, Taylor AM. Diagnostic accuracy of post-mortem magnetic resonance imaging in fetuses, children and adults: a systematic review. *Eur J Radiol.* 2010 Jul;75(1):e142-148.
95. Votino C, Cannie M, Segers V, Dobrescu O, Dessy H, Gallo V, Cos T, Damry N, Jani J. Virtual autopsy by computed tomographic angiography of the fetal heart: a feasibility study. *Ultrasound Obstet Gynecol.* 2012 Jun;39(6):679–84.
96. Breeze ACG, Jessop FA, Whitehead AL, Set PAK, Berman L, Hackett GA, Lees CC, Cambridge post-mortem MRI Study Group. Feasibility of percutaneous organ biopsy as part of a minimally invasive perinatal autopsy. *Virchows Arch Int J Pathol.* 2008 Feb;452(2):201–7.
97. Breeze ACG, Jessop FA, Set PAK, Whitehead AL, Cross JJ, Lomas DJ, Hackett GA, Joubert I, Lees CC. Minimally-invasive fetal autopsy using magnetic resonance imaging and percutaneous organ biopsies: clinical value and comparison to conventional autopsy. *Ultrasound Obstet Gynecol.* 2011 Mar;37(3):317–23.
98. Sebire NJ, Weber MA, Thayyil S, Mushtaq I, Taylor A, Chitty LS. Minimally invasive perinatal autopsies using magnetic resonance imaging and endoscopic postmortem examination (“keyhole autopsy”): feasibility and initial experience. *J Matern-Fetal Neonatal Med.* 2012 May;25(5):513–8.
99. Kang X, Cos Sanchez T, Arthurs OJ, Bevilacqua E, Cannie MM, Segers V, Lecomte S, Sebire NJ, Jani JC. Postmortem fetal imaging: a prospective blinded comparison study of 2-dimensional ultrasound with MR imaging. *Ultrasound Obstet Gynecol.* 2019 Dec;54(6):790-8.
100. Arthurs OJ, Guy A, Thayyil S, Wade A, Jones R, Norman W, Scott R, Robertson NJ, Jacques TS, Chong WK, Gunny R, Saunders D, Olsen OE, Owens CM, Offiah

- AC, Chitty LS, Taylor AM, Sebire NJ; Magnetic Resonance Imaging Autopsy Study (MaRIAS) Collaborative Group. Comparison of diagnostic performance for perinatal and paediatric post-mortem imaging: CT versus MRI. *Eur Radiol*. 2016 Jul;26(7):2327–36.
101. Kang X, Cannie MM, Arthurs OJ, Segers V, Fourneau C, Bevilacqua E, Cos Sanchez T, Sebire NJ, Jani JC. Post-mortem whole-body magnetic resonance imaging of human fetuses: a comparison of 3-T vs. 1.5-T MR imaging with classical autopsy. *Eur Radiol*. 2017 Aug;27(8):3542-53.
 102. Ashwin C, Hutchinson JC, Kang X, Langan D, Jones R, Norman W, Cannie M, Jani J, Sebire NJ, Arthurs OJ. Learning effect on perinatal post-mortem magnetic resonance imaging reporting: single reporter diagnostic accuracy of 200 cases. *Prenat Diagn*. 2017 Jun;37(6):566–74.
 103. Sandaite I, Dymarkowski S, De Catte L, Moerman P, Gewillig M, Fedele L, Deprest J, Claus F. Fetal heart pathology on postmortem 3-T magnetic resonance imaging. *Prenat Diagn*. 2014 Mar;34(3):223–9.
 104. Arthurs OJ, Thayyil S, Addison S, Wade A, Jones R, Norman W, Scott R, Robertson NJ, Chitty LS, Taylor AM, Sebire NJ, Offiah AC; Magnetic Resonance Imaging Autopsy Study (MaRIAS) Collaborative Group. Diagnostic accuracy of postmortem MRI for musculoskeletal abnormalities in fetuses and children. *Prenat Diagn*. 2014 Dec;34(13):1254–61.
 105. Taylor AM, Sebire NJ, Ashworth MT, Schievano S, Scott RJ, Wade A, Chitty LS, Robertson N, Thayyil S; Magnetic Resonance Imaging Autopsy Study Collaborative Group. Post-Mortem Cardiovascular Magnetic Resonance Imaging in Fetuses and Children: A Masked Comparison Study with Conventional Autopsy. *Circulation*. 2014 May;129(19):1937-44.
 106. Arthurs OJ, Thayyil S, Olsen OE, Addison S, Wade A, Jones R, Norman W, Scott RJ, Robertson NJ, Taylor AM, Chitty LS, Sebire NJ, Owens CM; Magnetic Resonance Imaging Autopsy Study (MaRIAS) Collaborative Group. Diagnostic

- accuracy of post-mortem MRI for thoracic abnormalities in fetuses and children. *Eur Radiol*. 2014 Nov;24(11):2876–84.
107. Arthurs OJ, Thayyil S, Pauliah SS, Jacques TS, Chong WK, Gunny R, Saunders D, Addison S, Lally P, Cady E, Jones R, Normal W, Scott R, Robertson NJ, Wade A, Chitty L, Taylor AM, Sebire NJ; Magnetic Resonance Imaging Autopsy Study (MaRIAS) Collaborative Group. Diagnostic accuracy and limitations of post-mortem MRI for neurological abnormalities in fetuses and children. *Clin Radiol*. 2015 Aug;70(8):872–80.
108. Arthurs OJ, Thayyil S, Owens CM, Olsen OE, Wade A, Addison S, Jones R, Norman W, Scott RJ, Robertson NJ, Taylor AM, Chitty LS, Sebire NJ. Diagnostic accuracy of post mortem MRI for abdominal abnormalities in foetuses and children. *Eur J Radiol*. 2015 Mar;84(3):474–81.
109. Vullo A, Panebianco V, Cannavale G, Aromatario M, Cipolloni L, Frati P, Santurro A, Vullo F, Catalano C, Fineschi V.. Post-mortem magnetic resonance foetal imaging: a study of morphological correlation with conventional autopsy and histopathological findings. *Radiol Med*. 2016 Nov;121(11):847–56.
110. Shruthi M, Gupta N, Jana M, Mridha AR, Kumar A, Agarwal R, Sharma R, Deka D, Gupta AK, Kabra M. Conventional vs virtual autopsy with postmortem MRI in phenotypic characterization of stillbirths and fetal malformations. *Ultrasound Obstet Gynecol*. 2018 Feb;51(2):236–45.
111. Staicu A, Albu C, Popa-Stanila R, Chiriac L, Boitor-Borza D, Bondor C, et al. Potential clinical benefits and limitations of fetal virtopsy using high-field MRI at 7 Tesla versus stereomicroscopic autopsy to assess first trimester fetuses. *Prenat Diagn*. 2019 Jun;39(7):505–18.
112. Shelmerdine SC, Arthurs OJ, Gilpin I, Norman W, Jones R, Taylor AM, Sebire NJ, Chitty LS. Is traditional perinatal autopsy needed after detailed fetal ultrasound and post-mortem MRI? *Prenat Diagn*. 2019 Aug;39(9):818-29.

113. O'Donnell C, Woodford N. Post-mortem radiology--a new sub-speciality? *Clin Radiol.* 2008 Nov;63(11):1189–94.
114. Addison S, Arthurs OJ, Thayyil S. Post-mortem MRI as an alternative to non-forensic autopsy in foetuses and children: from research into clinical practice. *Br J Radiol.* 2014 Apr;87(1036):20130621.
115. Arthurs OJ, van Rijn RR, Taylor AM, Sebire NJ. Paediatric and perinatal postmortem imaging: the need for a subspecialty approach. *Pediatr Radiol.* 2015 Apr;45(4):483–90.
116. Sebire NJ, Miller S, Jacques TS, Taylor AM, Rennie JM, Kendall G, Chitty LS. Post-mortem apparent resolution of fetal ventriculomegaly: evidence from magnetic resonance imaging. *Prenat Diagn.* 2013 Apr;33(4):360–4.
117. Arthurs OJ, Hutchinson JC, Sebire NJ. Current issues in postmortem imaging of perinatal and forensic childhood deaths. *Forensic Sci Med Pathol.* 2017 Mar;13(1):58–66.
118. Kang X, Cannie M, Jani JC. Effect of staining using gadolinium and formaldehyde on fetal whole-body postmortem 3-T magnetic resonance imaging. *Ultrasound Obstet Gynecol.* 2019 May;doi:10.1002/uog.20356.
119. Furness ME, Weckert RC, Parker SA, Knowles S. Ultrasound in the perinatal necropsy. *J Med Genet.* 1989 Jun;26(6):368–72.
120. Votino C, Cos Sanchez T, Bessieres B, Segers V, Kadhim H, Razavi F, Condorelli M, Votino R, D'Ambrosio V, Jani J. Minimally invasive fetal autopsy using ultrasound: a feasibility study. *Ultrasound Obstet Gynecol.* 2018 Dec;52(6):776–83. Prodhomme O, Baud C, Saguintah M, Béchard-Sevette N, Bolivar J, David S, Taleb-Arrada I, Couture A. Comparison of postmortem ultrasound and X-Ray with autopsy in fetal death: Retrospective study of 169 cases. *J Forensic Radiol Imaging.* 2015 Jun;3(2):120–30.
121. Kang X, Shelmerdine SC, Hurtado I, Bevilacqua E, Hutchinson C, Mandalia U, Segers V, Cos Sanchez T, Cannie MM, Carlin A, Sebire NJ, Arthurs OJ, Jani JC.

- Postmortem examination of human fetuses: a comparison of 2-dimensional ultrasound with invasive autopsy. *Ultrasound Obstet Gynecol.* 2019 Feb;53(2):229-38.
122. Prodhomme O, Baud C, Saguintaa M, Béchard-Sevette N, Bolivar J, David S, Taleb-Arrada I, Couture A. Comparison of postmortem ultrasound and X-Ray with autopsy in fetal death: Retrospective study of 169 cases. *J Forensic Radiol Imaging.* 2015 Jun;3(2):120–30.
 123. Tuchtan L, Lesieur E, Bartoli C, Delteil C, Sarda-Quarello L, Torrents J, Sigaudy S, Piercecchi MD, Gorincour G. Diagnosis of congenital abnormalities with postmortem ultrasound in perinatal death. *Diagn Interv Imaging.* 2018 Mar;99(3):143–9.
 124. Shelmerdine SC, Hickson M, Sebire NJ, Arthurs OJ. Post-Mortem Magnetic Resonance Imaging Appearances of Feticide in Perinatal Deaths. *Fetal Diagn Ther.* 2019;45(4):221-229.
 125. Kang X, Resta S, Cos T, Carlin A, Bevilacqua E, Jani JC. Impact of the delay between fetal death and delivery on the success of postmortem ultrasound following termination of pregnancy. *J Matern Fetal Neonatal Med.* 2019 Jul;1-6. doi:10.1080/14767058.2019.1642868.
 126. Metscher BD. MicroCT for developmental biology: a versatile tool for high-contrast 3D imaging at histological resolutions. *Dev Dyn.* 2009 Mar;238(3):632–40.
 127. Freeth T, Bitsakis Y, Moussas X, Seiradakis JH, Tselikas A, Mangou H, Zafeiropoulou M, Hadland R, Bate D, Ramsey A, Allen M, Crawley A, Hockey P, Malzbender T, Gelb D, Ambrisco W, Edmunds MG. Decoding the ancient Greek astronomical calculator known as the Antikythera Mechanism. *Nature.* 2006 Nov;444(7119):587–91.

128. Lombardi CM, Zambelli V, Botta G, Moltrasio F, Cattoretti G, Lucchini V, Fesslova V, Cuttin MS. Postmortem microcomputed tomography (micro-CT) of small fetuses and hearts. *Ultrasound Obstet Gynecol.* 2014 Nov;44(5):600–9.
129. Hutchinson JC, Arthurs OJ, Ashworth MT, Ramsey AT, Mifsud W, Lombardi CM, Sebire NJ. Clinical utility of postmortem microcomputed tomography of the fetal heart: diagnostic imaging vs macroscopic dissection. *Ultrasound Obstet Gynecol.* 2016 Jan;47(1):58–64.
130. Hutchinson JC, Barrett H, Ramsey AT, Haig IG, Guy A, Sebire NJ, Arthurs OJ. Virtual pathological examination of the human fetal kidney using micro-CT. *Ultrasound Obstet Gynecol.* 2016 Nov;48(5):663–5.
131. Hutchinson JC, Ashworth MT, Sebire NJ, Arthurs OJ. Multiple Cardiac Rhabdomyomas Visualised Using Micro-CT in a Case of Tuberous Sclerosis. *Fetal Diagn Ther.* 2017;41(2):157–60.
132. Apps JR, Hutchinson JC, Arthurs OJ, Virasami A, Joshi A, Zeller-Plumhoff B, Moulding D, Jacques TS, Sebire NJ, Martinez-Barbera JP. Imaging Invasion: Micro-CT imaging of adamantinomatous craniopharyngioma highlights cell type specific spatial relationships of tissue invasion. *Acta Neuropathol Commun.* 2016 Jun;4(1):57.
133. Hutchinson JC, Kang X, Shelmerdine SC, Segers V, Lombardi CM, Cannie MM, Sebire NJ, Jani JC, Arthurs OJ. Postmortem microfocus computed tomography for early gestation fetuses: a validation study against conventional autopsy. *Am J Obstet Gynecol.* 2018 Apr;218(4):445.e1-445.e12.
134. Shelmerdine SC, Hutchinson JC, Kang X, Suich JD, Ashworth M, Cannie MM, Segers V, Sebire NJ, Jani JC, Arthurs OJ. Novel usage of microfocus computed tomography (micro-CT) for visualisation of human embryonic development-Implications for future non-invasive post-mortem investigation. *Prenat Diagn.* 2018 Jun;38(7):538–42.

135. Hutchinson JC, Shelmerdine SC, Simcock IC, Sebire NJ, Arthurs OJ. Early clinical applications for imaging at microscopic detail: microfocus computed tomography (micro-CT). *Br J Radiol.* 2017 Jul;90(1075):20170113.
136. Shelmerdine SC, Singh M, Simcock I, Calder AD, Ashworth M, Beleza A, Sebire NJ, Arthurs OJ. Characterisation of Bardet Biedl Syndrome by post-mortem microfocus computed tomography (micro-CT). *Ultrasound Obstet Gynecol.* 2019 Jan;53(1):132-4.
137. Vymazalová K, Vargová L, Zikmund T, Kaiser J. The possibilities of studying human embryos and foetuses using micro-CT: a technical note. *Anat Sci Int.* 2017 Mar;92(2):299–303.
138. Arthurs OJ, Calder AD, Kiho L, Taylor AM, Sebire NJ. Routine perinatal and paediatric post-mortem radiography: detection rates and implications for practice. *Pediatr Radiol.* 2014 Mar;44(3):252–7.
139. Kamphuis-van Ulzen K, Koopmanschap DHJLM, Marcelis CLM, van Vugt JMG, Klein WM. When is a post-mortem skeletal survey of the fetus indicated, and when not? *J Matern-Fetal Neonatal Med.* 2016 Mar;29(6):991–7.
140. O'Donoghue K, O'Regan KN, Sheridan CP, O'Connor OJ, Benson J, McWilliams S, Moore N, Murphy MJ, Chopra R, Higging JR, Maher MM. Investigation of the role of computed tomography as an adjunct to autopsy in the evaluation of stillbirth. *Eur J Radiol.* 2012 Jul;81(7):1667–75.
141. Sarda-Quarello L, Tuchtan L, Torrents J, Piercecchi-Marti M-D, Bartoli C, Laurent P-E, Bourlière-Najean B, Petit P, Quarello E, Gorincour G. Perinatal death: Is there a place for post-mortem angio-CT? *J Forensic Radiol Imaging.* 2015 Mar;3(1):1–4.
142. Hutchinson C, Shelmerdine SC, Lewis C, Parmenter J, Simcock IC, Ward L, Ashworth MT, Chitty LS, Arthurs OJ, Sebire NJ. Feasibility and experience of the MinIMAL procedure: Minimally Invasive perinatal and paediatric

- Autopsies with Laparoscopically assisted tissue sampling. *Ultrasound Obstet Gynecol.* 2019 Jan;doi:10.1002/uog.20211.
143. Kang X, Cos T, Guizani M, Cannie MM, Segers V, Jani JC. Parental acceptance of minimally invasive fetal and neonatal autopsy compared with conventional autopsy. *Prenat Diagn.* 2014 Nov;34(11):1106-10.
 144. Man J, Hutchinson JC, Ashworth M, Judge-Kronis L, Levine S, Sebire NJ. Stillbirth and intrauterine fetal death: role of routine histological organ sampling to determine cause of death. *Ultrasound Obstet Gynecol.* 2016;48(5):596–601.
 145. Benn P, Cuckle H, Pergament E. Non-invasive prenatal testing for aneuploidy: current status and future prospects. *Ultrasound Obstet Gynecol.* 2013 Jul;42(1):15–33.
 146. Bevilacqua E, Resta S, Carlin A, Kang X, Cos Sanchez T, de Marchin J, Jani JC. Profile of women choosing the Harmony® Prenatal Test. *Expert Rev Mol Diagn.* 2018;18(7):591–9.
 147. Benachi A, Caffrey J, Calda P, Carreras E, Jani JC, Kilby MD, Klein HG, Rizzo G, Yaron Y. Understanding attitudes and behaviors towards cell-free DNA-based noninvasive prenatal testing (NIPT): A survey of European health-care providers. *Eur J Med Genet.* 2019 Jan;doi:10.1016/j.ejmg.2019.01.006.
 148. Norton ME, Baer RJ, Wapner RJ, Kuppermann M, Jelliffe-Pawlowski LL, Currier RJ. Cell-free DNA vs sequential screening for the detection of fetal chromosomal abnormalities. *Am J Obstet Gynecol.* 2016 Jun 1;214(6):727.e1-727.e6.
 149. Struksnaes C, Blaas H-GK, Eik-Nes SH, Vogt C. Correlation between prenatal ultrasound and postmortem findings in 1029 fetuses following termination of pregnancy. *Ultrasound Obstet Gynecol.* 2016 Aug;48(2):232–8.

150. Society for Maternal-Fetal Medicine (SMFM), Dugoff L, Norton ME, Kuller JA. The use of chromosomal microarray for prenatal diagnosis. *Am J Obstet Gynecol.* 2016 Oct;215(4):B2-9.
151. Best S, Wou K, Vora N, Van der Veyver IB, Wapner R, Chitty LS. Promises, pitfalls and practicalities of prenatal whole exome sequencing. *Prenat Diagn.* 2018;38(1):10–9.
152. Rauch A, Wieczorek D, Graf E, Wieland T, Endele S, Schwarzmayr T, Albrecht B, Bartholdi D, Beygo J, Di Donato N, Dufke A, Cremer K, Hempel M, Horn D, Hoyer J, Joset P, Röpke A, Moog U, Riess A, Thiel CT, Tzschach A, Wiesener A, Wohlleber E, Zweier C, Ekici AB, Zink AM, Rump A, Meisinger C, Grallert H, Sticht H, Schenck A, Engels H, Rappold G, Schröck E, Wieacker P, Riess O, Meitinger T, Reis A, Strom TM. Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. *Lancet.* 2012 Nov 10;380(9854):1674–82.
153. de Ligt J, Willemsen MH, van Bon BWM, Kleefstra T, Yntema HG, Kroes T, Vullo-van Silfhout AT, Koolen DA, de Vries P, Gilissen C, del Rosario M, Hoischen A, Scheffer H, de Vries BB, Brunner HG, Veltman JA, Vissers LE. Diagnostic exome sequencing in persons with severe intellectual disability. *N Engl J Med.* 2012 Nov;367(20):1921–9.
154. Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, Braxton A, Beuten J, Xia F, Niu Z, Hardison M, Person R, Bekheirnia MR, Leduc MS, Kirby A, Pham P, Scull J, Wang M, Ding Y, Pion SE, Lupski JR, Beaudet AL, Gibbs RA, Eng CM. Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N Engl J Med.* 2013 Oct;369(16):1502–11.
155. Sonnemans LJP, Vester MEM, Kolsteren EEM, Erwich JJHM, Nikkels PGJ, Kint , van Rijn RR, Klein WM; Dutch post-mortem imaging guideline group. Dutch guideline for clinical foetal-neonatal and paediatric post-mortem radiology, including a review of literature. *Eur J Pediatr.* 2018 Jun;177(6):791–803.

156. Weustink AC, Hunink MGM, van Dijke CF, Renken NS, Krestin GP, Oosterhuis JW. Minimally invasive autopsy: an alternative to conventional autopsy? Radiology. 2009 Mar;250(3):897–904.