

Oxygenated C₁₇ polyacetylene metabolites from Algerian *Eryngium tricuspidatum* L. roots: Structure and biological activity



A. Djebara^{a,b}, M.L. Ciavatta^{a,*}, V. Mathieu^{c,d}, M. Colin^c, F. Bitam^e, M. Carbone^a, M. Gavagnin^a

^a Consiglio Nazionale delle Ricerche (CNR), Istituto di Chimica Biomolecolare (ICB), Via Campi Flegrei, 34, 80078 Pozzuoli, Napoli, Italy

^b Laboratoire de Chimie et Chimie de l'Environnement, Faculté des Sciences de la Matière, Département de Chimie, Université de Batna-1, 05000, Algeria

^c Department of Pharmacotherapy and Pharmaceutics, Faculty of Pharmacy, ULB, Campus de la Plaine, Blvd du Triomphe, 1050 Brussels, Belgium

^d U-CRC, ULB Cancer Research Center, ULB, route de Lennik, 808, 1070 Brussels, Belgium

^e Faculté de Médecine, Département de Pharmacie, Université de Batna-2, 05000, Algeria

ARTICLE INFO

Keywords:

Eryngium tricuspidatum: Apiaceae

Polyacetylenes

Falcarindiol

Antiproliferative activity

ABSTRACT

The secondary metabolite pattern of *Eryngium tricuspidatum* has been found to be dominated by C₁₇ acetylene oxylipins, according to the chemistry reported in the literature for the genus *Eryngium*. Two new oxylipins, 11-acetoxy-falcarindiol (4) and 1,2-dihydro-11-acetoxy-falcarindiol (5) have been isolated, along with main related polyacetylenes 1–3 and the already known monoterpene aldehydes 6–10, from the petroleum ether extract of roots. The structure and the absolute configuration of compounds 4 and 5 have been determined by spectroscopic methods as well as by comparison with related known compounds. Polyacetylenes 1–4 inhibited significantly the in vitro growth of a series of cancer cell lines, ranging from 0.3 to 29 μM, whereas 5 was inactive.

1. Introduction

The genus *Eryngium* is the largest and arguably the most taxonomically complex genus of Apiaceae family (Umbelliferae) [1]. The genus comprises approximately 300 species throughout the world including species cultivated as ornamental, vegetable, or medicinal crops for folk uses. *Eryngium tricuspidatum* L. (syn. *Eryngium zanonii*) that is native to North Africa, Malta, Sicily, and Sardinia, is commonly known as sea holly and is characterized by dark blue flowers appearing in Summer and silver foliage. The perennials reach heights of 50 to 150 cm and produce achenes whereas the leaves are arranged opposite to one another and are pinnate. The plant grows in sunny sites and can withstand sandy and drought-prone soils as well as temperatures down to −12 °C [2].

E. tricuspidatum is one of seven *Eryngium* species endemic in Algeria, mainly distributed in the northern regions of the country [3]. Analogous with other plants of this genus, *E. tricuspidatum* (vernacular name: M'ghizla) is used by local populations as folk remedy for the treatment of various disorders. In particular, the decoction of its roots is effective against poisoning and constipation [4], and also to relieve labor pain [5].

Several studies on the chemistry and pharmacological activities of plants of *Eryngium* genus have been reported to date in the literature

[6]. A large array of chemicals, that are terpenoids, polyacetylenes, steroids, triterpenoid saponins, flavonoids, and coumarins, displaying a variety of biological properties such as antitumour, antibacterial, antimicrobial, antifungal, and phototoxic activities have been described from distinct species of *Eryngium* genus [6]. The phytochemical constituents of *E. tricuspidatum* have been investigated in recent years with regards to the aerial parts whereas, to the best of our knowledge, no chemical analysis of the roots of the plant has been reported to date. In particular, two papers described the chemical composition and biological activity, including antibacterial, antioxidant, tyrosinase inhibitory, and antifungal properties, of the essential oil [7] and phenolic glycoside [8] mixtures obtained from the extracts of the aerial parts of *E. tricuspidatum* collections from northwestern and northeastern Algeria, respectively.

In continuing our phytochemical studies on Algerian plants [9–12], we have investigated the content of the petroleum ether extract of the roots of *E. tricuspidatum* collected in North Algeria. This study has revealed a secondary metabolite pattern dominated by oxygenated C₁₇ polyacetylenes (compounds 1–5) and monoterpene aldehydes (compounds 6–9). Compounds 4 and 5 were new enynols structurally related to co-occurring (3S,8S)-falcarindiol (3). Falcarindiol-3,8-diacetate (1) was never reported as naturally occurring metabolite. The in vitro growth inhibitory properties of polyacetylenes 1–5 on a panel of cancer

* Corresponding author.

E-mail addresses: lcivatta@icb.cnr.it (M.L. Ciavatta), Veronique.Mathieu@ulb.ac.be (V. Mathieu), Margaux.Colin@ulb.ac.be (M. Colin), f.bitam@univ-batna2.dz (F. Bitam), mcarbone@icb.cnr.it (M. Carbone), mgavagnin@icb.cnr.it (M. Gavagnin).

<https://doi.org/10.1016/j.fitote.2019.104355>

Received 22 July 2019; Received in revised form 6 September 2019; Accepted 10 September 2019

Available online 11 September 2019

0367-326X/© 2019 Elsevier B.V. All rights reserved.

cell lines has been also evaluated and here reported.

2. Experimental

2.1. General experimental procedures

Optical rotations were measured on a Jasco DIP 370 digital polarimeter and a A. KRÜSS OPTRONIC P3000 polarimeter. ESIMS were performed on a Micromass Q-TOF MicroTM coupled with a HPLC Waters Alliance 2695. The instrument was calibrated by using a PEG mixture from 200 to 1000 MW (resolution specification 5000 FWHM, deviation < 5 ppm RMS in the presence of a known lock mass). High resolution mass spectra (HRESIMS) were acquired on a Q-Exactive hybrid quadrupole-orbitrap mass spectrometer (Thermo Scientific). NMR experiments were recorded at ICB-NMR Service Centre. Chemical shifts values are reported in ppm and referenced to internal signals of residual protons (CDCl₃, ¹H δ 7.26, ¹³C 77.0 ppm; DMSO-*d*₆, ¹H δ 2.50, ¹³C 40.0 ppm). 1D and 2D NMR spectra were acquired on a Bruker Avance-400 spectrometer using an inverse probe fitted with a gradient along the Z-axis, on a Bruker Avance III HD spectrometer equipped with a CryoProbe Prodigy, and on a Bruker DRX-600 operating at 600 MHz, using an inverse TCI CryoProbe fitted with a gradient along the Z-axis. HPLC separation was performed on a Shimadzu high-performance liquid chromatography using a Shimadzu liquid chromatograph LC-10 CE equipped with an UV SPD-10A wavelength detector, with a reversed-phase (RP-18) column (4.6 × 250 mm, Kromasil-Phenomenex). Silica-gel chromatography was performed using pre-coated Merck F254 plates (TLC) and Merck Kieselgel 60 powder (70–230 mesh). The spots on TLC were visualized under UV light (254 nm) and then spraying them with 10% H₂SO₄ in water followed by heating.

2.2. Plant material

The plant *Eryngium tricuspidatum* was collected in El Madher, in the northeastern region of Batna (Algeria) and identified by Prof. Bachir Oudjehih, Institute of Agronomy of University of Batna (Algeria). A voucher specimen is deposited in the herbarium of the department of the same University with code 05/ISAV/D.A.G./2015.

2.3. Extraction and isolation

Air-dried aerial parts (1.5 kg) and roots (0.8 kg) of *E. tricuspidatum* were accurately separated, powdered, and exhaustively extracted at room temperature with an hydroalcoholic solution (EtOH/H₂O, 8:2) for five times (2.5 L × 5). The extracts obtained for each part were filtered, combined and concentrated under vacuum to afford two distinct aqueous solutions which were sequentially partitioned with solvents of increasing polarity: petroleum ether (500 mL × 3), EtOAc (500 mL × 3), and *n*-BuOH (500 mL × 3). After removing the organic solvents, crude residues from petroleum ether (14.0 g for aerial parts, 7.0 g for roots), EtOAc (18.0 g for aerial parts, 6.0 g for roots), and *n*-BuOH (35.0 g for aerial parts, 20.0 g for roots) were obtained, respectively. All these extracts were analysed by TLC chromatography. A portion (4.0 g) of the petroleum ether extract of roots was fractionated by silica gel column chromatography (column diameter: 4.5 cm, height: 120 cm, silica gel: 200 g), eluting with an increasing polarity gradient of petroleum ether/Et₂O, and then CHCl₃/MeOH. A total of 62 fractions (volume of each tube: 50 mL) were collected starting from petroleum ether/Et₂O, 8:2 (500 mL of solvent, collected 10 tubes), petroleum ether/Et₂O, 7:3 (400 mL of solvent 8 tubes), petroleum ether/Et₂O, 6:4 (350 mL of solvent, 7 tubes), petroleum ether/Et₂O, 1:1 (300 mL of solvent, 6 tubes), petroleum ether/Et₂O, 3:7 (800 mL of solvent, 16 tubes), Et₂O (300 mL of solvent, 6 tubes), CHCl₃/MeOH 9:1 (350 mL of solvent, 7 tubes), and only MeOH (100 mL, 2 tubes). After TLC chromatography all these fractions were combined to give 21 final fractions

(F1/1 → F1/21) some of which were subjected to subsequent purifications aided by ¹H NMR analysis. A portion of F1/2 (30 mg), eluted with petroleum ether/ Et₂O 8:2 (800 mg), was further purified on semipreparative TLC (petroleum ether/ Et₂O 9:1) to give compounds 1 (Rf 0.45, 6.0 mg), 6 (Rf 0.4, 5.0 mg) and 7 (Rf 0.6, 9.0 mg), and a mixture of fatty acid methyl esters. Also a portion (110 mg) of F1/3, eluted with petroleum ether/ Et₂O 8:2 (340 mg), was purified by semipreparative TLC (petroleum ether/ Et₂O 8:2) yielding compound 6 (Rf 0.55, 18.0 mg), 8 (Rf 0.5, 1.7 mg) and 2 (Rf 0.25, 25 mg). Compound 2 was also the main component of fractions F1/4 (190 mg), F1/5 (270 mg), and F1/6 (600 mg), whereas compound 3 was present in both F1/8 (640 mg) and F1/9 (250 mg) from which it has been purified by two subsequent purifications. An aliquot (50 mg) of F1/8, eluted from the first column in petroleum ether/ Et₂O 7:3, was loaded into a SPE cartridge C18 using as eluent MeOH/H₂O 8:2. This subfraction (25 mg) was further purified by semipreparative TLC chromatography, using as solvent system CH₂Cl₂/MeOH 98:2 to obtain compound 3 (Rf 0.55, 14.5 mg) and 9 (Rf 0.85, 7.0 mg). Fractions F1/10 (89 mg), and F1/12 (180 mg), eluted from the first column with petroleum ether/ Et₂O from 6:4 to 1:1, were combined and subjected to a first purification on SiO₂ gel column (column diameter: 1.8 cm, height: 120 cm, silica gel: 15 g, collected tubes: 41 tubes, each tube volume: 15 mL) using a gradient solvent system of petroleum ether/ Et₂O (100:0 to 0:100, v/v). Subfractions eluted with 30% to 40% of Et₂O in petroleum ether were combined (110 mg) and loaded onto a SPE cartridge C18 packed in MeOH/H₂O 3:7 and eluted first with MeOH/H₂O 3:7, then with MeOH/H₂O 1:1, MeOH/H₂O 6:4, MeOH/H₂O 8:2 and finally with only MeOH. Subfraction eluted with MeOH/H₂O 1:1 (14.5 mg) was further purified by analytical reversed phase HPLC column (Kromasil C18, eluent MeOH/H₂O 75:25, flow 1.0 mL/min, UV detector 210 and 254 nm) obtaining compounds 4 (1.5 mg, Rt 11.9) and 5 (1.0 mg, Rt 13.9). Finally a portion (40.0 mg) of F1/11 was loaded onto a semipreparative SiO₂ TLC plate in CHCl₃/MeOH 98:2 to give pure ferulol (10, Rf 0.2, 10.0 mg).

2.3.1. 11-Acetoxy-falcarindiol (4)

Pale yellow oil; [α]_D²⁵ + 141.1 (c 0.39, CHCl₃); HRESIMS: *m/z* 341.1733 [M + Na]⁺ (calcd. 341.1723 for C₁₉H₂₆O₄Na); ¹H and ¹³C NMR data, see Table 1.

2.3.2. 1,2-Dihydro-11-acetoxy-falcarindiol (5)

Pale yellow oil; [α]_D²⁵ + 41.3 (c 0.06, CHCl₃); HRESIMS: *m/z* 343.1888 [M + Na]⁺ (calcd. 343.1880 for C₁₉H₂₈O₄Na); ¹H and ¹³C NMR data, see Table 1.

2.4. Biological evaluation

2.4.1. Cell lines

Human cancer cell lines were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA), the European Collection of Cell Culture (ECACC, Salisbury, UK) and the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ, Braunschweig, Germany). B16F10 mouse melanoma (ATCC code CRL 6475) cells were cultured in RPMI culture medium (Gibco, Thermofisher, Dilbeek, Belgium) supplemented with 10% heat-inactivated fetal bovine serum (Gibco), like human melanoma SKMEL-28 cells (ATCC HTB72), human mammary carcinoma MCF-7 (ATCC HTB22), A549 non small cell lung carcinoma cells (DSMZ ACC107), Hs683 anaplastic oligodendroglioma cells (ATCC HTB-138), and U373 glioblastoma cells (ECACC 08061901). Cell culture media were supplemented with 0.6 mg/mL L-glutamine (Gibco), 100 µg/mL gentamicin (Gibco), and penicillin-streptomycin (200 units/mL; Gibco). All cell lines were cultured in T25 flasks (Sarstedt AG & CO, Nümbrecht, Germany), maintained and grown at 37 °C, 95% humidity and 5% CO₂.

Table 1
¹³C (150 MHz) and ¹H (600 MHz) NMR data^a for compounds **4** and **5** in CDCl₃.

Position	4		5	
	δ _c type	δ _H (J, Hz)	δ _c type	δ _H (J, Hz)
1a	117.3, CH ₂	5.49 (br d, 16.4)	9.3, CH ₃	1.01 (t, 7.5)
1b		5.26 (br d, 10.2)		
2	135.7, CH	5.94 (ddd, 16.4, 10.2, 5.4)	30.6, CH ₂	1.74 (m)
3	63.5, CH	4.94 (m)	64.1, CH	4.38 (app t, 6.4)
4	77.9, C	–	nd	–
5	69.6, C	–	nd	–
6	68.8, C	–	69.2, C	–
7	79.1, C	–	78.8, C	–
8	58.6, CH	5.45 (dd, 8.7, 3.6)	58.6, CH	5.45 (d, 8.6)
9	132.0, CH	5.74, (br dd, 9.8, 8.7)	132.2, CH	5.74, (br dd, 9.8, 8.6)
10	130.2, CH	5.35 (overlapped)	130.2, CH	5.36 (overlapped)
11	70.5, CH	5.35 (overlapped)	70.6, CH	5.36 (overlapped)
12	33.8, CH ₂	1.69 (m)	33.8, CH ₂	1.68 (m)
		1.54 (m)		1.54 (m)
13	24.9, CH ₂	1.30 (m)	24.9, CH ₂	1.29 (m)
14	28.9, CH ₂	1.30 (m)	29.0, CH ₂	1.30 (m)
15	31.6, CH ₂	1.28 (m)	31.7, CH ₂	1.28 (m)
16	22.5, CH ₂	1.29 (m)	22.5, CH ₂	1.29 (m)
17	14.0, CH ₃	0.89 (t, 6.9)	14.1, CH ₃	0.89 (t, 6.6)
OAc-11	171.5, CO		171.5, CO	
	21.3, CH ₃	2.03 (s)	21.3, CH ₃	2.03 (s)

^a Assignments aided by COSY, edHSQC, HMBC ($J = 8$ Hz).

2.4.2. Antiproliferative properties

Antiproliferative properties of the compounds were evaluated by the MTT assay. All compounds were dissolved in DMSO at a concentration of 10 mM prior to cell treatment. The cells were trypsinized and seeded at various cell concentrations depending on the cell type into 96-well plates (Sarstedt). The cells were grown for 24 h, treated with compounds at concentrations ranging from 0.001 to 100 μM and incubated for 72 h in 100 μL media depending. After the treatment, culture media were replaced by 100 μL of MTT reagent (Sigma-Aldrich, St. Louis, MO) in serum free medium (0.5 mg/mL) and incubated further for 4 h. Formazan crystals formed by viable mitochondrial reduction were resolubilized in 100 μL of DMSO. Absorbance was measured at 570 nm with a spectrophotometer (680XR, Bio-Rad Laboratories, Berkeley, CA, USA; reference wavelength 610 nm). The experiments were performed once in six replicates.

2.4.3. Computer-assisted phase contrast microscopy

Computer-assisted phase contrast microscopy was performed as previously described [13]. Briefly, U373 or SKMEL-28 cells were seeded in 25 cm² culture flasks (Sarstedt) and left untreated or treated with the compounds at their IC₅₀ concentrations determined with the MTT colorimetric assay. Pictures of one field were taken every four min during a 72 h period and further compiled into a short movie [13]. Experiments were performed once in triplicates.

3. Results and discussion

The petroleum ether extracts of *E. tricuspidatum* aerial parts and roots were compared by TLC in different eluent systems. Significant differences in the composition and in the relative distribution of the metabolites were observed between the two extracts. In particular, the presence of a rich metabolite pattern characterized by a series of UV visible metabolites was observed almost exclusively in the roots. Thus, the petroleum ether extract of roots was taken into consideration for the chemical analysis. An aliquot (4.0 g) was fractionated as before described (Experimental) to give after sequential chromatographic steps five acetylene oxylipins, falcarindiol-3,8 diacetate (**1**) [14], falcarindiol-8-acetate (**2**) [15], falcarindiol (**3**) [16,17], new compounds 11-

acetoxy-falcarindiol (**4**) and 1,2-dihydro-11-acetoxy-falcarindiol (**5**), along with four monoterpenoid aldehydes, isoferulol angelate (**6**) [18], 2,3,4-trimethyl-benzaldehyde (**7**) [19], ferulol-O-methyl ether (**8**) [20], isoferulol-5'-acetoxy angelate (**9**) [21], and finally ferulol (**10**) [21,22]. Known compounds were identified by ¹H and ¹³C NMR and MS spectroscopic data (Supplementary Material). As NMR data of monoterpenoid aldehydes were only partially reported in the literature [18–22], the complete NMR assignment was also made (Supplementary Material). In addition, stereochemical aspects of falcarindiol (**3**) and derivatives were investigated. The structures of compounds **4** and **5** were determined by spectroscopic analysis and correlation with co-occurring polyacetylenes.

The polyacetylene pool was predominant in the extract (ca. 60%) and compounds **2** and **3** were the main components of this fraction. Polyacetylenes of falcarindiol series are typical metabolites of several members of the Apiaceae and Arialiaceae families [23–25] including edible plants such as carrots. [26,27] (Fig. 1).

Falcarindiol-3,8-diacetate (**1**) was a minor component of the extract that has never been isolated as naturally occurring compound to date. It has been previously reported in the literature only as derived by acetylation of falcarindiol (**3**) [14]. Falcarindiol-8-acetate (**2**) has been identified in a limited number of plants [15,28] whereas well-known falcarindiol (**3**), first isolated from *Falcaria vulgaris* [16], is a very common metabolite widely distributed in several species including important food vegetables [23–25].

The physical and chemical properties including the absolute configuration of **3** have been described in a number of papers [23–25] even though some conflicting data have been reported with regards to the stereochemical determination of C-3 and C-8 stereogenic centers [29–33]. The uncertainty in the stereochemical assignment was probably due to the difficulty of distinguishing the isomers by comparison of NMR and optical rotation data. The stereochemistry of **3** has been also investigated by a series of synthesis studies [27,34–36]. All four stereoisomers [(3*R*,8*S*), (3*S*,8*S*), (3*S*,8*R*), (3*R*,8*R*)] of falcarindiol have been synthesized [27,36] and their optical rotation have been measured showing that the sign of [α]_D is mainly affected by C-8 configuration (8*R* in levorotatory isomers, 8*S* in dextrorotatory isomers) [36]. Both (3*R*)- and (3*S*)- falcarindiol stereoisomers have been found in nature whereas the *S* configuration at C-8 has been assigned for all samples isolated from different sources despite the negative [α]_D value observed in some cases [32]. In addition, falcarindiol (**3**) isolated from an *Oplopanax* species has been found to be a mixture of C-3 epimers by careful NMR analysis of the Mosher ester derivatives [33]. Based on this, we have thus investigated the stereochemistry of falcarindiol (**3**) isolated in this work by analysis of NMR, [α]_D and CD data as well as by application of the Mosher method. Comparison of the ¹H and ¹³C NMR spectra with those described for synthetic stereoisomers [36] along with evaluation of the [α]_D value [36] and CD profile [33] strongly suggested the (3*S*,8*S*) configuration. In order to further support this assignment, the modified Mosher method was applied. The NMR analysis of both *S* and *R* MTPA esters of **3** confirmed the *S* absolute configuration at both stereogenic centers (see Supplementary Material) and also revealed the presence of small amounts of (3*R*,8*S*)-falcarindiol (**3**), co-occurring with the major stereoisomer. In fact, the spectra of both *S* and *R* MTPA esters contained two sets of H₂-1 and H-2 multiplets easily distinguishable and attributable to major (3*S*)- and minor (3*R*)-falcarindiol stereoisomers. Integration of these multiplets indicated a 3*S*:3*R* ratio as approximately 10:1. The finding of C-3 epimers of falcarindiol was in accordance with the previous report on *Oplopanax elatus* [33] even though it should be evidenced that in *E. tricuspidatum* the relative ratio of 3*S*/3*R* isomers is significantly higher with respect to the literature [33]. Falcarindiol-8-acetate (**2**) was also investigated for the stereochemical aspects and the Mosher method was applied to secure the absolute configuration of C-3. Analogous with falcarindiol (**3**), the *S* and *R* MTPA esters of **2** contained small proton signals, which were readily recognizable from H₂-1 and H-2 multiplets by about 10%




Fig. 1. Chemical structures of compounds isolated from *Eryngium tricuspdatum*.

smaller integral, confirming the expected 3*S* absolute configuration for the main stereoisomer (3*S*/3*R*, 10:1 ca.) (see Supplementary Material). Based on biogenetic grounds, the absolute configuration at C-8 was assigned to be *S*, the same as falcarindiol (**3**). Similar considerations were made for falcarindiol-3,8-diacetate (**1**) for which the 3*S*,8*S* absolute configuration was also proposed. Samples of falcarindiol-8-acetate (**2**) and falcarindiol (**3**) were acetylated and the diacetyl derivatives obtained were identical (^1H and ^{13}C NMR, $[\alpha]_D$, and MS) with natural **1**.

The HRESIMS spectrum of 11-acetoxy-falcarindiol (**4**) contained a sodium adduct ion at m/z 341.17333 $[\text{M} + \text{Na}]^+$, which was consistent with the molecular formula $\text{C}_{19}\text{H}_{26}\text{O}_4$ and implying an additional acetoxy group with respect to falcarindiol (**3**). Careful analysis of ^1H and ^{13}C NMR data (Table 1) indicated the close similarity with **3**, in particular suggesting the presence of typical enynol partial structure C-1/C-7 with a terminal double bond, a carbinol methine and two triple bonds, sequentially. The remaining part of the C_{17} chain contained a double bond along with a hydroxy and an acetoxy group. The inspection of the ^1H - ^1H COSY spectrum of **4** secured the expected C-1/C-3 spin-system [δ_{H} 5.94 (1*H*, ddd, $J = 16.4, 10.2, 5.4$ Hz, H-2), 5.49 (1*H*, br d, $J = 16.4$ Hz, H-1a), 5.26 (1*H*, br d, $J = 10.2$ Hz, H-1b), and 4.94 (1*H*, m, H-3)] and led to the definition of the subsequent carbon sequence establishing the position of the acetoxy group at C-11. The carbinol methine C-8 [δ_{H} 5.45 (1*H*, dd, $J = 8.7, 3.6$ Hz, H-8)] was

connected to vinyl carbon C-9 [δ_{H} 5.74 (1*H*, br dd, $J = 9.8, 8.7$ Hz, H-9)] whereas the second vinyl carbon C-10 [δ_{H} 5.35 (1*H*, overlapped, H-10)] linked the methine bearing the acetoxy group [δ_{H} 5.35 (1*H*, overlapped, H-11) and 2.03 (3*H*, s, -OAc)]. Analysis of HMBC experiments showed diagnostic cross-peaks from H-8 to C-6 (δ_{C} 68.8) and C-7 (δ_{C} 79.1), from H-9 to C-6 and C-11 (δ_{C} 70.5), from H-11 to C-8 (δ_{C} 58.6) and C-12 (δ_{C} 33.8) confirming this assignment. The *Z*-geometry of Δ^9 double bond was inferred by the coupling constant value of olefinic protons ($J_{\text{H}_9, \text{H}_{10}} = 9.8$ Hz) whereas the absolute configuration of C-3 and C-8 was suggested to be 3*S*,8*S*, the same as co-occurring main polyacetylenes **1**-**3**, by biogenetic considerations. The absolute configuration of C-11 remained undetermined.

1,2-Dihydro-11-acetoxy-falcarindiol (**5**) had the molecular formula $\text{C}_{19}\text{H}_{28}\text{O}_4$ as it was deduced from the sodium adduct ion at m/z 343.18887 $[\text{M} + \text{Na}]^+$ in the HRESIMS spectrum. Comparison of ^1H and ^{13}C NMR spectra with those of co-occurring compounds **1**-**4** revealed that **5** was lacking in the terminal double bond which was replaced by an ethyl moiety [δ_{H} 1.01 (3*H*, t, $J = 7.5$ Hz, H₃-1) and 1.74 (2*H*, m, H₂-2); δ_{C} 9.3 (C-1) and 30.6 (C-2)] linked to C-3 carbinol methine [δ_{H} 4.38 (1*H*, apparent t, $J = 6.4$ Hz, H-3)].

On the other side, the structure of **5** exhibited the fragment C-8/C-11 containing Δ^9 double bond [δ_{H} 5.74 (1*H*, br dd, $J = 9.8, 8.6$ Hz, H-9) and 5.36 (1*H*, m, overlapped, H-10); δ_{C} 132.2 (C-9) and 130.2 (C-10)] connected to CH(OH)- [δ_{H} 5.45 (1*H*, d, $J = 8.6$ Hz, H-8); δ_{C} 58.6 (C-8)]

Table 2
Determination of the IC₅₀ (μM) in vitro growth inhibitory concentration by means of the MTT colorimetric assay in six cancer cell lines.

Cancer cell lines versus compounds (IC ₅₀ , μM)	Melanoma		Carcinoma		Glioma		Mean ± SEM
	B16F10	SKMEL28	A549	MCF-7	HS683	U373	
1	4 ± 0,3	9 ± 0,2	27 ± 0,5	18 ± 0,9	29 ± 0,9	3 ± 0,1	15 ± 5
2	0,3 ± 0,01	0,6 ± 0,01	2,5 ± 0,1	3 ± 0,7	3 ± 0,4	0,3 ± 0,02	1 ± 0,6
3	1 ± 0,04	0,5 ± 0,04	3 ± 0,09	21 ± 1	4 ± 0,07	0,7 ± 0,06	5 ± 3
4	2 ± 0,1	3 ± 0,09	3 ± 0,1	26 ± 0,6	16 ± 2,3	13 ± 4	10 ± 4
5	88 ± 3	> 100	> 100	> 100	> 100	> 100	98 ± 2
Cisplatin^a	4 ± 0,1	9 ± 0,2	NT	NT	4 ± 0,1	2 ± 0,2	5 ± 2

Average concentration required to reduce the viability of cells by 50% after a 72 h treatment relative to a control (untreated cells) as determined by MTT assay performed once in sextuplicates., NT: not tested.

^a Tested in a separated experiment but with the same protocol and by the same experimenter.

and CH(OAc)- [δ_{H} 5.36 (1H, m, overlapped, H-11) and 2.01 (3H, s, -OAc); δ_{C} 70.6 (C-11), 171.5 (-OAc) and 21.3 (-OAc)] moieties, the same as **4**. The Δ^9 double bond was suggested to be Z by the coupling constant value of olefinic protons ($J_{\text{H9,H10}} = 9.8$ Hz). Analogous to co-occurring metabolites, the 3*S*,8*S* absolute configuration could be proposed whereas the stereochemistry at C-11 was undefined.

Various and significant biological properties have been reported in the literature for plant acetylenic metabolites [37]. Among these, faltarindiol (**3**) is one of the most bioactive dietary polyacetylene having a vast array of biological properties including anti-inflammation, antibacterial, antifungal, antimutagenic, antiproliferative and anticancer activities [37]. Interestingly, faltarindiol (**3**) has been shown to affect cell proliferation in a biphasic manner (hormesis) thus implying that benefits were achieved at non-toxic concentrations [38]. Recent pharmacological studies confirmed the anticancer activity of **3** against some selected cancer cells such as colonrectal cancer [39], breast cancer [40],

and glioblastoma cells [41]. Based on the literature data, we decided to investigate the effects on the cell proliferation of new faltarindiol-derived compounds **4** and **5** compared with faltarindiol (**3**) and acetyl derivatives **1** and **2**. A panel of six cancer cell lines containing melanoma, carcinoma and glioma models was considered and the in vitro growth inhibitory concentration for each tested compound, as determined by MTT assay, is reported in Table 2. Possible interference in terms of intrinsic absorbance and chemical reactivity with MTT salt has been ruled out for each compound (data not shown). All faltarindiol-type polyacetylenes **1–4** showed significant inhibition of cell proliferation against the cell lines under study with IC₅₀ ranging from 0.3 to 29 μM whereas the dihydro-derivative **5** was inactive against all of them. These results were in agreement with previous reports [26,42] and supported the presence of the terminal double bond as an important structural requirement for the activity.

Importantly, faltarindiol-8-acetate (**2**) displayed significant higher




Fig. 2. Quantitative videomicroscopy analyses of compounds **1–3**-induced effects on SKMEL-28 melanoma (time-lapse pictures over time). Vacuoles (white arrows) appear at different timepoint according to the compound tested at its IC₅₀. Faltarindiol (**3**) appeared to trigger cell stress earlier than compounds **1** and **2**. Cell death is consequently observed in all cases. Morphological characteristics are suggestive of apoptosis (black arrows).

inhibition on the proliferation of the cell lines than falcarindiol-3,8-diacetate (**1**) ($p = .017$; 2-tailed comparison conducted on the active compounds of this chemical series i.e. compounds **1–4**) but such statistical difference could not be found with falcarindiol (**3**) suggesting a substantial structural equivalence for the activity of 3-OH and 8-OAc moieties. The ANOVA analysis of the data of compounds **1–4** between the cell lines also showed significant variation in their level of sensitivity to the different polyacetylenes. Although A549 appeared for instance less affected by compound **1** (IC₅₀ ten-fold higher with respect to compound **2**) and MCF-7 less affected after treatment with **4** (IC₅₀ ten-fold higher with respect to compound **2**), the 2-tailed analysis did not evidenced one cell model to be more or less sensitive to the whole chemical series.

To further confirm the anti-cancer effects observed by MTT assay, morphological evaluation of those of compounds **1–3** has been assessed by time-lapse phase contrast videomicroscopy. As illustrated in Fig. 2, all compounds **1–3** induced vacuoles in melanoma cells (SKMEL-28) at their own IC₅₀ concentration but cell death occurs faster with compound **3**, i.e. within 24 h compared to 48 h for both compounds **1** and **2**. Vacuoles observed with compounds **1–3** (Fig. 2, white arrows) could derive from ER swelling due to the ER stress which may lead in turn to apoptosis in a similar manner to previously deciphered anti-cancer effects induced by falcarindiol (**3**) in colon cancer models [39].

In summary, five C₁₇ polyacetylenes **1–5** of the falcarindiol series, including two new compounds **4** and **5**, and five monoterpene aldehydes **6–10** were isolated from the lipophilic extract of the roots of Algerian plant *E. tricuspidatum*. The structures of new oxylipins **4** and **5** were established by spectroscopic methods, mainly by NMR analysis, and by chemical correlation with co-occurring known compounds. Stereochemical aspects were also investigated and the absolute configuration of selected stereogenic centers was secured by the Mosher method. Full NMR assignment of aldehydes **6–10**, the spectroscopic characterization of which was not previously reported in the literature, was also made. Finally, the effects on the cell proliferation against a panel of cancer cell lines of new oxylipins **4** and **5** were investigated compared with falcarindiol (**3**) and acetyl derivatives **1** and **2**. Compound **1–4** showed significant inhibition against all cell lines under study whereas the dihydro-derivative **5** was inactive against all of them. In addition, the morphological evaluation by time-lapse phase contrast videomicroscopy of the anti-cancer effects induced by oxylipins **1–3** against SKMEL-28 melanoma cells suggested the apoptosis mechanism, analogous with that reported in the literature for falcarindiol (**3**) in colon cancer model.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. They declare don't have any conflict of interest.

Acknowledgment

The authors acknowledge Dr. A. Cutignano and Mr. M. Zampa for the ICB-MS spectrometry service and Mr. V. Mirra of the ICB-NMR service. Thanks are also due to Prof. Bachir Oudjehih for identification of the plant. Authors acknowledge Miss Rita Roberta Serino for lab assistance. This research work was granted by MIUR-ITALY PRIN2015 “Top-down and bottom-up approach in the development of new bioactive chemical entities inspired on natural products scaffolds” (Project No. 2015MSCKCE_004). A.D. is deeply grateful to the Ministry of Higher Education and Scientific Research, Algeria for financial support. Part of this work was financially supported by the Belgium Brain Tumor Support (BBTS, Belgium).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fitote.2019.104355>.

References

- [1] A. Wörz, A taxonomic index of the species of *Eryngium* L. (Apiaceae: Saniculoideae), Stuttgart. Beitr. Naturk. Ser. A 596 (1999) 1–48.
- [2] C. Brickell, Encyclopedia of Garden Plants, 3rd ed, Dorling Kindersley, London, 2003.
- [3] P. Quezel, S. Santa, Nouvelle Flore de l'Algérie et des Régions Désertiques Méridionales, 1–2 CNRS, Paris, 1963, p. 650.
- [4] J. Bamm, A. Douira, Les plantes médicinales dans la forêt de l'Achach (Plateau Central, Maroc), Acta Bot. Malacit. 27 (2002) 131–145.
- [5] S. Salhi, M. Fadli, L. Zidane, A. Douira, Etudes floristique et ethnobotanique des plantes médicinales de la ville de Kénitra (Maroc), Lazaroa 31 (2010) 133–146, https://doi.org/10.5209/rev_LAZA.2010.v31.9.
- [6] S.A. Erdem, S.F. Nabavi, I.E. Orhan, M. Daglia, M. Izadi, S.M. Nabavi, Blessings in disguise: a review of phytochemical composition and antimicrobial activity of plants belonging to the genus *Eryngium*, DARU J. Pharm. Sci. 23 (2015) e53, <https://doi.org/10.1186/s40199-015-0136-3>.
- [7] D. Merghache, Z. Boucherit-Otmani, S. Merghache, I. Chikhi, C. Selles, K. Boucherit, Chemical composition, antibacterial, antifungal and antioxidant activities of Algerian *Eryngium tricuspidatum* L. essential oil, Nat. Prod. Res. 28 (2014) 795–807, <https://doi.org/10.1080/14786419.2014.883392>.
- [8] A. Benmerache, A. Alabdul Magid, D. Berrehal, A. Kabouche, L. Voutquenne-Nazabadioko, S. Messaili, A. Abedini, D. Harakat, Z. Kabouche, Chemical composition, antibacterial, antioxidant and tyrosinase inhibitory activities of glycosides from aerial parts of *Eryngium tricuspidatum* L., Phytochem. Lett. 18 (2016) 23–28, <https://doi.org/10.1016/j.phytol.2016.08.018>.
- [9] F. Bitam, M.L. Ciavatta, E. Manzo, A. Dibi, M. Gavagnin, First chemical characterisation of the terpenoid constituents of the Algerian plant *Launaea arborescens*, Phytochemistry 69 (2008) 2984–2992, <https://doi.org/10.1016/j.phytochem.2008.09.025>.
- [10] F. Bouzergoune, M.L. Ciavatta, F. Bitam, M. Carbone, M.C. Aberkane, M. Gavagnin, Phytochemical study of *Eryngium triquetrum*: isolation of polyacetylenes and lignans, Planta Med. 82 (2016) 1438–1445, <https://doi.org/10.1055/s-0042-110316>.
- [11] M. Boumaraf, M. Carbone, M.L. Ciavatta, S. Benyahia, S. Ameddah, A. Menad, S. Benayache, F. Benayache, M. Gavagnin, Exploring the bioactive terpenoid content of an Algerian plant of genus *Pulicaria*: the ent-series of asteriscunolides, J. Nat. Prod. 80 (2017) 82–89, <https://doi.org/10.1021/acs.jnatprod.6b00717>.
- [12] N. Zergainoh, M.L. Ciavatta, M. Carbone, F. Bitam, M.C. Aberkane, M. Gavagnin, Exploring the chemical diversity of Algerian plants: three new pentacyclic triterpenoids from *Launaea acanthoclada* roots, Molecules 23 (2018) e80, <https://doi.org/10.3390/molecules23010080>.
- [13] O. Debeir, V. Mégalizzi, N. Warzée, R. Kiss, C. Decaestecker, Videomicroscopic extraction of specific information on cell proliferation and migration *in vitro*, Exp. Cell Res. 314 (2008) 2985–2998.
- [14] K.W. Tan, D.P. Killeen, Y. Li, J.W. Paxton, N.P. Birch, A. Scheepens, Dietary polyacetylenes of the falcarinol type are inhibitors of breast cancer resistance protein (BCRP/ABCG2), Eur. J. Pharmacol. 723 (2014) 346–352, <https://doi.org/10.1016/j.ejphar.2013.11.005>.
- [15] T. Fujioka, K. Furumi, H. Fujii, H. Okabe, K. Mihashi, Y. Nakano, H. Matsunaga, M. Katano, M. Mori, Antiproliferative constituents from Umbelliferae plants. V. a new furanocoumarin and falcarindiol furanocoumarin ethers from the root of *Angelica japonica*, Chem. Pharm. Bull. 47 (1999) 96–100, <https://doi.org/10.1248/cpb.47.96>.
- [16] F. Bohlmann, U. Niedballa, K.-M. Rode, Polyacetylenverbindungen, CXVIII. Über neue polyne mit C17-Kette, Chem. Ber. 99 (1966) 3552–3558, <https://doi.org/10.1002/cber.19660991122>.
- [17] S. Deng, Y. Wang, T. Inui, S.-N. Chen, N.R. Farnsworth, S. Cho, S.G. Franzblau, G.F. Pauli, Anti-TB polyynes from the roots of *Angelica sinensis*, Phytother. Res. 22 (2008) 878–882, <https://doi.org/10.1002/ptr.2303>.
- [18] A.F. Barrero, A. Haïdour, F. Reyes, Monoterpene aldehydes from *Bupleurum gibbraltarium*, J. Nat. Prod. 61 (1998) 506–507, <https://doi.org/10.1021/np960732y>.
- [19] F. Bohlmann, C. Zdero, Über terpenoide aus *Ferula hispanica*, Chem. Ber. 102 (1969) 2211–2215, <https://doi.org/10.1002/cber.19691020708>.
- [20] J. Lemmich, P.A. Pedersen, M.S. Sood, B.E. Nielsen, Constituents of Umbelliferous plants. XVI. Terpenoids from the root of *Selinum carvifolium* L., Acta Chem. Scand. 25 (1971) 995–999.
- [21] F. Bohlmann, M. Grenz, Über weitere terpenoide aus umbelliferen, Tetrahedron Lett. (1970) 1453–1456, [https://doi.org/10.1016/S0040-4039\(01\)97993-6](https://doi.org/10.1016/S0040-4039(01)97993-6).
- [22] F. Bohlmann, G. Weickgenannt, Synthese der *Ferula*-terpene, Chem. Ber. 107 (1974) 1769–1772.
- [23] R. Negri, Polyacetylenes from terrestrial plants and fungi: recent phytochemical and biological advances, Fitoterapia 106 (2015) 92–109, <https://doi.org/10.1016/j.fitote.2015.08.011>.
- [24] L. Hansen, P.M. Boll, Polyacetylenes in Araliaceae: their chemistry, biosynthesis and biological significance, Phytochemistry 25 (1986) 285–293, [https://doi.org/10.1016/S0031-9422\(00\)85468-0](https://doi.org/10.1016/S0031-9422(00)85468-0).
- [25] Y. Chen, S. Peng, Q. Luo, J. Zhang, Q. Guo, Y. Zhang, X. Chai, Chemical and pharmacological progress on polyacetylenes isolated from the family Apiaceae, Chem. Biodivers. 12 (2015) 474–502, <https://doi.org/10.1002/cbdv.201300396>.

- [26] C. Zidorn, K. Jöhrer, M. Ganzera, B. Schubert, E.M. Sigmund, J. Mader, R. Greil, E.P. Ellmerer, H. Stuppner, Polyacetylenes from the Apiaceae vegetables carrot, celery, fennel, parsley, and parsnip and their cytotoxic activities, *J. Agric. Food Chem.* 53 (2005) 2518–2523, <https://doi.org/10.1021/jf048041s>.
- [27] L. Schmiech, C. Alayrac, B. Witulski, T. Hofmann, Structure determination of bisacetylenic oxylipins in carrots (*Daucus carota* L.) and enantioselective synthesis of faltarindiol, *J. Agric. Food Chem.* 57 (2009) 11030–11040, <https://doi.org/10.1021/jf9031475>.
- [28] H.Q. Duan, Y. Takaishi, Y. Fujimoto, C. Garzon, C. Osorio, C. Duque, Chemical constituents from the Colombian medicinal plant *Niphogeton ternate*, *Chem. Pharm. Bull.* 50 (2002) 115–117, <https://doi.org/10.1248/cpb.50.115>.
- [29] E. Lemmich, The absolute configuration of the acetylenic compound faltarindiol, *Phytochemistry* 20 (1981) 1419–1420, [https://doi.org/10.1016/0031-9422\(81\)80053-2](https://doi.org/10.1016/0031-9422(81)80053-2).
- [30] M.W. Bernart, J.H. Cardellina, M.S. Balaschak, M.R. Alexander, R.H. Shoemaker, M.R. Boyd, Cytotoxic faltarinol oxylipins from *Dendropanax arboreus*, *J. Nat. Prod.* 59 (1996) 748–753, <https://doi.org/10.1021/np960224o>.
- [31] M. Kobaisy, Z. Abramowski, L. Lermer, G. Saxena, R.E.W. Hancock, G.H.N. Towers, Antimycobacterial polyynes of Devil's Club (*Oplopanax horridus*), a North American native medicinal plant, *J. Nat. Prod.* 60 (1997) 1210–1213, <https://doi.org/10.1021/np970182j>.
- [32] D. Lechner, M. Stavri, M. Oluwatuyi, R. Pereda-Miranda, S. Gibbons, The anti-staphylococcal activity of *Angelica dahurica* (Bai Zhi), *Phytochemistry* 65 (2004) 331–335, <https://doi.org/10.1016/j.phytochem.2003.11.010>.
- [33] M.C. Yang, H.C. Kwon, Y.-J. Kim, K.R. Lee, H.O. Yang, Oploxynes a and B, polyacetylenes from the stems of *Oplopanax elatus*, *J. Nat. Prod.* 73 (2010) 801–805, <https://doi.org/10.1021/np900628j>.
- [34] G. Zheng, W. Lu, J. Cai, Stereoselective total synthesis of (3R,8S)-faltarindiol, a common polyacetylenic compound from Umbellifers, *J. Nat. Prod.* 62 (1999) 626–628, <https://doi.org/10.1021/np980418z>.
- [35] A.S. Ratnayake, T. Hemscheidt, Olefin cross-metathesis as a tool in natural product degradation. The stereochemistry of (+)-faltarindiol, *Org. Lett.* 4 (2002) 4667–4669, <https://doi.org/10.1021/ol027033z>.
- [36] S. Tamura, T. Ohno, Y. Hattori, N. Murakami, Establishment of absolute stereostructure of faltarindiol, algicidal principle against *Heterocapsa circularisquama* from *Notopterygii* rhizome, *Tetrahedron Lett.* 51 (2010) 1523–1525, <https://doi.org/10.1016/j.tetlet.2010.01.047>.
- [37] D.V. Kuklev, A.J. Domb, V.M. Dembitsky, Bioactive acetylenic metabolites, *Phytomedicine* 20 (2013) 1145–1159.
- [38] J.F. Young, L.P. Christensen, P.K. Theil, N. Oksbjerg, The polyacetylenes faltarinol and faltarindiol affect stress responses in myotube cultures in a biphasic manner, *Dose – Response* 6 (2008) 239–251.
- [39] H.R. Jin, J. Zhao, Z. Zhang, Y. Liao, C.-Z. Wang, W.-H. Huang, S.-P. Li, T.-C. He, C.-S. Yuan, W. Du, The antitumor natural compound faltarindiol promotes cancer cell death by inducing endoplasmic reticulum stress, *Cell Death Dis.* 3 (2012) e376, <https://doi.org/10.1038/cddis.2012.122>.
- [40] T. Lu, M. Gu, Y. Zhao, X. Zheng, C. Xing, Autophagy contributes to faltarindiol-induced cell death in breast cancer cells with enhanced endoplasmic reticulum stress, *PLoS ONE* 12 (2017) e0176348, <https://doi.org/10.1371/journal.pone.0176348>.
- [41] T.-J. Kim, H.-S. Kwon, M. Kang, H.H. Leem, K.-H. Lee, D.-Y. Kim, The antitumor natural compound faltarindiol disrupts neural stem cell homeostasis by suppressing notch pathway, *Int. J. Mol. Sci.* 19 (2018) e3432, <https://doi.org/10.3390/ijms19113432>.
- [42] W.H. Huang, L. Shao, C.Z. Wang, C.S. Yuan, H.H. Zhou, Anticancer activities of polyynes from the root bark of *Oplopanax horridus* and their acetylated derivatives, *Molecules* 19 (2014) 6142–6162, <https://doi.org/10.3390/molecules19056142>.