

Science 11 Dec 2020:

Vol. 370, Issue 6522, pp. 1348-1352

DOI: 10.1126/science.abd2115

<https://science.sciencemag.org/content/370/6522/1348>

**Southern Ocean upwelling, Earth's obliquity, and glacial-interglacial
atmospheric CO₂ change**

Authors: Xuyuan E. Ai^{1,2*}, Anja S. Studer³, Daniel M. Sigman¹, Alfredo Martínez-García²,
Francois Fripiat⁴, Lena M. Thöle⁵, Elisabeth Michel⁶, Julia Gottschalk⁷, Laura Arnold⁸,
Simone Moretti², Mareike Schmitt², Sergey Oleynik¹, Samuel L. Jaccard^{9†}, Gerald H. Haug².

Affiliations:

¹Department of Geosciences, Princeton University, Princeton, NJ 08544, USA.

²Climate Geochemistry Department, Max Planck Institute for Chemistry, Mainz 55128,
Germany.

³Department of Environmental Sciences, University of Basel, Basel.

⁴Department of Geosciences, Environment and Society, Université Libre de Bruxelles,
Brussels, Belgium.

⁵Department of Earth Sciences, Utrecht University, Utrecht, the Netherlands.

⁶Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Gif-sur-Yvette, France.

⁷Lamont-Doherty Earth Observatory, Palisades, NY 10964, USA.

⁸Department of Earth Sciences, ETH Zurich, Zurich, Switzerland.

⁹Institute of Geological Sciences & Oeschger Center for Climate Change Research,
University of Bern, 3012 Bern, Switzerland.

[†]now at Institute of Earth Sciences, University of Lausanne, Switzerland

*Correspondence to: xuyuana@princeton.edu.

Abstract:

Previous studies suggest that during the late Pleistocene ice ages, surface-deep exchange was somehow weakened in the Southern Ocean's Antarctic Zone, which reduced the leakage of deeply sequestered carbon dioxide and thus contributing to the lower atmospheric carbon dioxide levels of the ice ages. Here, high-resolution diatom-bound nitrogen isotope measurements from the Indian sector of the Antarctic Zone reveal three modes of change in Southern Westerly Wind-driven upwelling, each affecting atmospheric carbon dioxide. Two modes, related to global climate and the "bipolar seesaw", have been proposed previously. The third mode – which arises from the meridional temperature gradient as affected by Earth's "obliquity" (axial tilt) - can explain the lag of atmospheric CO₂ behind climate during glacial inception and deglaciation. This obliquity-induced lag, in turn, makes carbon dioxide a delayed climate amplifier in the late Pleistocene glacial cycles.