南極資料

2019-2020年ベルギー南極観測隊による東南極ナンセン氷原における隕石探査報告

--Manuscript Draft--

受付番号:	AR-D-20-00006R2
論文種別:	報告（report）
標題:	2019-2020年ベルギー南極観測隊による東南極ナンセン氷原における隕石探査報告
標題（英語）:	Detailed record of the BELARE 2019-2020 meteorite recovery expedition on the Nansen Ice Field, East Antarctica
ランニングタイトル:	2019-2020年ベルギー南極観測隊による隕石探査
ランニングタイトル（英語）:	Detailed record of the BELARE 2019-2020 meteorite recovery expedition
筆頭著者:	Goderis Steven
全著者:	Goderis Steven
	Yesiltas Mehmet
	Pourkhorsandi Hamed
	白井 直樹, 博士
	Poudelet Manu
	Leitl Martin
	山口 亮
	Debaille Vinciane
	Claeys Philippe
責任著者:	白井 直樹, 博士 東京都立大学 八王子市, 東京都 JAPAN
抄録:	2019-2020年の夏期に，東南極セール・ロンダーネ山地南部においてベルギー南極観測隊 (BELARE) により隕石探査を実施した．ナンセン氷原には，2020年1月15日から2月6日まで23日間滞在し，採取した隕石の総数は66個，合計重量は約8kgであった．ナンセン氷原での隕石集積機構を解明するために，隕石の他に氷，火山灰層や岩石の破片も採取した．採取した隕石は，凍結したまま国立極地研究所に輸送された．これら採取した隕石が国際隕石学会の隕石命名委員会に認可された後，分類データはMeteorite Newsletterで公開される．
抄録（英語）:	This report summarizes the Belgian Antarctic Expedition (BELARE) 2019-2020 meteorite search and recovery expedition near the Sør Rondane Mountains of East Antarctica during the 2019-2020 field season. This expedition took place from 15 January to 6 February 2020 within the area defined as “C” of the Nansen Ice Fields (S72°38’-72°48’S, 24°35’-25°06’E). The expedition team consisted of four scientists and two field guides, who systematically searched the ice field area and collected 66 meteorites. The total weight of the meteorites was determined to be ~8 kg. In addition to meteorites, blue ice samples, volcanic ash layers, and wind-blown terrestrial rock fragments were collected from the area to study in detail the nature of the mechanisms concentrating meteorites on the Nansen Ice Fields. The recovered meteorites were transported in frozen state to the National Institute of Polar Research, Japan for dry-thawing and subsequent classification. The newly collected meteorites will be presented to the Meteorite Nomenclature Committee of the Meteoritical Society for approval, after which their classification will be published in the Meteorite Newsletter.
分野:	900: 隕石（Meteoritics）
キーワード:	隕石探査；南極隕石；ナンセン氷原
キーワード (英語):	Meteorite search; Antarctic meteorite, Nansen Ice Field

査読者への回答

Dear editor,

We truly appreciate the comments from editor. In revising our report, comments are all considered. Let us describe here how we responded to them one by one. The revised portions are shown in blue in the revised manuscript. We believe that our responses satisfactorily meet the comments given by a editor.

編集者のコメント：

前回査読者からしてきされた点は修正が行われていますが、若干のミス（日付の記述の統一など）が本文にありますので微修正を行ってください。その他、気が付いた微細な点を修正しましたので、著者側で再度確認の上、必要に応じて修正を行ってください。微修正となりますので対応よろしくお願いします。

line50, line391

The expedition team consisted of four scientists and two field guides,

数字ではなく2章のように単語に統一してはいかが？

Expedition members and their roles

The BELAM 2019-2020 expedition team consisted of four researchers in the fields of meteoritics, planetary sciences, or cosmochemistry, and two field guides, trained as medical doctor or paramedic. All six BELAM 2019-2020 team members and their roles are listed in Table 2.

Response: We modified it.

line51

a total of は不要では？

collected 66 meteorites

Response: We modified it.

line152

layoverでは？

Response: We modified it.

line165

Antarctic cold environmentではなく、cold Antarctic environmentでは？

Response: We modified it.

line187

airstripでは？

Response: We modified it.

line190

snowmobile suitesではなく、snowmobile suitsでは？

Response: We modified it.

line193-4
at Cape Town or PEAでは？
Response: We modified it.

line242-245
この部分の日付の表記が統一されていない。例えば、13 Februaryと13th of February。
(13 to 14 January and 6 to 13 February), and 22 days were spent on the Nansen Ice Field (15 January to 6 February, Table 4). Following the expedition, the BELAM 2019-2020 team left PEA at 8:00 UTC+1 on the 12th of February 2020, departed from Perseus on an Ilyushin-76 airplane, and arrived at Cape Town at 15:00 UTC+2 on 13th of February 2020
Response: We modified it throughout the revised manuscript.

line277
setupではなくset upでは？
Response: We modified it.

line303
17:00ｐｍからｐｍを削除
Response: We modified it.

line327
existent databaseではなくexisting databaseでは？
Response: We modified it.

Detailed record of the BELARE 2019-2020 meteorite recovery expedition on the Nansen Ice Field, East Antarctica

Steven Goderis[1], Mehmet Yesiltas[2], Hamed Pourkhorsandi[3], Naoki Shirai[4*], Manu Poudelet[5], Martin Leitl[5], Akira Yamaguchi[6], Vinciane Debaille[3], Philippe Claeys[1]

2019-20202 年ベルギー南極観測隊による東南極ナンセン氷原における隕石探査報告

Steven Goderis[1], Mehmet Yesiltas[2], Hamed Pourkhorsandi[3], 白井直樹 [4*], Manu Poudelet[5], Martin Leitl[5], 山口亮 [6], Vinciane Debaille[3], Philippe Claeys[1]

[1] ブリュッセル自由大学. Analytical-, Environmental-, and Geo-Chemistry, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium.

[2] クルクラーレリ大学. Faculty of Aeronautics and Space Sciences, Kirklareli University, Kirklareli, Turkey 39100.

[3] ブリュッセル自由大学. Laboratoire G-Time, Université Libre de Bruxelles, CP 160/02, 50, Av. F.D. Roosevelt, 1050 Brussels, Belgium.

[4] 東京都立大学理学研究科化学専攻. Department of Chemistry, Tokyo Metropolitan

University, 1-1 Minamiosawa, Hachioji, Tokyo 192-0397.

[5] 国際極地基金. International Polar Foundation, Rue des vétérinaires, 42c/1 1070, Brussels, Belgium.

[6] 国立極地研究所. National Institute of Polar Research, 10-3 Midoricho, Tachikawa, Tokyo 190-8518.

＊ Corresponding author. E-mail: shirai-naoki@tmu.ac.jp

要旨：2019-2020 年の夏期に，東南極セール・ロンダーネ山地南部においてベルギー南極観測隊 (BELARE) により隕石探査を実施した．ナンセン氷原には，2020 年 1 月 15 日から 2 月 6 日まで 23 日間滞在し，採取した隕石の総数は 66 個，合計重量は約 8kg であった．ナンセン氷原での隕石集積機構を解明するために，隕石の他に氷，火山灰層や岩石の破片も採取した．採取した隕石は，凍結したまま国立極地研究所に輸送された．これら採取した隕石が国際隕石学会の隕石命名委員会に認可された後，分類データは Meteorite Newsletter で公開される．

キーワード：隕石探査, 南極隕石, ナンセン氷原

Abstract: This report summarizes the Belgian Antarctic Expedition (BELARE) 2019-2020 meteorite search and recovery expedition near the Sør Rondane Mountains of East Antarctica during the 2019-2020 field season. This expedition took place from 15 January to 6 February 2020 within the area defined as “C” of the Nansen Ice Fields (S72°38’-72°48’S, 24°35’-25°06’E). The expedition team consisted of four scientists and two field guides, who systematically searched the ice field area and collected 66 meteorites. The total weight of the meteorites was determined to be ~8 kg. In addition to meteorites, blue ice samples, volcanic ash layers, and wind-blown terrestrial rock fragments were collected from the area to study in detail the nature of the mechanisms concentrating meteorites on the Nansen Ice Fields. The recovered meteorites were transported in frozen state to the National Institute of Polar Research, Japan for dry-thawing and subsequent classification. The newly collected meteorites will be presented to

the Meteorite Nomenclature Committee of the Meteoritical Society for approval, after which their classification will be published in the Meteorite Newsletter.

Keywords: Meteorite search, Antarctic meteorite, Nansen Ice Field

1. Introduction

The Sør Rondane Mountains and blue ice fields surrounding these mountains in East Antarctica have previously been visited by research expedition teams to collect meteorites. One of the first expeditions to East Antarctica was JARE-10, the 10th Japanese Antarctic Research Expedition, which recovered 9 meteorites from the ice fields surrounding the Yamato Mountains (Yoshida *et al.,* 1971). The meteorites collected around the Sør Rondane Mountains (mainly from the Nansen and Balchen Ice Fields) are called Asuka meteorites. The first Asuka meteorites were found by accident in 1986 by a glaciological party of JARE-27 on bare ice fields near Mt. Balchen (Nishio *et al.,* 1987). During the 1987-1988 and 1988-1989 field seasons, JARE-29 and JARE-30 team members visited the Nansen and Balchen Ice Fields and recovered nearly 2000 meteorites (Naraoka *et al.,* 1990; Yanai, 1993). Following this expedition, the nomenclature committee of the Meteoritical Society designated this region as an official meteorite dense collecting area (DCA), designated Asuka after the former Japanese research station in this area. All collected meteorites from this DCA are defined by the name Asuka (abbreviated as A), followed by the year of discovery and a number (e.g., A 87251). During the JARE-51 in the 2009–2010 field season with participation of a Belgian researcher and logistical support from the Belgian Princess Elisabeth Antarctica (PEA) station, a total of 635 meteorites were recovered from the Mt.

78 Balchen area (Kaiden *et al.,* 2010). By systematically searching various sections of Nansen Ice
79 Fields, the JARE and Belgian Antarctic Expedition (BELARE) joint expeditions recovered a total
80 of 908 Asuka meteorites in the 2010-2011 and 2011-2012 seasons (Goderis *et al.,* 2011; Tsuchiya
81 *et al.,* 2012; Imae *et al.,* 2015). The Asuka meteorites include several large specimens such as A
82 87251 (46 kg) and A 12389 (18 kg), rare meteorites such as angrites (A 881371, A 12209), a
83 lunar meteorite (A 881757), a shergottite (A 12325), unique carbonaceous chondrites (CH3: A
84 881020, A 881541, A 881691), a Rumuruti chondrite (A 881988), mesosiderites (A 87106, A
85 881154, and A 882023), 19 diogenites, 11 ureilites, 2 acapulcoites, and 2 ungrouped
86 carbonaceous chondrites. During the 2018-2019 Antarctic season, 3 meteorites (A 18001, A
87 18002, and A 18003), classified as ordinary chondrites, were recovered from the area A of
88 Nansen Ice Field by MY, as part of the 3rd Turkish Antarctic Expedition (TAE-III) (Yesiltas *et*
89 *al.,* 2019). All meteorite search expeditions in the vicinity of the Sør Rondane Mountains as well
90 as the current repositories for the recovered meteorites are summarized in Table 1.

91 (Table 1 could be here)

92 In this report, we report in detail on the BELARE 2019-2020 meteorite search and recovery
93 expedition (BELAM), which took place between the 15 January and the 6 February 2020 within
94 the area C of Nansen Ice Field and in which a Turkish scientist participated as part of the TAE-
95 IV. The expedition team recovered 66 meteorites, totaling ~8 kg. The collected meteorites
96 include carbonaceous chondrites, ordinary chondrites, and achondrites (notably HED meteorites),
97 as identified in the field. These meteorites will be referred to as Asuka 19 meteorites. The
98 average weight of each meteorite is ~120 g. The collected meteorites were securely packaged and
99 transported in frozen state to the National Institute of Polar Research (NIPR) in Tokyo, Japan. At

100 the NIPR, the meteorites are dry-thawed in a controlled environment in order to avoid any
101 contamination due to melting of ice/snow. Following the thawing process, each meteorite will be
102 shared between Japan and Belgium. In addition, the expedition team members have priority
103 access to the recovered meteorites for scientific research, as specified in the Memorandum of
104 Understanding (MoU) signed by all parties involved.

105

106 **2. Expedition members and their roles**

107 The BELAM 2019-2020 expedition team consisted of four researchers in the fields of
108 meteoritics, planetary sciences, or cosmochemistry, and two field guides, trained as medical
109 doctor or paramedic. All six BELAM 2019-2020 team members and their roles are listed in Table
110 2.

111 (Table 2 could be here)

112 **3. Preparation for the expedition**

113 *3.1. Pre-expedition meetings in Brussels and Cape Town:* On 8 May 2019, BELARE 2019-2020
114 members (including MY, VD, and SG) met for the first time in Brussels at the headquarters of
115 the International Polar Foundation (IPF) with the scientists and logistics staff to receive and
116 exchange general information on BELARE 2019-2020, including the number of scientific teams,
117 field guides, preliminary traveling dates, and the registration process for participation in the
118 expedition. With regard to the BELAM 2019-2020 campaign, the length of the expedition to the
119 Nansen Ice Field and the logistical support required were first discussed elaborately. This general

meeting was preceded by shorter meetings between MY, SG, VD, and PC on 7 May in 2020 at both the Vrije Universiteit Brussel (VUB) and Université Libre de Bruxelles (ULB).

A second BELARE 2019-2020 meeting was organized at the IPF on 20 September 2019 (attended by HP and SG), with more detailed information regarding the various scientific projects, field guides, and cargo shipping. Particular attention was paid to the BELAM program and logistics, which included detailed discussion with Alain Hubert of the IPF on the targeted search area on the blue ice fields.

The first BELAM 2019-2020 team meeting was organized in Cape Town, South Africa on 6 January 2020, after the arrival of all the scientific team members on 5 January. During various meetings in the following days, the schedule of the expedition was discussed after studying the targeted blue ice area of Nansen C (Figure 1-d) and considering the details presented by Imae *et al.* (2015). The main objective of recovering meteorites and sampling blue ice to better constrain the meteorite concentration mechanisms of the Nansen Ice Fields as a whole was discussed in great detail.

3.2. Dronning Maud Land Air Network (DROMLAN) flights: The Belgian, Japanese, and Turkish team members flew to Cape Town, South Africa independently using commercial airlines, from where DROMLAN operates flights to and from Antarctica. The flight schedules for outgoing (and return) flights from Cape Town to the Novolazarevskaya (Novo) Air Base (using a Boeing 757) and then to PEA Station (using a Basler Turbo) were arranged by the IPF staff members. The Antarctic Logistics Centre International (ALCI) briefing was organized on 8 January at

10:00 (UTC+2) to inform on general flight procedures, cargo requirements, polar clothing, and the general infrastructure at Novo.

3.3. Itinerary: While the first departure on the Boeing 757 took place on 9 January (UTC+2), the airplane returned to Cape Town after approximately 3.5h of flight (reaching the point-of-no-return) due to (suddenly developed) poor weather conditions at Novo (total of 7 hours flight). The next and successful departure took place from Cape Town on 13 January at 10:00 (UTC+2) on an Ilyushin-76 airplane. After two hours of layover at Novo, the consecutive flight to PEA took place using a Basler Turbo airplane, which arrived at PEA on 13 January at 19:00 (UTC+1). A general discussion of the BELARE 2019-2020 scientific activities during the second part of the Antarctic season was organized on 14 January at 9:00 (UTC+1), led by Alain Hubert and other members of the IPF scientific and logistics staff. Details of the meteorite search and recovery expedition were further discussed during this meeting to ensure a safe and successful field campaign.

3.4. Safety trainings: After arrival from Novo and before departure to the Nansen Ice Field, the pre-expedition safety training for the BELAM 2019-2020 team members took place near PEA. On 14 January, all team members participated in crevasse rescue training, by descending into a well-studied crevasse near PEA (S71.9170˚, E23.5885˚) and constructing a pulley-string system to extract each team member out of the crevasse in rotation. Before and after this crevasse rescue training, technical details regarding the use of crampons as well as snowmobile driving and maintenance were reviewed. The team participated during the following afternoon and evening in presentation sessions on medical issues related to the cold Antarctic environment and high altitude (including first aid, rescue training, hypothermia, frostbite, and altitude sickness).

Additional training and explanations were provided by the field guides once the team arrived at the base camp (area C of Nansen Ice Field).

3.5. Logistics in Antarctica: At the PEA station, most of the expedition necessities had already been prepared by ML (fuel, containers, snowmobile, and communication devices) and MP (food, general equipment, and medical kits) between 5 and 10 January 2020. As a result, BELAM 2019-2020 team members and IPF support team (Alain Hubert, Gigi Johnson-Amin, Louis Greindl, Pierre Dumont) were able to leave PEA for the Nansen Ice Field on 15 January 2020 at 9:30 (UTC+1). After 13 hours of travel, the convoy of two large snow vehicles (Prinoth snow tractors), each pulling 3 Lehmann sledges carrying living modules, fuel, food, and extra snowmobiles (Ski-Doo, BRP), arrived on the Nansen C Ice Field (at 22:30, UTC+1). Upon arrival, the "Nansen C" base camp (BC) was set up (S72.79754°, E24.86002°; elevation 3100 m) and the IPF support team returned to PEA with the two Prinoth snow tractors, one living module and an empty Lehmann sledge on 17 January. The team planned to spend approximately 3 weeks at the BC to search for meteorites within area C, based on the considered search area (Figure 1-d). After completion of the meteorite recovery mission, the return journey to PEA took place on the 6th of February. Upon returning to PEA, the team prepared to travel outbound, for which they first moved to the Perseus landing and departure airstrip near Romnœsfjellet. The flight to Cape Town took place on the 13th of February. The full BELAM 2019-2020 team arrived at their home countries in the period of 16 to 18 February 2020.

3.6. Clothing: Outer clothes that were used during the meteorite search (snowmobile suits, snow boots, helmets, and crampons) were borrowed from the IPF, from the stock kept either at Cape

183 Town or PEA. Other items such as base layer clothing, goggles, gloves, and balaclavas were
184 prepared by the individual team members.

185 *3.7. Snowmobiles*: The fuel requirement for snowmobiles was estimated based on the following
186 assumptions: 6 snowmobiles traveling 50 km/day during 20 days with a fuel consumption of 3.5
187 km/liter/snowmobile requires ~1715 liters of fuel. The expedition foresaw a total of 1880 liters of
188 fuel for the field work. The snowmobiles also require engine oil to operate. Oil consumption was
189 estimated to be ~20 liters/snowmobile during the period of 20 days, for which a total of ~440
190 liters of engine oil was stored for use throughout the expedition.

191 *3.8. Heater*: A total of 280 liters of JET-A1 fuel was brought to the base camp for the heating
192 system (Webasto).

193 *3.9. Food*: Twenty loaves of bread was estimated for 20 days to be consumed at breakfast. For
194 lunch, 120 packs of instant soups and noodles, and 360 chocolate bars were brought. One
195 hundred and twenty frozen meals for dinner were prepared at PEA. These numbers were
196 estimated as for 6 persons per day during 20 expedition days. Plenty of fresh and dried fruits,
197 vegetables, snacks, and nuts were also brought to the base camp.

198

199 **4. Scientific matters**

200 *4.1. Definition, division, and subdivision of search areas:* Following previous expeditions to the
201 Nansen Ice Field, three areas have been defined for meteorite searches on the Nansen Ice Field:
202 areas A, B, and C (Figure 1-d). Areas A and B were previously searched by JARE-29 in 1987-
203 1988 and 1988-1989 as well as BELARE 2010-2011 and JARE-54/BELARE 2012-2013

204 expeditions. Area C had been previously searched during JARE-29, while only the northern part
205 had been covered by JARE-54/BELARE 2012-2013. The remaining southern part of Nansen area
206 C had thus not been searched in over 30 years. During this expedition, 3 weeks were dedicated to
207 Nansen area C to cover the entire field for searching meteorites and collecting blue ice samples.
208 A short excursion to Nansen area B also took place at the end of the 3 weeks to search for
209 meteorites and terrestrial rocks in a previously encountered moraine. Another day trip allowed to
210 collect additional ice samples from area A, which will be used to extend (or not) previously
211 observed spatial patterns in the H and O isotopic compositions of Nansen area B across the entire
212 Nansen Ice Field (Zekollari *et al.*, 2019).

213 *4.2. Expected number of meteorite finds:* The expedition by JARE-29 in 1987-1989 collected 573
214 meteorites from area A, 698 from area B, and 311 from area C (Naraoka *et al.*, 1990; Imae *et al.*,
215 2012). During the BELARE 2010-2011, 218 meteorites were collected from area A (Goderis *et*
216 *al.*, 2011), while 368 meteorites were recovered from area B and 56 from area C during JARE-
217 54/BELARE 2012-2013 (Debaille *et al.*, 2013; Imae *et al.*, 2015). Assuming similar blue ice
218 conditions and extraction efficiencies for each expedition, an approximate number of meteorites
219 to be recovered was estimated based on the following ratios between the different areas and
220 expeditions (i.e., 573:698:311=218:368:x), where x was the expected number of meteorites to be
221 recovered from area C during the 2019-2020 season. Therefore, the expected number of
222 meteorites for area C ranged from 118 to 164. Taking into account the number of meteorites that
223 had already been recovered from the northern part of area C during JARE-54/BELARE 2012-
224 2013, the expected number of meteorites for the remaining part of Nansen C ranged from 62 to
225 108.

5. Expedition log

A typical daily schedule on the Nansen Ice Field is summarized in Table 3. The duration of the BELAM 2019-2020 expedition was 31 days (Table 4). Of the 31 days in Antarctica, 9 days were spent at PEA (13 to 14 January and 6 to 13 February), and 22 days were spent on the Nansen Ice Field (15 January to 6 February, Table 4). Following the expedition, the BELAM 2019-2020 team left PEA at 8:00 UTC+1 on 12 February 2020, departed from Perseus on an Ilyushin-76 airplane, and arrived at Cape Town at 15:00 UTC+2 on 13 February 2020 after spending a single night at Novo.

(Figure 1, and Tables 3 and 4 could be here)

6. Nansen Ice Field

6.1. Base camp: The location of our base camp (S72.79754°, E24.86002°) was determined by Alain Hubert during his reconnaissance trip. The living quarters were arranged in a straight line so that each solar panel mounted on the living quarters faced the sun. The arrangement of the three living quarters is shown in Figure 2. Snowmobiles were parked about 20 m north of the living quarters. Snow drift that developed at the camp site is clearly visible in Figure 2.

(Figure 2 could be here)

6.2. Evening briefings: Every evening, activities performed during the day were reviewed by the team members. Magnetic susceptibility of the newly collected meteorites was determined. All meteorite images, track lines, and coordinates were gathered. These geographical data were

246 imported in the QGIS-Quantarctica (Matsuoka *et al.,* 2008) and Garmin BaseCamp software
247 packages, and maps of the visited regions were produced. The weather forecast for the next few
248 days was also discussed (Section 6.3). Then, the next day was planned based on discussions
249 between all team members.

250

251 *6.3. Weather observations:* Weather conditions in the Nansen Ice Field were measured three
252 times a day using a handheld weather station (Kestrel-4500), before and after the breakfast (at
253 7:30 and 9:00 UTC+1, respectively) as well as in the evening (at 21:30 UTC+1). Temperature
254 with and without the windchill effect, wind speed, humidity, and pressure was recorded. Based
255 on the collected weather data, we observed that absolute temperature varied between -26.5°C and
256 -18.6°C, which corresponds to -47.1°C and -31.8°C when the windchill effect is taken into
257 account (Figure 3, Table 5). Generally, wind and snow drift were experienced every day. Wind
258 speeds ranged between 5.5 and 13.4 m/s. The humidity level varied between 44.3 and 100%,
259 while the atmospheric pressure ranged between 984.9 and 1009.5 hPa. The weather observation
260 log is given in Table 5. Weather conditions were ideal for meteorite search for 12 days out of the
261 22 days spent on the Nansen Ice Field. During the other days, strong winds caused snow drifts,
262 which reduced visibility and prevented any field work.

263 (Figure 3 and Table 5 could be here)

264 *6.4. Meteorite search:* Our search mostly focused on the Nansen Ice Field C (Figure 1-d).
265 Therefore, our base camp was set up in this region. Every morning, after breakfast and the
266 weather observation, the departure time for the field work was decided. In the case of good

267 weather, the usual departure time was around 10:00. After each snowmobile was checked and
268 prepared, the team members left the base camp following the guide (MP) in a straight line with
269 an appropriate distance from each other that depended on the visibility. After arrival at the
270 targeted search area, the team members took their designated position, forming a V-shaped
271 orientation with each member in his fixed position and one field guide (MP) at the apex of the V-
272 formation (Figure 4), while the other field guide (ML) was positioned at one end of the V-shape.
273 The field guide then followed predetermined GPS tracks in a zigzag pattern. Following previous
274 campaigns, working and searching in this pattern provides the highest surface coverage as well as
275 maximum safety.

276 (Figure 4 could be here)

277 When a meteorite was found, previously defined procedures were followed (Goderis *et al.*, 2011;
278 Imae *et al.*, 2015). First, the find was reported to other members over the radio. Then, the field
279 sample number consisting of the initial of the finder's last name, the year, month, day, and the
280 number of the meteorite found on that day by the member (e.g., Y20012501) was written with a
281 marker pen on the surrounding ice/snow near the meteorite and a scale was placed beside it.
282 Pictures of the meteorite (with a scale) were taken from various angles (Figure 5). The meteorite
283 was then picked up using zip lock polyethylene bags while avoiding any direct contact. In case
284 carbonaceous chondrites were identified in the field, these meteorites were placed in Teflon bags
285 instead of regular polyethylene zip lock bags due to possible contamination of organic
286 compounds from polyethylene zip lock bags. After sealing the zip lock bag, it was placed in a
287 larger field bag. Finally, the location of the recovered meteorite was recorded using a handheld
288 GPS unit. In order to save time, these steps were usually performed with the help of another

member of the field campaign. Return time to the base camp was determined based on the weather conditions. In good conditions, this typically was around 17:00 (UTC+1). After 17:00 (UTC+1), even during good weather conditions, the angle of the sunlight and its reflectance from the surface of the blue ice made the search strenuous, lowering the effectiveness of the systematic searches.

(Figure 5 could be here)

6.5. Mass distribution: All recovered meteorites (n=66) were weighted upon returning to the PEA station. Although the samples were weighted within their bags, the weight of empty bags was later subtracted from the initial measurements. The total mass of the collected meteorites was calculated to be ~8280 g, with individual specimens range from 1 to 889 g. The average, median, and mode weights were ~125, 58, and 30 g, respectively. Figure 6-left summarizes the mass distribution of the collected samples. Meteorites with masses between 10 to 50 g exhibit the highest relative abundance (34.85%, n=23), while those heavier than 500 g are the least common ones (4.55%, n=3).

6.6. Magnetic measurements: Magnetic susceptibilities of the recovered meteorites were measured on whole samples using a handheld A*METMET susceptibility meter developed at CEREGE (Aix-en-Provence) by Dr. M. Uehara and Dr. J. Gattacecca. Magnetic susceptibility is expressed as the decimal logarithm of χ in 10^{-9} m^3/kg in order to account for the five orders of magnitude variation in rocks. For strongly magnetic material ($\log \chi > 3$), χ is proportional to the amount of metal, magnetite, maghemite, cohenite, and schreibersite, i.e., minerals with practically equal specific χ. A combination of magnetic susceptibility data with microscopic

observations is an effective way to classify ordinary chondrites and evaluate pairing (Pourkhorsandi *et al.*, 2019). Comparison of in-situ magnetic susceptibility measurements data with the existing database (Folco *et al.*, 2006, Rochette *et al.*, 2003, 2008, 2009) can provide insights on the possible meteorite types. While detailed mineralogical and textural investigation as well as the approval of the meteorite Nomenclature Committee of the Meteoritical Society is needed to narrow down this preliminary classification, the log χ values of the meteorites collected on the Nansen area C Ice Field are presented in Figure 6-right. A more detailed classification of these samples will take place at the NIPR.

(Figure 6 could be here)

7. Problems encountered

7.1. Snowmobiles: Two snowmobiles broke down during the course of the expedition. On the 26th of January, a JARE snowmobile (number 49-1) had to be towed back to the base camp by another snowmobile due to battery failure. Two days later, another snowmobile broke down due to engine failure during the meteorite search and was also towed back to the base camp by another snowmobile.

7.2. Heaters and generators: Each living quarter had a diesel-fueled heater (Webasto heater - Air Top Evo 3900). The heater in the bedroom did not work for unknown reasons, instead an electric-powered heater was used in the bedroom. The day before the end of the expedition, the generator stopped working because of a low battery following failure of the alternator of the generator. This issue was solved by using a power booster/jumper to start the generator.

7.3. Medical issues: Some of the expedition team members suffered from the strong winds during meteorite searches. This resulted in colder than usual fingers and toes. Multiple instances of light frostbite (frostnip) on the face were experienced on windy days. However, severe frostbite did not occur, due to the use of appropriate clothing, chemical warmers and wind-protected helmets.

8. Post-expedition matters

8.1. Retrieval of the base camp: On the evening of 5 February 2020, an IPF support team composed of Alain Hubert and Pierre Dumont arrived at the base camp location with two Prinoth snow tractors to retrieve the meteorite recovery team members and base camp. Prior to their arrival, the interiors of the living quarters and containers were cleaned and prepared by the field guides and the expedition team members for the return. Due to constant snow drift near the base camp, the containers accumulated significant amounts of snow on the leeward side. The accumulated snow had to be removed by hand and using the Prinoth vehicles to free the containers from accumulated and compacted snow. Next, all containers were disassembled and separated from each other. Subsequently, these containers were attached to the Prinoth trucks for the return trip. After departing from the Nansen Ice Field at 16:00 (UTC+1), all containers and equipment, along with the expedition team members, who travelled inside one of the living quarters, arrived at the PEA station at 4:00 (UTC+1) on 7 February 2020.

8.2. Return of food supplies and equipment: The returned containers and living quarters were emptied by the field guides upon arrival to the PEA station. All remaining food and used equipment were returned to the PEA station.

355

356 *8.3. Storage and transportation of the meteorites:* All individually packed meteorites were stored
357 in larger zip lock and cotton bags, which were stored inside a cool box placed outside of the
358 containers throughout the expedition (Figure 7). Although the NIPR was not directly involved in
359 the field collection of meteorites in this expedition, the MoU defined during the previous
360 expeditions since 2009-2010 was upheld and all collected meteorites were sent to NIPR in
361 Tokyo, Japan for defrosting and classification before sharing with the Royal Belgian Natural
362 History Museum (RBINS). As such, NIPR scientists prepared 0.08 mm-thick polyethylene zip
363 lock sample bags of various sizes to store the collected meteorites, similar to what had been done
364 during previous joint expeditions (e.g., Imae *et al.*, 2015): size A (70×50 mm), size C (100×70
365 mm), size E (140×100 mm), size F (170×120 mm), and size K (400×280 mm). In addition, 0.05-
366 mm-thick powder-free polyethylene bags of 40×60 cm and 100×120 cm were used. Due to
367 logistical issues and the outbreak of COVID-19, the collected meteorites were only transported at
368 the beginning of June (departure date from Cape Town on 4 June). In continuously frozen state,
369 the World Courier shipping company transported the meteorites from the Trade Universal storage
370 facilities near the airport in Cape Town to the NIPR in Tokyo, where the package arrived on 11
371 June and was kept in the cold temperature (-30˚C) room until the meteorites were processed.

372 (Figure 7 could be here)

373 **9. Summary**

374 During the 2019-2020 field season, an expedition team consisting of four scientists and two field
375 guides conducted systematic meteorite recovery activities within the previously defined area C of

the Nansen Ice Fields. Sixty-six meteorites with a combined mass of ~8 kg was recovered. This number falls within the range of meteorites (62 to 108) estimated to be collected within Nansen C. These meteorites will be referred to as Asuka 19 meteorites and classified following controlled dry-thawing at the NIPR. Volcanic ash layers and wind-blown terrestrial rock samples as well as blue ice samples were also collected from the Nansen Ice Fields, which will improve our current understanding of the meteorite concentration mechanisms in this area.

Acknowledgments

We gratefully acknowledge the support provided by the IPF at PEA and in the field, with a special thanks to Alain Hubert and Pierre Dumont for their support throughout the expedition. The BELARE 2019-2020 meteorite search and recovery expedition was supported by the Belgian Science Policy (BELSPO) under research grants "AMUNDSEN" and "DIABASE". MY was a member of the TAE-IV, which was carried under the auspices of Presidency of The Republic of Turkey, supported by the Ministry of Industry and Technology, and coordinated by TUBITAK MAM Polar Research Institute.

References

Debaille, V., Imae, N., Yamaguchi, A., Goderis, S., Mikouchi, T., Debouge, W., Hublet, G., Van Roosbroek, N., Zekollari, H., Kojima, H. and Claeys. Ph. (2013): The 2012-2013 joint field campaign for collecting meteorites in Antarctica: an efficient collaboration between Japan

396 and Belgium. Antarctic Meteorites XXXVI. Tokyo, 2013-11-14/15. National Institute of
397 Polar Research, 11–12.

398 Folco, L., Rochette, P., Gattacceca, J. and Perchiazzi, N. (2006): In situ identification, pairing,
399 and classification of meteorites from Antarctica through magnetic susceptibility
400 measurements. Meteoritics & Planetary Science, **41**, 343–353.

401 Goderis, S., Kaiden, H., Debaille, V., Kojima, H. and Claeys, Ph. (2011): Belgian-Japanese
402 search for Antarctic meteorites during the 2010-2011 field season. Antarctic Meteorites
403 XXXIV. Tokyo, 2011-11-17/18. National Institute of Polar Research, 12.

404 Imae, N., Akada, Y., Clayes, Ph., Debaille, V., Goderis, S., Hublet, G., Kojima, H., Martin, C.,
405 Mikouchi, T., Van Roosbroek, N., Yamaguchi, A. and Zekollari, H. (2012): The plan of the
406 search for Antarctic meteorites on the Nansen Ice Fields by the joint expedition between
407 KARE-54 and BELARE 2012-2013. Antarctic Meteorites XXXV. Tokyo, 2012-11-29/30.
408 National Institute of Polar Research, 22-23.

409 Imae, N., Debaille, V., Akada, Y., Debouge, W., Goderis, S., Hublet, G., Mikouchi, T., Van
410 Roosbroek, N., Yamaguchi, A., Zekollari, H., Claeys, Ph. and Kojima, H. (2015): Report of
411 the JARE-54 and BELARE 2012-2013 joint expedition to collect meteorites on the Nansen
412 Ice Field, Antarctica. Nankyoku Shiryô (Antarctic Record), **59**, 38–72.

413 Kaiden, H., Kojima, H. and Goderis, S. (2010): Collection of the Asuka 09 meteorites by the 51st
414 Japanese Antarctic Research Expedition: a preliminary report. Antarctic Meteorites
415 XXXIII. Tokyo, 2010-06-08/09. National Institute of Polar Research, 34–35.

Matsuoka, K., Skoglund, A. and Roth, G. (2018): Quantarctica [Data set]. Norwegian Polar Institute. https://doi.org/10.21334/npolar.2018.8516e961

Naraoka, H., Yanai, K. and Fujita, S. (1990): Report on Antarctic meteorites search around the Sør Rondane Mountains, JARE-29 1988-1989. Nankyoku Shiryô (Antarctic Record), **34**, 216-224 (in Japanese with English abstract).

Nishio, F., Ohmae, H., Mori, K., Osada, K. and Urazuka, S. (1987): Collection of Yamato and Sør Rondane meteorites in the 1986-87 field season, Antarctica. Antarctic Meteorites XII. Tokyo, 1987-06-8/10. National Institute of Polar Research, 1–2.

Pourkhorsandi, H., Gattacceca, J., Rochette, P., D'Orazio, M., Kamali, H., de Avillez, R., Letichevsky, S., Djamali, M., Mirnejad, H., Debaille, V. and Jull, A. J. T. (2019): Meteorites from the Lut Desert (Iran). Meteoritics & Planetary Science, **54**, 1737–1763.

Rochette, P., Sagnotti, L., Bourot-Denise, M., Consolmagno, G., Folco, L., Gattacceca, J., Osete, M. L. and Pesonen, L. (2003): Magnetic classification of stony meteorites: 1. Ordinary chondrites. Meteoritics & Planetary Science, **38**, 251–268.

Rochette, P., Gattacceca, J., Bonal, L., Bourot-Denis, M., Chevrier, V., Clerc, J. P., Consolmagno, G., Folco, L., Gounelle, M., Kohout, T., Pesonen, L., Quirico, E., Sagnotti, L. and Skripnik, A. (2008): Magnetic classification of stony meteorites: 2. Non-ordinary chondrites. Meteoritics & Planetary Science, **43**, 959–980.

Rochette, P., Gattacceca, J., Bourot-Denise, M., Consolmagno, G., Folco, L., Kohout, T., Pesonen, L. and Sagnotti, L. (2009): Magnetic classification of stony meteorites: 3. Achondrites. Meteoritics & Planetary Science, **44**, 405–427.

Tsuchiya, N., Ishikawa, M., Satish-Kumar, M., Kawakami, T., Kojima, H., Kaiden, H., Miura, H., Suganuma, Y., Abe, M., Sasaki, D., Chiba, M., Okada, Y., Hashizume, F., Grantham, G. and Goderis, S. (2012): Report on geological, geomorphological and meteorite fieldwork in the Sør Rondane Mountains, Eastern Dronning Maud Land, 2009-2010 (JARE-51). Nankyoku Shiryô (Antarctic Record), **56**, 295–379 (in Japanese with English abstract).

Yanai, K. (1993): The Asuka-87 and Asuka-88 collections of Antarctic meteorites: Preliminary examination with brief descriptions of some typical and unique-unusual specimens. Proceedings of the NIPR Symposium on Antarctic Meteorites, **6**, 148–170.

Yanai, K., Shiraishi, K. and Kojima, H. (1994): The Asuka-90 meteorites collection from Antarctica: Searching, initial processing and preliminary identification. Proceedings of the NIPR Symposium on Antarctic Meteorites, **7**, 1–8.

Yesiltas, M., Zolensky, M. and Glotch, T. D. (2019): The first Turkish Antarctic meteorite search expedition. 82nd Annual Meeting of the Meteoritical Society. Sapporo, 2019-07-07/12. LPI Contrib. No. 2157, abstract 6161.

Yoshida, M., Ando, H., Omoto, K., Naruse, R. and Ageta, Y. (1971): Discovery of meteorites near Yamato Mountains, East Antarctica. Nankyoku Shiryô (Antarctic Record), **39**, 62–65.

Zekollari, H., Goderis, S., Debaille, V., van Ginneken, M., Gattacceca, J., ASTER Team, Jull, A. J. T., Lenaerts, J., Yamaguchi, A., Huybrechts, Ph. and Claeys, Ph. (2019): Unravelling the high altitude Nansen blue ice field meteorite trap (East Antarctica) and implications for regional palaeo-conditions. Geochimica et Cosmochimica Acta, **248**, 289–310.

Figure captions

Figure 1. Maps outlining the route of this expedition, from Cape Town to the Nansen Ice Field. (a) Cape Town to Novolazarevskaya Air Base (Novo). The area within the rectangle is enlarged in (b). (b) Novo to Princess Elisabeth Antarctica station (PEA). The area within the rectangle is enlarged in (c). (c) Sør Rondane Mountains and the Nansen Ice Field. "BC" denotes the basecamp setup in the Nansen Ice Field. The area within the rectangle is enlarged in (d). (d) Divisions of the Nansen Ice Field. Red and blue dots denote locations of meteorites and blue ice samples collected during the expedition, respectively. Solid grey lines highlight routes and tracks.

Figure 2. Base camp on the Nansen Ice Field at S72.79754°, E24.86002°.

Figure 3. Daily weather data for the Nansen Ice Field. Gray vertical lines indicate days with poor weather that prevented any field work. Morning data (red triangles) represent average of the two measurements that were done before and after breakfast.

Figure 4. Expedition members and their positions in the V-formation during the meteorite search. Initials inside the circles denote the names of the members (see Table 2), while circles denote the positions of the snowmobiles. One field guide (MP) was always at the apex of the V-formation during the search, while the other field guide (ML) was positioned at one end of the V-shape.

Figure 5. (a) Picture of the team members leaving the base camp in a line with an appropriate distance from each other until the search location is reached, (b) the V-formation of the team while searching for meteorites in the field, (c) the field name of the meteorite was written on blue ice and its picture was taken with a scale for documentation, (d) the meteorite was placed in a zip lock bag (Section 6.3) without touching the sample, and the meteorite field name was written on the bag. C-type meteorites were placed in Teflon bags instead of regular polyethylene zip lock bags.

Figure 6. Mass distribution (left) and log χ values (right) of the recovered meteorites in this expedition.

Figure 7. (a) The recovered meteorites were placed in a cool box at the end of each field day. (b) All samples were appropriately packed and secured in the cool box before shipment to Cape Town.



Figure 1. S. Goderis et al.

38 Figure 2. S. Goderis et al.



48 Figure 3. S. Goderis et al.



49

50

51 Figure 4. S. Goderis et al.



Figure 5. S. Goderis et al.



Figure 6. S. Goderis et al.



60 Figure 7. S. Goderis et al.

61

Table 1. Previous and current meteorite search expeditions in the vicinity of the Sør Rondane Mountains and repositories.

Expedition*	Search Location	Field Season	Number and Weight of Recovered Meteorites	Current Repository[+]	Literature
JARE-27	Mt. Balchen	1986-1987	3 - 2.2 kg	NIPR	Nishio *et al.* (1987)
JARE-29	Mt. Balchen & Nansen	1987-1989	1897 - 375 kg	NIPR	Naraoka *et al.* (1990)
JARE-31	Mt. Balchen	1990-1991	48 - 10.6 kg	NIPR	Yanai *et al.* (1994)
JARE-51/BELARE-SAMBA	Mt. Balchen	2009-2010	635 - 13 kg	NIPR	Kaiden *et al.* (2010)
BELARE-SAMBA	Nansen A	2010-2011	218 - 6 kg	NIPR & RBINS	Goderis *et al.* (2011)
JARE-54/BELARE-SAMBA	Nansen B	2012-2013	424 - 70 kg	NIPR & RBINS	Imae *et al.* (2015)
TAE-III	Nansen A	2018-2019	3 - 64.4 kg	Kirk U	Yesiltas *et al.* (2019)
BELARE-BELAM/TAE-IV	Nansen C	2019-2020	66 - 8.3 kg	NIPR & RBINS	

*JARE, Japanese Antarctic Research Expedition; BELARE, Belgian Antarctic Expedition; SAMBA, Search for Antarctic Meteorites, Belgian Activities; TAE, Turkish Antarctic Expedition; BELAM, Belgian Antarctic Meteorites.

[+]NIPR, National Institute of Polar Research; RBINS, Royal Belgian Institute of Natural Sciences; KirkU, Kirklareli Univeristy.

Table 2. BELAM 2019-2020 joint meteorite search expedition team members, their affiliation and previous Antarctic experience.

Team member	Nationality	Affiliation	Antarctic Experience	Responsibility
Steven Goderis (SG)	Belgium	Vrije Universiteit Brussel	Total of 5 expeditions	Team leader, planning, ice sampling, packaging.
Hamed Pourkhorsandi (HP)	Iran	Université Libre de Bruxelles	1st expedition	Samples, magnetic susceptibility measurements.
Mehmet Yesiltas (MY)	Turkey	Kirklareli University	Total of 2 expeditions	Weather observation, daily logs, packaging.
Naoki Shirai (NS)	Japan	Tokyo Metropolitan University	1st expedition	GPS, transportation.
Manu Poudelet (MP)	France	International Polar Foundation	Total of 2 expeditions	Logistics, communications.
Martin Leitl (ML)	Germany	International Polar Foundation	Total of 2 expeditions	Logistics, medical, mechanical problems.

Table 3. Typical daily schedule at the Base Camp.

Time	Item
7:20	Weather observation (MY)
7:30 - 8:30	Breakfast (all members)
8:30	Weather observation (MY)
8:30 - 9:00	Briefing (all members)
9:00 - 10:00	Preparation for departure (all members)
10:00	Departure for meteorite search (all members)
10:00-17:00	Meteorite search (all members)
17:00 - 19:30	Daily tasks (all members)
19:30 - 20:30	Dinner (all members)
20:30	Weather observation (MY)
20:30 - 21:00	Briefing (all members)
21:00 -	Personal tasks and sleep (all members)

Table 4. Daily log of the 2019-2020 meteorite search expedition.

Date (2020)	Location	Recovered meteorites	Distance driven by snowmobile (km)	Comments
12 Jan	Cape Town			Meeting of expedition members
13 Jan	Cape Town			Departure, D10 flight
13 Jan	Novo Airbase			Arrival, Novo Airbase
13 Jan	Novo Airbase			Departure, Novo Airbase
13 Jan	PEA station			Arrival, PEA station
14 Jan	PEA station			Preparation at PEA station
15 Jan	Nansen Ice Field			Arrival, Base Camp (BC)
16 Jan	Nansen Ice Field			Preliminary search
17 Jan	Nansen Ice Field		7.6	Reconnaissance trip
18 Jan	Nansen Ice Field	6	51.4	Field work
19 Jan	Nansen Ice Field	1	26.4	Field work
20 Jan	Nansen Ice Field	2	55.3	Field work
21 Jan	Nansen Ice Field	9	51.2	Field work
22 Jan	Nansen Ice Field	16	37.8	Field work
23 Jan	Nansen Ice Field	12	67.4	Field work
24 Jan	Nansen Ice Field			Poor weather. No field work
25 Jan	Nansen Ice Field			Poor weather. No field work
26 Jan	Nansen Ice Field	2	72.2	
27 Jan	Nansen Ice Field	18	73.0	
28 Jan	Nansen Ice Field			Poor weather. No field work
29 Jan	Nansen Ice Field			Poor weather. No field work
30 Jan	Nansen Ice Field		39.0	Half day field work
31 Jan	Nansen Ice Field			Poor weather. No field work
1 Feb	Nansen Ice Field		56.8	Half day field work
2 Feb	Nansen Ice Field		52.7	Visit to Nansen B
3 Feb	Nansen Ice Field			Poor weather. No field work
4 Feb	Nansen Ice Field			Poor weather. No field work
5 Feb	Nansen Ice Field			Poor weather. No field work
6 Feb	Nansen Ice Field			Departure from base camp
7 Feb	PEA station			Arrival, PEA station
8 Feb	PEA station			Unloading containers
9 Feb	PEA station			Unloading containers
10 Feb	PEA station			Weighing meteorites
11 Feb	PEA station			Packaging meteorites/cargo
12 Feb	PEA station			Departure, PEA station
12 Feb	Novo Airbase			Arrival, Novo Airbase
13 Feb	Cape Town			Arrival, Cape Town, D13 flight
Total		66	301.6	

Table 5. Daily average weather observation log at the Base Camp in Nansen Ice Field.

Day (2020)	Temperature (°C)*	Wind speed (m/s)	Temperature (°C)+	Humidity (%)	Pressure (hPa)
16 Jan	-22.0	9.9	-38.4	95.5	994.2
17 Jan	-22.0	7.8	-37.0	77.5	995.6
18 Jan	-21.5	7.8	-36.2	67.1	1003.2
19 Jan	-21.8	8.7	-35.7	96.2	1003.5
20 Jan	-20.3	10.0	-36.7	97.7	1004.1
21 Jan	-19.3	9.9	-35.5	86.4	1004.8
22 Jan	-21.2	9.6	-37.7	69.7	1003.7
23 Jan	-22.0	8.2	-36.2	92.2	1005.5
24 Jan	-22.5	10.6	-40.8	94.6	1007.3
25 Jan	-22.2	2.9	-40.2	87.5	1005.7
26 Jan	-22.2	7.9	-36.0	95.5	1006.2
27 Jan	-21.9	7.9	-37.3	96.2	1006.2
28 Jan	-24.7	12.1	-43.5	99.9	985.0
29 Jan	-21.0	10.4	-36.3	95.4	994.4
30 Jan	-21.8	10.3	-38.5	96.1	999.6
31 Jan	-24.4	11.0	-42.8	97.6	999.3
1 Feb	-25.6	11.7	-45.4	99.5	998.9
2 Feb	-25.2	11.3	-44.7	98.1	995.3
3 Feb	-21.6	9.9	-37.7	99.4	995.5
4 Feb	-21.8	11.1	-41.3	99.3	994.6
5 Feb	-22.6	11.3	-37.5	99.4	997.7
6 Feb	-23.5	9.1	-36.5	99.3	998.8

*Average absolute temperature. +Average temperature with the windchill effect.