

<u>Abstract</u>	i
<u>Abbreviations</u>	vii
<u>Chapter 1 Introduction</u>	1
<u>1.1 Function of oxygen in living organisms</u>	1
<u>1.2 Adaptation of animal life to variable environmental oxygen concentrations</u>	1
<u>1.3 Concept of hypoxia</u>	3
<u>1.3.1 Environmental hypoxia</u>	4
<u>1.3.2 Functional hypoxia</u>	4
<u>1.4 Systemic response to hypoxia</u>	6
<u>1.4.1 Increasing oxygen supply</u>	6
<u>1.4.2 Decreasing oxygen demand</u>	6
<u>1.5 Cellular response to hypoxia</u>	8
<u>1.5.1 Increasing metabolic enzyme activity in hypoxia</u>	8
<u>1.5.2 Switch from aerobic to anaerobic metabolism</u>	10
<u>1.5.3 Inhibition of ATP-dependent ion pumping by hypoxia</u>	12
<u>1.5.4 Translation in eukaryotic cells</u>	14
<u>1.5.5 Global translation inhibition under cellular hypoxic condition</u>	17
<u>1.5.6 Transcriptional regulation of gene expression by hypoxia inducible factors (HIF)</u>	24
<u>1.5.7 HIF regulation</u>	24
<u>1.5.8 Families of HIF factors</u>	26
<u>1.5.9 The evolution of HIFs and HIF components</u>	28
<u>1.5.10 Conservation of HIFs and HIF components</u>	28
<u>1.5.11 HIF therapeutics</u>	30
<u>1.5.12 HIF-independent transcriptional regulation in hypoxia</u>	32
<u>1.6 Mechanisms of translation under hypoxia</u>	36
<u>1.6.1 Function of uORF, 5'-TOP and IRES and in specific 5'UTR mediated translation under hypoxia</u>	36
<u>1.6.2 Influence of 3'UTR on translation</u>	38
<u>1.6.3 Role of 3'UTR binding proteins</u>	38
<u>1.6.4 Function of 3'UTR localized sequences in specific mRNA translation under hypoxia</u>	39
<u>1.7 Hypoxia in invertebrates</u>	41
<u>1.7.1 Variable environmental oxygen concentrations in invertebrates</u>	41
<u>1.7.2 Oxygen delivery system in invertebrates</u>	41
<u>1.7.3 HIFs in invertebrates</u>	43
<u>1.7.4 <i>Drosophila melanogaster</i> as a model to study translational control in hypoxia</u>	44
<u>1.7.5 Life cycle of <i>Drosophila melanogaster</i></u>	46
<u>Chapter 2 Objectives</u>	48
<u>2.1 Characterize general features of transcripts induced by hypoxia</u>	48
<u>2.2 Investigate the molecular mechanisms of translational control under hypoxia in <i>Drosophila melanogaster</i></u>	48
<u>2.3 Identification of initiation factors required for translation in hypoxia</u>	49
<u>2.4 Identification of RNA-binding proteins involved in hypoxia translation</u>	49
<u>Chapter 3 Results</u>	51

<u>3.1 Translation of hypoxia-induced mRNAs in <i>Drosophila</i></u>	51
<u>3.1.1 Identification of upregulated mRNAs under hypoxia in S2 cells</u>	53
<u>3.1.2 Translation of mRNAs candidates under hypoxia</u>	55
<u>3.1.3 Candidates genes are translated in a 3'UTR dependent manner</u>	57
<u>3.1.4 “CA-rich” motif is essential for 3'UTR mediated translation in hypoxia</u>	57
<u>3.1.5 “CA-rich” motifs in 3'UTRs of candidate genes are sufficient to induce polysomes formation on a reporter mRNA</u>	59
<u>3.1.6 Adenosines in “CA-rich” motifs play a key role in the expression of reporter gene under hypoxia</u>	59
<u>3.1.7 3'UTR CA-rich motifs are enriched in subsets of hypoxia induced mRNA</u>	61
<u>3.2 4EHP drives the translation under hypoxia</u>	64
<u>3.2.1 <i>Drosophila</i> initiation factor</u>	66
<u>3.2.2 Expression of eIF4E family genes in S2 cells under hypoxia</u>	66
<u>3.2.3 4EHP is necessary for the production of LDH under hypoxia</u>	68
<u>3.2.4 Reduced accumulation of candidate mRNAs in polysomes of 4ehp KO cells in hypoxia</u>	70
<u>3.2.5 4EHP is required for the efficient 3'UTR-mediated translation of mRNA under hypoxia</u>	70
<u>3.2.6 4EHP protein is recruited to polysome fractions under hypoxia</u>	75
<u>3.2.7 4EHP represses global translation under hypoxia in S2 cells</u>	75
<u>3.2.8 4EHP is required for <i>Drosophila melanogaster</i> development in a hypoxic environment</u>	78
<u>3.3 Investigating the potential binding of <i>Drosophila</i> LARK to the 3'UTR of Ldh mRNA under hypoxia</u>	80
<u>3.3.1 Long-term inactivation of Lark is lethal to S2 cells</u>	80
<u>3.3.2 Inactivation of LARK reduces LDH synthesis in hypoxic <i>Drosophila</i> S2 cells</u>	82
<u>3.3.3 Localization of LARK and 4EHP in 293T cells</u>	82
<u>3.3.4 Proteins LARK and 4EHP are partially recruited to polysomal fractions in <i>Drosophila</i> S2 cells</u>	84
<u>3.3.5 Investigating the potential interaction of LARK and 4EHP</u>	84
<u>3.3.6 Identification of factors binding to Ldh and Sirup 3'UTR under hypoxia</u>	86
<u>3.3.7 Investigating the possible binding of LARK to the Ldh mRNA 3'UTR</u>	86
<u>Chapter 4 Conclusions, Discussion and Perspectives</u>	89
<u>4.1 Conclusions</u>	89
<u>4.2 Discussion and Perspectives</u>	89
<u>Chapter 5. Methods and Materials</u>	101
<u>5.1 Reagents and instruments</u>	101
<u>5.2 Cells culture and treatment</u>	102
<u>5.3 Fly strains and maintenance</u>	102
<u>5.4 Construction of stable cell lines</u>	105
<u>5.5 Generation of 4EHP antibody</u>	105
<u>5.6 Sucrose density gradient polysome analysis</u>	106
<u>5.7 RNA and protein extraction</u>	106
<u>5.8 Northern blot and RT-qPCR</u>	107
<u>5.9 RNA interference</u>	107
<u>5.10 Western blot</u>	108

<u>5.11 Dual luciferase assay</u>	109
<u>5.12 Inhibition of protein synthesis (Puromycin labelling)</u>	109
<u>5.13 Co-immunoprecipitation</u>	111
<u>5.14 EMSA</u>	111
<u>5.14 Surveyor nuclease assay</u>	112
<u>5.16 Bioinformatic analysis</u>	112
<u>5.17 Secondary structure of Ldh 3'UTR</u>	113
<u>5.18 Immunofluorescence</u>	113
<u>5.19 Establishment of 4ehp rescue stable cell line</u>	114
<u>5.20 Statistics and graphs</u>	114
<u>Funding and support</u>	114
<u>Appendix</u>	115
<u>References</u>	122