Kcnk3 dysfunction exaggerates the development of pulmonary hypertension induced by left ventricular pressure overload

Mélanie Lambert ^{1,2,3†}, Pedro Mendes-Ferreira ^{4,5†}, Maria-Rosa Ghigna ^{1,2,3},
Hélène LeRibeux^{1,2,3}, Rui Adão⁴, Angèle Boet^{1,2,3}, Véronique Capuano^{1,2,3},
Catherine Rucker-Martin^{1,2,3}, Carmen Brás-Silva⁴, Rozenn Quarck ^{5,6},
Valérie Domergue⁷, Jean-Luc Vachiéry⁸, Marc Humbert ^{1,2,3}, Frédéric Perros ^{1,2,3},
David Montani ^{1,2,3}, and Fabrice Antigny ^{1,2,3*}

¹Univ. Paris-Sud, Faculté de Médecine, Université Paris-Saclay, Le Kremlin Bicêtre, France; ²Assistance Publique Hôpitaux de Paris, Service de Pneumologie, Centre de Référence de l'Hypertension Pulmonaire, Hôpital Bicêtre, Le Kremlin Bicêtre, France; ³Inserm UMR_S 999, Hôpital Marie Lannelongue, 133, Avenue de la Résistance, F-92350 Le Plessis Robinson, France; ⁴Cardiovascular R&D Center, Faculty of Medicine of the University of Porto, Porto, Portugal; ⁵Laboratory of Respiratory Diseases & Thoracic Surgery (BREATHE), Department of Chronic Diseases & Metabolism (CHROMETA), KU Leuven—University of Leuven, Leuven, Belgium; ⁶Clinical Department of Respiratory Diseases, University Hospitals of Leuven, Leuven, Belgium; ⁷Animal Facility, Institut Paris Saclay d'Innovation Thérapeutique (UMS IPSIT), Université Paris-Saclay, Châtenay-Malabry, France; and ⁸Department of Cardiology, Cliniques Universitaires de Bruxelles—Hôpital Erasme, Brussels, Belgium

Received 17 June 2020; editorial decision 4 January 2021; accepted 9 January 2021; online publish-ahead-of-print 23 January 2021

Aims

Pulmonary hypertension (PH) is a common complication of left heart disease (LHD, Group 2 PH) leading to right ventricular (RV) failure and death. Several loss-of-function (LOF) mutations in *KCNK3* were identified in pulmonary arterial hypertension (PAH, Group 1 PH). Additionally, we found that *KCNK3* dysfunction is a hallmark of PAH at pulmonary vascular and RV levels. However, the role of *KCNK3* in the pathobiology of PH due to LHD is unknown.

Methods and results

We evaluated the role of *KCNK3* on PH induced by ascending aortic constriction (AAC), in WT and *Kcnk3*-LOF-mutated rats, by echocardiography, RV catheterization, histology analyses, and molecular biology experiments. We found that *Kcnk3*-LOF-mutation had no consequence on the development of left ventricular (LV) compensated concentric hypertrophy in AAC, while left atrial emptying fraction was impaired in AAC-*Kcnk3*-mutated rats. AAC-animals (WT and *Kcnk3*-mutated rats) developed PH secondary to AAC and *Kcnk3*-mutated rats developed more severe PH than WT. AAC-*Kcnk3*-mutated rats developed RV and LV fibrosis in association with an increase of *Col1a1* mRNA in right ventricle and left ventricle. AAC-*Kcnk3*-mutated rats developed severe pulmonary vascular (pulmonary artery as well as pulmonary veins) remodelling with intense peri-vascular and peri-bronchial inflammation, perivascular oedema, alveolar wall thickening, and exaggerated lung vascular cell proliferation compared to AAC-WT-rats. Finally, in lung, right ventricle, left ventricle, and left atrium of AAC-*Kcnk3*-mutated rats, we found a strong increased expression of *Il-6* and *periostin* expression and a reduction of lung *Ctnnd1* mRNA (coding for p120 catenin), contributing to the exaggerated pulmonary and heart remodelling and pulmonary vascular oedema in AAC-*Kcnk3*-mutated rats.

Conclusions

Our results indicate that *Kcnk3*-LOF is a key event in the pathobiology of PH due to AAC, suggesting that *Kcnk3* channel dysfunction could play a potential key role in the development of PH due to LHD.

*Corresponding author. Tel: +33 1 40 94 22 99; fax: +33 1 40 94 25 22, E-mail: fabrice.antigny@université-paris-saclay.fr

† The first two authors contributed equally to the work.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2021. For permissions, please email: journals.permissions@oup.com.

Graphical Abstract

Keywords

PH due to left heart diseases • Ascending-aortic constriction • Proliferation • K2P3.1 • Task-1

1. Introduction

Pulmonary hypertension (PH), which constitutes an abnormal increase in blood pressure in the pulmonary circulation, has seen its definition revised during the last PH world symposium, as mean pulmonary arterial pressure (mPAP) >20 mmHg.¹ PH can be divided into five sub-groups, being Group 1, or pulmonary arterial hypertension (PAH). Other forms of PH, such as PH due to left heart disease (PH-LHD), have much higher incidence and prevalence, and although constitutes one of the most common forms of PH, roughly representing 50% of PH cases.² However, molecular mechanisms involved in the development of PH-LHD are poorly documented.

The pathophysiology of PH-LHD is complex and is caused by several mechanisms: (i) an initial passive increase in left ventricular (LV) filling pressures more specifically left atrial (LA) pressure; (ii) pulmonary artery endothelial dysfunction; (iii) pulmonary vascular (artery and vein) remodelling; and (iv) right ventricular (RV) dysfunction and altered RV/pulmonary artery coupling.²

Pulmonary endothelial dysfunction seems to play a central role in PH-LHD in both preclinical models^{3–5} and patients.⁶ Endothelium-derived vasorelaxation is markedly impaired in pulmonary arteries from animals with PH-LHD induced by ischaemic⁷ or pressure overload conditions.³

Similarly, exacerbated pulmonary arterial and venous remodelling is observed in patients with both heart failure (HF) with reduced or preserved (HFpEF) ejection fraction,^{8,9} mostly explained by endothelial dysfunction and an imbalance between vasodilation/vasoconstriction and growth factor signalling.¹⁰ To date, most of the molecular pathways associated with PH-LHD are assumed to be common with PAH namely endothelial dysfunction, mitochondrial dysfunction, exacerbated inflammation.¹¹ In PH-LHD, endothelial dysfunction is at least characterized by an increased endothelin-1 (ET-1) production in the pulmonary vasculature,¹² as well as in ascending aortic constriction (AAC)-induced PH in rat.⁴ ET-1 inhibits KCNK3 function in pulmonary artery smooth muscle cell (PASMC) through Rho kinase-mediated phosphorylation,¹³ linking one of the main modulators of PH-LHD (ET-1) and KCNK3-dysfunction. KCNK3 protein, also known as TASK-1 (Twik-related-acid-sensitive-K⁺ channel) is an outward-K⁺ channel. KCNK3 regulates the resting membrane potential in several cell types, including PASMCs¹⁴ and RV cardiomyocytes.¹⁵ The role of KCNK3 dysfunction in the development of PAH was previously highlighted by the identification of several loss-of-function (LOF) mutations in the KCNK3 gene, in PAH patients.^{16,17} In PH Group 1, 16 different mutations were identified in PAH patients, 93.6% of them were missense mutations, most of them were heterozygous, and only one patient was identified with a

homozygous mutation,¹⁶ and by the fact that *KCNK3* mutation carriers' are younger and have higher mPAP at diagnosis.¹⁴

We have recently developed the first *Kcnk3*-mutated rat line with a *Kcnk3*-LOF-mutation (*Kcnk3*^{Δ94ex1/Δ94ex1}), which develops exaggerated PH when challenged with either monocrotaline or chronic hypoxia.¹⁸ Furthermore, *Kcnk3*-LOF-mutated rats are predisposed to pulmonary artery constriction, and have compromised endothelium-dependent and independent relaxation.¹⁸

Assuming the importance of KCNK3 in the modulation of pulmonary vascular tone, and the fact that endothelial dysfunction in PH-LHD should induce KCNK3 dysfunction, we hypothesize that KCNK3 dysfunction could be a key player in PH-LHD pathobiology and that *Kcnk3*-mutated rats are more susceptible to the development of PH secondary to LHD induced by LV pressure overload (AAC).

2. Methods

Due to space limitation, an expanded methods section is provided in the [Supplementary material online](#).

2.1 Animals and surgical procedures

The animal facility is licensed by the French Ministry of Agriculture (agreement N° C92-019-01). This study was approved by the Committee on the Ethics of Animal Experiments (CEEA26 CAP Sud). The animal experiments were approved by the French Ministry of Higher Education, Research and Innovation (N°7757). The animal experiments were performed conforming to the guidelines from Directive 2010/63/EU on 22 September 2010 of the European Parliament on the protection of animals used for scientific purposes and complied with the French institution's guidelines for animal care and handling.

AAC was induced in rats under anaesthesia and analgesia, by placing a stainless steel hemoclip (0.6 mm internal diameter) on the ascending aorta via thoracic incision. The surgical procedure was carried out on 3-week-old rats (WT/*Kcnk3*^{Δ94ex1/Δ94ex1}, *Kcnk3*^{Δ94ex1/+} rats), under anaesthesia with a ketamine-xylazine mix (respectively, 75–10 mg/kg and 0.3 mL/100 g, i.p.). Buprenorphine chlorhydrate (0.03 mg/kg, 0.2 mL/100g, s.c.) was administered twice daily for 3 days beginning at the end of the surgery. Age-matched Sham-operated animals (WT and *Kcnk3*-mutated) underwent the same procedure without placement of the clip. Echocardiographic measurement and closed-chest RV catheterization were performed in operated (AAC and Sham) animals 7 weeks after surgery. After RV catheterization, under profound anaesthesia (isoflurane 5%), the animals were euthanized by exsanguination (section of the descending abdominal aorta) and tissues harvest were performed in all animals.

2.2 Statistical analyses

All statistical tests were performed using GraphPad Prism software (GraphPad, version 8.0 for Windows). All data were verified for normal distribution using Shapiro–Wilk normality. Positively skewed data was log-transformed for analysis, while for clarity reasons graphical representations are shown in their original scale. All values are reported as mean ± standard error of the mean. For experiments with more than five samples, the difference between groups was assessed by two-way ANOVA analysis completed by Sidak's multiple comparisons test for *post hoc* analyses. For experiments with less than five samples, the difference between two groups was assessed by one-way ANOVA analysis completed by

Dunn's multiple comparisons test for *post hoc* analyses after non-parametric ANOVA (Kruskal–Wallis). Differences were considered statistically significant at *P*-values <0.05.

3. Results

3.1 AAC-induced LV pressure overload leads to concentric hypertrophy

To determine the role of *Kcnk3* channel in the development PH-LHD, we attempted to induce PH secondary to LV pressure overload by performing ascending aorta constriction (AAC), in WT and *Kcnk3*-mutated rats (homozygous and heterozygous *Kcnk3*-mutated rat, *Kcnk3*^{Δ94ex1/Δ94ex1} and *Kcnk3*^{Δ94ex1/+}, respectively) as compared to littermate Sham animals, as previously described.¹⁹ For clarity reasons, we present all results obtained in heterozygous *Kcnk3*-mutated rats in the [Supplementary material online](#).

Three weeks old animals were submitted to AAC, and after 7 weeks, cardiac structure and function was evaluated by echocardiography. LV cavity sizes remained within the normal range, resulting in unaltered LV ejection fraction (LVEF) in our different genotypes (Figure 1A and B). AAC resulted in significant LV wall hypertrophy [as evaluated by LV wall thickness, as relative wall thickness] was also increased (Figure 1C). LV hypertrophy were confirmed at the whole organ and cardiomyocyte level (Figure 1D–F), and revealed a concentric hypertrophy pattern (Figure 1G), in both WT and *Kcnk3*-mutated animals submitted to pressure overload (AAC). Similar results were obtained in heterozygous *Kcnk3*-mutated rats (see [Supplementary material online, Figure S1](#)).

Of note, the aortic mean and peak gradient across the constriction were not different between WT and homozygous or heterozygous *Kcnk3*-mutated animals (see [Supplementary material online, Figure S2](#)). No changes in *Kcnk3* mRNA expression were detected in either ventricles or lung after AAC-surgery (Figure 1H).

3.2 *Kcnk3*-LOF mutation leads to LV diastolic dysfunction

Despite similar LV hypertrophic responses and preserved LVEF, diastolic function was compromised in *Kcnk3*-mutated rats (Figure 2A). Doppler-derived trans-mitral flow showed an increase in peak early diastolic flow velocity (E-wave) in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (Figure 2B). Furthermore, tissue Doppler imaging-derived mitral annular early diastolic peak myocardial velocity (*e'*) was significantly decreased in *Kcnk3*-mutated rats (Sham or AAC conditions), resulting in a quasi-significant increase in the *E/e'* ratio in *Kcnk3*-mutated rats, suggesting increased LV filling pressures. Myocardial relaxation dynamics (IVRT, Figure 2C) were affected by surgery, but was uninfluenced by the genotype. A similar pattern was observed in *Kcnk3*^{Δ94ex1/+} rats, in which *E/e'* was significantly increased, despite an apparently *e'* velocity (see [Supplementary material online, Figure S3A and B](#)). The diastolic alterations present in Sham-*Kcnk3*-mutated rats suggest that *Kcnk3*-LOF mutation compromises myocardial compliance, predisposing *Kcnk3*-mutated rats to diastolic dysfunction.

Underlying LV diastolic dysfunction, LV fibrosis was present in AAC rats, as shown by an increased proportion of interstitial fibrotic deposition (Figure 2D and E), even more pronounced in homozygous (Figure 2E) and heterozygous (see [Supplementary material online, Figure S3C](#)) AAC-*Kcnk3*-mutated rats. This increase was associated with an up-regulation of pro-fibrotic transcripts (*Col3a1* and *Periostin*) in left ventricle

Figure 2 *Kcnk3*-LOF mutation leads to LV diastolic dysfunction. (A) Mitral inflow and mitral annulus velocity analysis, using pulse wave Doppler (PWD) and tissue-Doppler imaging (TDI). Vertical bars, 50 (upper) and 5 (lower panel) cm/s, horizontal bars represent 100 ms. (B) Left panel: E-wave, middle panel: e', right panel: E/e' from Sham-WT ($n = 6-7$), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} ($n = 7-8$), AAC-WT ($n = 7-8$), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats ($n = 8-9$). (C) IVRT from Sham-WT ($n = 5$), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} ($n = 8$), AAC-WT ($n = 8$), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats ($n = 9$). (D) Representative images of interstitial collagen deposition in LV tissue sections stained with sirius red from WT and *Kcnk3*-mutated sham or AAC rats. Scale bar, 100 μ m. (E) Analysis of the percentage of LV interstitial collagen staining in Sham-WT ($n = 3$), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} ($n = 3$), AAC-WT ($n = 4$), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats ($n = 3$). (F) Relative mRNA expression (2^{-ΔΔCt}) of *Myh6*, *Myh7*, *Nppa*, *Nppb*, *Col1a1*, *Col3a1*, *Periostin*, and *IL-6* (heatmap representation of the mean) in LV from Sham-WT ($n = 6$), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} ($n = 5$), AAC-WT ($n = 6$), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats ($n = 6$). Experiments were normalized by 18S mRNA and analysed using Δ Ct values (for a full statistical analysis refer to the [Supplementary material online](#)). Data are represented as scatter dot plots, with mean $\pm$ SEM. Experiments presented in panels A–D and F were analysed using two-way ANOVA followed by Sidak's *post hoc* test, and experiments presented in panel E were analysed using one-way ANOVA followed by Dunn's *post hoc* test after Kruskal–Wallis test. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. Fibrosis was not normally distributed and was log-transformed prior to statistical analysis.

from AAC-*Kcnk3*-mutated rats (Figure 2F, and see [Supplementary material online, Table S1](#)). Also associated with diastolic dysfunction, the left ventricle of AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (re)expressed the pathological hypertrophic markers atrial and brain natriuretic peptides (*Nppa* and *Nppb*), and shifted towards the HF prone myosin heavy chain beta gene (*Myh7*) (Figure 2F, and see [Supplementary material online, Table S1](#)). The pro-inflammatory cytokine, *Il6*, gene expression was up-regulated (six-fold).

3.3 Left atrial dysfunction underlines *Kcnk3* LOF mutation-associated with LV diastolic dysfunction

In parallel to LV stiffness, we observed a significant dilation of the left atrium, associated with compromised LA function evaluated by LA emptying fraction (Figure 3A–C, and see [Supplementary material online, Figure S3D](#) and E), in homozygous and heterozygous *Kcnk3*-mutated rats submitted to AAC. As previously shown in human and rat heart,²⁰ we observed an atrial predominance for *Kcnk3* expression (Figure 3D). This indicates that KCNK3 plays an important role in modulating LA function, and its LOF could affect atrioventricular flow, impaired in Sham-*Kcnk3*-mutated rats (Figure 2B, see [Supplementary material online, Figure S3A](#)). Indeed, altered gene expression, with activation of pro-remodelling signalling in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats, could either trigger or result from LA structural and functional changes. Natriuretic peptides (*Nppa* and *Nppb*) and *Myh7* up-regulation in LA tissue (Figure 3E, and see [Supplementary material online, Table S1](#)), as surrogates of hypertrophic and myocardial stretch signalling, and massive (>100-fold) increase in *Periostin* and *Il6*, markers of fibrosis and inflammation, fits with the enlarged and dysfunctional LA seen in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1}. These results demonstrate that *Kcnk3*-LOF mutation promotes LA remodelling and diastolic dysfunction in LV pressure overload condition, a common consequence of LV dysfunction playing a major role in patients with HFrEF.²¹

3.4 Pulmonary hypertension is exaggerated in *Kcnk3*-mutated rats with AAC

At the 10-week endpoint, animals were submitted to right heart catheterization, using a closed-chest approach to evaluate severity of PH. Compared to Sham groups, AAC resulted in the development of mild PH in AAC-WT rats, which was more severe in heterozygous and homozygous *Kcnk3*-mutated rats (Figure 4A and B, and see [Supplementary material online, Figure S4B](#)). Right ventricular systolic pressure (RVSP) was strongly correlated with LA dilation (LA maximal area), when looking at AAC rats, irrespectively of genotype, suggesting that PH had a passive/post-capillary component (Figure 4C), as a consequence of LA enlargement/dysfunction by *Kcnk3* LOF-mutation. Sham-*Kcnk3*-mutated rats showed no significant increase in RVSP as compared to Sham-WT (Figure 4A and B, and see [Supplementary material online, Figure S4B](#)), contrarily to what we have previously shown in older *Kcnk3*-mutated rats.¹⁸ This is likely due to the low number of sham animals.

As a consequence of the development of PH, AAC-*Kcnk3*-mutated rats developed more RV hypertrophy compared to AAC-WT rats. RV free wall thickness (Figure 4D and E, and see [Supplementary material online, Figure S4C](#)), cardiomyocyte size, and normalized RV weight were increased in AAC-*Kcnk3*-mutated rats (Figure 4D–F). Although the Fulton index decreased in AAC animals (Figure 4F), as a consequence of LV hypertrophy, it was higher in AAC-*Kcnk3*-mutated rats when compared with AAC-WT (Figure 4F and see [Supplementary material online, Figure](#)

[S4C](#)), a sign of RV hypertrophy, fitting with the increased RVSP observed in these group. Despite increased PH severity and RV hypertrophy, no significant changes were observed in RV function, with only a mild non-significant decreased in tricuspid annular plane systolic excursion (TAPSE) (Figure 4D and E), when comparing all experimental groups. Interestingly, TAPSE was decreased in Sham and AAC-*Kcnk3*-mutated animals, when compared directly to Sham-WT rats ($P < 0.05$), suggesting that RV function was already compromised, albeit mildly, in animals with a *Kcnk3*-LOF mutation.

In AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats, the development of PH and RV hypertrophy was paralleled with increased expression of overload/hypertrophic response-related genes (*Nppa*, *Nppb* and a shift in the *Myh7/Myh6* relation, Figure 4G, and see [Supplementary material online, Table S1](#)), as well as overexpression of pro-inflammatory (*Il6*) and pro-fibrotic genes (*Col3a1* and *Periostin*), which resulted in increased interstitial fibrosis deposition in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (Figure 4D and E) compared to the AAC-WT rats. Overall, these results indicate that *Kcnk3*-dysfunction facilitates the development of PH-LHD.

Using frozen RV samples, we measured the myofilament properties of isolated skinned RV cardiomyocytes, where we found only a trend towards increased passive tension development in AAC-rats (WT and *Kcnk3*-mutated rats, see [Supplementary material online, Figure S4D](#)), without differences between genotypes. Active tension and calcium sensitivity were not changed (see [Supplementary material online, Figure S4D](#)).

Interestingly, and in agreement with the increase in RVSP, Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} animals showed a significant increase in RV cardiomyocyte size (Figure 4E). Unexpectedly, and without changes in the analysed pro-remodelling transcripts, Sham-*Kcnk3*-mutated rats had a significantly higher percentage of fibrosis deposition in the right ventricle.

No differences were observed in mean carotid arterial pressure, while cardiac output was decreased in AAC-rats, irrespectively of the genotype (see [Supplementary material online, Figure S4A](#) and B), which could be a consequence of LV diastolic dysfunction leading to compromised LV filling.

The development of PH resulted in mild right atrial dilation together with non-significant increase in HF and pro-fibrotic gene expression, in *Kcnk3*-mutated rats (homozygous and heterozygous) (see [Supplementary material online, Figure 5A–E](#)).

3.5 Extensive pulmonary vascular remodelling supports the development of PH in AAC- *Kcnk3*^{Δ94ex1/Δ94ex1} rats

Supporting the development of PH in AAC-rats and in comparison with Sham-group, we observed profound pulmonary structural remodelling in AAC-*Kcnk3*-mutated rats as media hypertrophy/hyperplasia in pulmonary arteries and neomuscularization of the small distal and normally not muscularized pulmonary vessels (Figure 5A and B). Semi-quantitative morphometric analysis of pulmonary vessels remodelling including arterioles and venules indicated a greater proportion of pulmonary vessel remodelling in Sham-*Kcnk3*-mutated rats vs. Sham-WT, and that all pulmonary vessels were remodelled in AAC-*Kcnk3*-mutated rats compared to AAC-WT rats (Figure 5C). The pulmonary vessels wall area was increased in AAC-*Kcnk3*-mutated rats (Figure 5C and see [Supplementary material online, Figure S6A](#)).

We observed the presence of adventitial/perivascular oedema in AAC-induced PH animals (Figure 5D and E, and see [Supplementary material online, Figure S6B](#)), as well as a decrease in *Ctnnd1* expression

Figure 3 Left atrial (LA) dysfunction underlines *Kcnk3*-LOF mutation-associated LV diastolic dysfunction. (A) Using an apical four chamber projection (A4C) we quantified left atrial (LA) dimensions. IAS, interatrial septum; IVS, interventricular septum; LV, left ventricle; RA, right atria; RV, right ventricle. Dashed white boxes represent the zoomed sections in the lower panel, and the blue shaded area delineates the LA. Vertical bars, 0.5 cm, horizontal bars represent 100 ms. (B) Left panel: LA Amax, right panel: LA emptying fraction from Sham-WT ($n = 7$), Sham- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ ($n = 7-8$), AAC-WT ($n = 8$), and AAC- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ rats ($n = 9$), and (C) LA/tibia length from Sham-WT ($n = 5$), Sham- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ ($n = 10$), AAC-WT ($n = 10$), and AAC- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ rats ($n = 15$). (D) Relative mRNA expression ($2^{-\Delta\Delta Ct}$) of *Kcnk3* in LV, RV, LA, and RA from Sham-WT ($n = 6$). Experiments were normalized by 18S mRNA and analysed using ΔCt values. (E) Relative mRNA expression ($2^{-\Delta\Delta Ct}$) of *Myh7*, *Nppa*, *Nppb*, *Periostin*, and *IL6* (heatmap representation of the mean) in LA from Sham-WT ($n = 5$), Sham- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ ($n = 6$), AAC-WT ($n = 5$), and AAC- $Kcnk3^{\Delta94Ex1/\Delta94Ex1}$ rats ($n = 7$). Experiments were normalized by 18S mRNA and analysed using ΔCt values (for a full statistical analysis refer to the [Supplementary material online](#)). Data are represented as scatter dot plots, with mean $\pm$ SEM. Experiments were analysed using two-way ANOVA followed by Sidak's *post hoc* test. * $P < 0.05$ and *** $P < 0.001$. LA Amax, LA/TL, and *Kcnk3* expression were not normally distributed and were log-transformed prior to statistical analysis.

Figure 4 Pulmonary hypertension is exaggerated in *Kcnk3*-mutated rats with AAC. (A) Representation of the record of right ventricular pressure (RVP in mmHg) using right heart catheterization. (B) Quantification of RVSP (in mmHg) (SHAM; WT = 5, $Kcnk3^{\Delta94ex1/\Delta94ex1}$ = 8, AAC; WT = 13, $Kcnk3^{\Delta94ex1/\Delta94ex1}$ = 19 rats). (C) Correlation (Spearman) between RVSP and LAA_{max} in AAC-WT ($n = 3$), AAC- $Kcnk3^{\Delta94ex1/+}$ ($n = 6$), and AAC- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ ($n = 5$) rats. (D) M-mode echocardiographic determination of RV free wall thickness (RVFW) and tricuspid annular plane systolic excursion (TAPSE, upper panels). Representative images of RV cardiomyocytes size (WGA staining) and RV interstitial collagen deposition (Sirius red staining, lower panels). Vertical bars, 0.5 cm (M-Mode) and horizontal bars represent 100 ms, 50 μ m (WGA staining), and 100 μ m (Sirius red). (E) First panel: RVFWT; and second panel: TAPSE from Sham-WT ($n = 7$), Sham- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ ($n = 7-8$), AAC-WT ($n = 8$), and AAC- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ rats ($n = 9$). Third panel: RV cross-section area; and fourth panel: analysis of the percentage of RV interstitial collagen staining from Sham-WT ($n = 5$), Sham- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ ($n = 5$), AAC-WT ($n = 5$), and AAC- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ rats ($n = 5$). (F) RV/tibia length ratio and Fulton index from Sham-WT ($n = 5-8$), Sham- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ ($n = 9-11$), AAC-WT ($n = 9-14$), and AAC- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ rats ($n = 14-17$). (G) Relative mRNA expression ($2^{-\Delta\Delta Ct}$) of *Myh6*, *Myh7*, *Nppa*, *Nppb*, *Col1a1*, *Periostin*, and *Il6* (heatmap representation of the mean) in RV from Sham-WT ($n = 6$), Sham- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ ($n = 5$), AAC-WT ($n = 6$), and AAC- $Kcnk3^{\Delta94ex1/\Delta94ex1}$ rats ($n = 7$). Experiments were normalized by *18S* mRNA and analysed using ΔCt values (for a full statistical analysis refer to the supplemental material). Data are represented as scatter dot plots, with mean $\pm$ SEM. Experiments were analysed using two-way ANOVA followed by Sidak's post hoc test after Kruskal-Wallis test. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. RVSP, RV CSA, RV Fibrosis, RV/TL, and Fulton were not normally distributed and were log-transformed prior to statistical analysis.

Figure 5 Analyses of pulmonary alteration in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats. (A) Representative photography of HES staining of lung from AAC-*Kcnk3*-mutated rats. Upper panel: Smooth muscle cell hypertrophy/hyperplasia in pulmonary arteries, Lower panel: neomuscularization of the small distal and normally not muscularized pulmonary vessels. Scale bars, 50 (upper) and 15 μm (lower). (B) HES staining on lung slice from Sham and AAC WT or *Kcnk3*^{Δ94ex1/Δ94ex1} rats. Scale bar, 100 μm. (C) Left panel: quantification of vessel wall area (μm²). Right panel: analysis of the vascular remodelling score in Sham and AAC WT or *Kcnk3*^{Δ94ex1/Δ94ex1} rats. Sham-WT (n = 4), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} (n = 4), AAC-WT (n = 8), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (n = 8). (D) Illustrative of lung perivascular oedema in AAC-*Kcnk3*-mutated rats. Scale bar, 50 μm. (E) Left panel, percentage of vessels with large adventitia (which could be measured). Right panel, quantification of adventitia surface area (μm²) in Sham-WT (n = 4), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} (n = 4), AAC-WT (n = 8), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (n = 9). (F) Relative mRNA expression (2^{-ΔΔCt}) of *Cttnnd1* (encoding p120Catenin) in lung from Sham-WT (n = 6), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} (n = 5), AAC-WT (n = 6), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (n = 5). Experiments were normalized by 18S mRNA and analysed using ΔCt values. (G) Left panel, illustrative images of severe lung injury with prominent enlargement of alveolar septa and inflammatory exsudates observed in AAC-*Kcnk3*-mutated rats. Scale bar, 500 and 100 μm. Right panel, analysis of the parenchymal alteration score in Sham-WT (n = 4), Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} (n = 4), AAC-WT (n = 4), and AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (n = 4). Data are represented as scatter dot plots, with mean ± SEM. Experiments presented in panels A–B and D–F were analysed using two-way ANOVA followed by Sidak's *post hoc* test, and experiments presented in panels C and G were analysed using one-way ANOVA followed by Dunn's *post hoc* test after Kruskal–Wallis test. **P* < 0.05, ***P* < 0.01. Vascular wall area, oedema, and adventitial area were not normally distributed and were log-transformed prior to statistical analysis.

(Figure 5F), which codes for p120-catenin, an important regulator of endothelial permeability. Despite that, adventitial area was significantly higher in *Kcnk3*^{Δ94ex1/Δ94ex1} rats (Figure 5E, and see [Supplementary material online, Figure S6B](#)).

Interestingly, both adventitial area was increased and *Ctnnd1* expression was decreased in Sham-*Kcnk3*-mutated rats, suggesting that the *Kcnk3*-LOF compromises endothelial integrity, leading to perivascular leakage, and ultimately supporting increased pulmonary pressures,²² and sensitizing the pulmonary circulation to an exaggerated response to passive pressure increases.

Not only the vasculature, but also the lung parenchyma was altered in AAC-induced PH, and aggravated with *Kcnk3*-LOF. In most of AAC-*Kcnk3*-mutated rats we observed severe lung injury with prominent enlargement of alveolar septa (Figure 5G), and inflammatory exudates (Figure 6D). Due to the lack of validated markers capable of differentiating arteries and veins, the assessment of vein remodelling in lung vascular diseases is an ambitious task, requiring accurate identification of venous vessels. In rodents, a reliable histologic item to identify pulmonary veins is based on the presence of a cardiomyocyte coat around pulmonary veins. Unfortunately, we did not find enough cardiomyocytes coat pulmonary veins to quantify their remodelling. However, when we succeeded to identify some cardiomyocytes-coated pulmonary veins in AAC-rats and we observed a severe interstitial oedema in the adventitial layer (see [Supplementary material online, Figure S7A](#)). In AAC-rats, we also observed that pulmonary veins wall thickness was increased with severe adventitial oedema and lymphangiectasis (see [Supplementary material online, Figure S7B](#)), indicating that large pulmonary veins were remodelled in AAC-rats.

Similarly to *Kcnk3*^{Δ94ex1/Δ94ex1}, we also found a significant increase in pulmonary vessel wall thickening, adventitia area in *Kcnk3*^{Δ94ex1/+} (see [Supplementary material online, Figure S6A and B](#)) and the presence of foci alveolar wall thickening in AAC-*Kcnk3*^{Δ94ex1/+} rats (data not shown), with significant parenchymal remodelling (see [Supplementary material online, Figure S6C](#)).

3.6 Characterization of pulmonary remodelling

To evaluate the consequence of *Kcnk3*-LOF in AAC context on *in situ* lung cell proliferation, we visualized cells undergoing DNA replication by imaging the incorporation of 5-ethynyl-2'-deoxyuridine (EdU; white or pink nucleus in Figure 6). We observed very few EdU-positive cells in the lung parenchyma of Sham-WT or Sham-*Kcnk3*^{Δ94ex1/Δ94ex1} rats (data not shown). Underlying the pulmonary structural changes, we found an increase in lung proliferative cells (Figure 6A–C), including PSMCs, as the amount of α -smooth-muscle actin and EdU positive cells was augmented in AAC-*Kcnk3*-mutated rats (*P*-value = 0.056) (Figure 6C). A larger accumulation of proliferating CD34⁺ cells was observed in lung from AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats compared with AAC-WT rats (see [Supplementary material online, Figure S7C](#)).

The analysis of lung histology (HES) revealed that moderate alveolar septa enlargement was mostly characterized by the affluence of inflammatory cells (mostly mononuclear: monocytes, lymphocytes). In more severe cases, we observed that the alveolar septa enlargement was hypercellular, with many inflammatory cells (monocytes, lymphocytes), macrophages, and interstitial oedema. Endo-alveolar macrophages were also visible (Figure 6D). Peri-bronchial inflammation was obvious in AAC-*Kcnk3*-mutated rats (Figure 6E), whose histological score was aggravated compared with WT animals (Figure 6F, and see [Supplementary material](#)

[online, Figure S6E](#)). Elevated levels of IL-6 are described to hyper-activate STAT3 signalling mediating the cell proliferation and cell survival.²³ In association with elevated lung IL-6 mRNA and pulmonary cell proliferation in AAC-*Kcnk3*^{Δ94ex1/Δ94ex1} rats, we found an increase of STAT3 phosphorylation (Figure 6G and H).

4. Discussion

Our findings, summarized in Figure 7, suggest that KCNK3 dysfunction plays a role in the development PH-LHD. In the context of LV pressure overload, we propose that *Kcnk3*-LOF mutation promotes (i) disruption of pulmonary endothelium integrity which leads to lung perivascular oedema and PA adventitial remodelling, (ii) inflammatory signalling (IL-6) which leads to increase PSMC proliferation promoting pulmonary vascular and parenchymal remodelling. (i) and (ii) contribute to the increase in pulmonary vascular resistances. (iii) Cardiac fibrosis links LV diastolic dysfunction to decreased LA compliance and abnormal pulmonary venous remodelling. All together these events act in favour of the aggravation of PH and the consequent RV hypertrophy and dysfunction in *Kcnk3*-mutated rats. Our results are of major interest since understanding of molecular mechanisms involved in PH-LHD pathobiology are yet to be defined.

KCNK3 is expressed in the rat and human heart,²⁰ with a clear atrial predominance, which is maintained in end-stage HF.²⁴ We have previously shown that *Kcnk3* expression is decreased in the right ventricle of PH animals and PAH patients, as well as in the pressure overloaded right ventricle.¹⁵ To understand the potential role of KCNK3 in LV pressure overload-induced PH, we measured its expression in heart and lungs. *Kcnk3* levels were unaltered by AAC surgery, in both left ventricle and right ventricle, and lungs. However, we observed enhanced expression in both right and left atria, in agreement with previously published literature, confirming that KCNK3 might play an important role in the function of atrial chambers. In human, both KCNK3 up-regulation and KCNK3-LOF mutations are associated with detrimental atrial electrophysiological alterations.²⁵ The important role of KCNK3 in left atrium could explain the LA enlargement in AAC-*Kcnk3*-LOF-mutated rats. Directly or indirectly, LV diastolic dysfunction was associated with LA enlargement and functional compromise in AAC-*Kcnk3*-mutated rats. Inactivation of KCNK3 can lead to atrial enlargement when induced in the experimental context.²⁶ Signs of adverse LA remodelling in HF include increased expression of myosin heavy chain- β (*Myh7*),²⁷ natriuretic peptides,²⁸ and *Periostin*,²⁹ all of which were up-regulated in the left atrium of *Kcnk3*-mutated rats with LV pressure overload, and underlie the molecular mechanisms that resulted in compromised LA structure and function in these animals. Increased pulmonary pressures were highly correlated with LA size, which reveals the link between LA structural remodelling and the severity of PH.

In the current work, a *Kcnk3*-LOF-mutation resulted in LV dysfunction, characterized by a pro-fibrotic remodelling and diastolic impairment, despite a similar degree of pressure-overload (gradient) and LV hypertrophic response. Bertero *et al.*,³⁰ recently demonstrated that KCNK3 knockdown up-regulated the expression of *Col3a1* and *Lox* in human pulmonary fibroblasts, which could explain the increased expression of pro-fibrotic genes and fibrosis in all cardiac chambers. In association with increased RV and LV fibrosis in AAC-*Kcnk3*-mutated rats, we found an increase in *Periostin* expression. *Periostin* is known to modulate the transition of fibroblasts to myofibroblasts, collagen fibrillogenesis, extracellular matrix synthesis, as well as the inflammatory

Figure 6 Characterization of pulmonary remodelling in AAC-rats. (A) Immunofluorescent staining of frozen rat lung sections and confocal imaging with Click-iT 5-ethynyl-2'-deoxyuridine (EdU; white or pink nuclei = EdU-positive nuclei = proliferating cells) in combination with α -smooth-muscle actin (α -SMA; in green) in AAC-WT and AAC-*Kcnk3*-mutated rats. Counterstain was DAPI (blue). EdU positive nuclei are indicated by yellow/white arrows, and vessel lumen is indicated (v). Scale bar, 50 μ m (left panels) and 20 μ m (right panels). (B) Quantification of the percentage of lung proliferating cells (% of EdU-positive nuclei) in Sham-WT ($n = 5$), Sham-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 6$), AAC-WT ($n = 6$), and AAC-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ rats ($n = 6$). (C) Quantification of the proportion of EdU positive + α SMA positive cells in Sham-WT ($n = 3$), Sham-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 3$), AAC-WT ($n = 4$), and AAC-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ rats ($n = 6$). (D) Upper panel, in a group of AAC-*kcnk3*-mutated rats alveolar wall appeared thickened. The alveolar septa enlargement is characterized by the affluence of inflammatory cells (mostly mononucleated), and at some degree to alveolar oedema. Lower panel, in severe cases the alveolar septa enlargement is hypercellular, with many inflammatory cells (red arrow), macrophages (blue arrow) and interstitial oedema (black star). Endoalveolar macrophages are also visible (dark blue arrows). No fibrin deposits nor alveolar necrosis are observed. Scale bars, 100 and 50 (zoom) μ m. (E) Severe peri-bronchial/peri-vascular inflammation, extending in peri-bronchiolar alveoli. Scale bars, 100 and 25 (zoom) μ m. (F) Analysis of the peri-bronchiolar inflammation score in Sham-WT ($n = 4$), Sham-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 4$), AAC-WT ($n = 4$), and AAC-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ rats ($n = 4$). (G) Western blot images of the phosphorylation of STAT3 protein in lung lysates. (H) Quantification of the phosphorylation of STAT3 protein in lung from Sham-WT ($n = 6$), Sham-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 6$), AAC-WT ($n = 6$), and AAC-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 7$) rats (see the complete unedited gel in see [Supplementary material online, Figure S8](#)). (I) Relative mRNA expression ($2^{-\Delta\Delta Ct}$) of *Il6* in lung from Sham-WT ($n = 6$), Sham-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ ($n = 6$), AAC-WT ($n = 4$), and AAC-*Kcnk3* $^{\Delta 94Ex1/\Delta 94Ex1}$ rats ($n = 4$). Experiments were normalized by *18S* mRNA and analysed using $\Delta\Delta Ct$ values. Data are represented as scatter dot plots, with mean $\pm$ SEM. Experiments presented in panels A–B, D–E, G–H, and I were analysed using two-way ANOVA followed by Sidak's *post hoc* test, and experiments presented in panels C and F were analysed using one-way ANOVA followed by Dunn's *post hoc* test after Kruskal–Wallis test, *** $P < 0.001$. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. EdU $^{+}$ and EdU $^{+}/\alpha$ SMA $^{+}$ and Stat3 phosphorylation were not normally distributed, and were log-transformed prior to statistical analysis.

Figure 7 Proposed sequence of events arising from *Kcnk3*-LOF mutation in context of LV pressure overload. In context of LV pressure overload, we propose that *Kcnk3*-LOF mutation promotes (i) the disruption of pulmonary endothelium integrity (with a decreased expression of p120 catenin in lung from Sham-*Kcnk3*-mutated vs Sham-WT rats) which leads to lung perivascular oedema and PA adventitial remodelling, (ii) the inflammatory signalling (Increased expression of IL-6 in lung, LV, RV and LA from AAC-*Kcnk3*-mutated rats) which leads to increase PASM C proliferation and promoting pulmonary vascular remodelling and parenchymal remodelling. (i) and (ii) contribute to the increase of pulmonary vascular resistances, and (iii) the cardiac fibrosis (with an increased expression of Col1A1 and periostin in LV and RV from AAC-*Kcnk3*-mutated rats) link to LV diastolic dysfunction, decreased of LA compliance and abnormal pulmonary venous remodelling. All together these deregulations act in favour of the aggravation of pulmonary hypertension and the consequent RV hypertrophy and dysfunction. All these events are responsible for the development of more severe PH-induced by LV pressure overload in *Kcnk3*-mutated rats.

response.³¹ Moreover, Bruns *et al.*,³² recently demonstrated that Periostin-secreted from cardiac fibroblast drives cardiomyocyte dedifferentiation, contributing to RV dysfunction in experimental PH. In our study increased Periostin in the four cardiac chambers could partly explain exaggerated peri-vascular inflammation and cardiac fibrosis.

By contrast to our findings, a recent work using global *kcnk3*-deficient mice, submitted to transverse aortic constriction (TAC), develop reduced cardiac hypertrophy and preserved cardiac function compared to WT-TAC.³³ These opposite effects could be explained either by species/strain-specific differences, as in Wistar rats *Kcnk3* expression is

two-fold higher in the right ventricle as compared to left ventricle,¹⁵ while in the current manuscript, using Sprague-Dawley rats, we found no differences between the two ventricles. Moreover, in mice pulmonary vascular KCNK3 is not functional in PSMCs (Manoury *et al.*, 2011),¹⁸ while KCNK3 plays a key role in the homeostasis of the pulmonary vasculature. Recently, detailed species-related differences in the response to pressure overload³⁴ have been described, fitting with the compromised ejection fraction in WT mice,³³ opposed to our WT rats.

Either as a consequence of increased pulmonary resistances, direct RV effects of *Kcnk3* decreased expression,¹⁵ or *Kcnk3*-LOF-mutation in

AAC animals, worsened RV hypertrophy and function in response to LV pressure overload, is an important observation, since a worse RV function predicts a deteriorated functional capacity in HF patients,³⁵ and affects treatment response in patients with HFpEF,³⁶ making its evaluation in PH-LHD³⁷ a requirement. RV cardiomyocyte hypertrophy and fibrosis in Sham-*Kcnk3*-mutated rats, suggests that the mutation itself promotes maladaptive remodelling, potentially having a direct consequence on RV adaptation in the context of LV pressure overload. The moderate deterioration RV function in our model could be explained by the improvement in RV perfusion secondary to AAC, recently shown to increase right coronary artery perfusion, leading to attenuation of RV functional and structural decline after MCT-exposure.³⁸

Generalized remodelling, involving most lung structures are associated with retrograde pressure build up-induced capillary injury.¹¹ Pulmonary arterial (pre-capillary component) and venous (post-capillary component) remodelling has been recently characterized in detail in patients with HF, irrespectively of EF, and correlated with disease severity.⁸ Accordingly, AAC-*Kcnk3*-mutated animals, showed significant alterations in both pulmonary arteries and veins, with both lumen-compromising medial hypertrophy and neomuscularization of pulmonary vessels.

Adventitial restructuring, characterized by perivascular cuffing (fluid accumulation in the interstitial surrounding the vessels), which was previously described as a consequence of elevated pulmonary venous pressure,²² was present in both AAC-groups (similar percentage of vessels with adventitial oedema), but amplified in homozygous *Kcnk3*-mutated rats (larger adventitial area), which could be explain by decreased *Ctnd1* expression. *Kcnk3*-LOF-mutation was associated with lower *Ctnd1* (p120-catenin) expression, a gatekeeper of endothelial barrier function and integrity,³⁹ even without LV pressure overload, predisposing these animals to endothelial injury, and asserting a causative role. We have previously shown that *Kcnk3*-mutated rats undergo endothelial-to-mesenchymal transition, with loss of CD31 expression, further contributing to endothelial barrier impairment. Unfortunately, we did not measure lung weight as a measure of congestion.

In parallel, extensive parenchymal remodelling, characterized by alveolar septal thickening was patent in overloaded *Kcnk3*-mutated lungs. Similar alterations contribute to limited respiratory efficiency and effort capacity in HF patients with PH.³⁵ As indicated by our present results, alveolar septal enlargement observed in AAC-*Kcnk3*-mutated rats is at least the consequence of the increased CD34⁺ cells proliferation. CD34 is a marker of endothelial cells which is also used as a marker of haematopoietic stem cells, haematopoietic progenitor cells as well other non-haematopoietic cell types, including vascular endothelial progenitors 1 and embryonic fibroblasts.⁴⁰ Since Dierick et al.,⁴¹ found that a population of pulmonary progenitor cells PW1⁺/CD34⁺ are recruited in mice and rat experimental PH models, we could hypothesized that alveolar septal enlargement could also due to a larger accumulation of PW1⁺/CD34⁺ cells.

In vivo, KCNK3 inhibition induced proliferation of CD4⁺ and CD8⁺ lymphocytes,⁴² suggesting an activation of inflammatory processes. Moreover, *kcnk3*-deficient mice and fibroblast-derived induced pluripotent stem cells from patients with a *Kcnk3*-LOF mutation, showed an increased sensitivity to inflammatory stimulus,⁴³ with increased inflammatory mediators expression and an exaggerated infiltration of inflammatory cells in the lung. In harmony, increased IL6/STAT3 activation, which was recently demonstrated to play an important role in experimental and human PH-LHD,⁴⁴ was evident in AAC-*Kcnk3*-mutated rats, leading to extensive peri-bronchial inflammatory infiltrates, and overall

increased pulmonary cellular proliferation, including PASMC, sustaining pulmonary vascular wall remodelling.

In AAC-*Kcnk3*-mutated rats, the inadequate pulmonary vascular remodelling and PH could shed some light on the development of unexplained disproportionate PH in LHD.⁴⁵ In some cases of PH-LHD, severe pulmonary vascular alterations persisted despite resolution of the underlying LV disease.¹¹ Our observation that *Kcnk3*-LOF exaggerates pulmonary vascular remodelling in the context of LV pressure overload, suggests that KCNK3-dysfunction might be associated with a lack of reverse remodelling, and therefore non-response to treatment. Recently, in AAC-induced LV pressure overload with pulmonary vascular remodelling, surgically removing the aortic constriction, normalized pulmonary vascular structure,⁴⁶ therefore using this approach would dissect if structural changes remain in the presence of a *Kcnk3*-mutation.

Importantly, vascular changes were already present in Sham-*Kcnk3*-mutated rats, which could explain the exaggerated pulmonary remodelling in response to LV pressure overload in a *Kcnk3*-LOF-mutated rats. Underlying remodelling in unstressed conditions describes a pre-capillary component in AAC-*Kcnk3*-mutated animals, which could explain the different response to LV pressure overload in the absence and presence of a *Kcnk3*-LOF mutation, and suggests that *Kcnk3*-dysfunction or reduced expression could not only worsen PH, but also compromise its reversibility/treatment in the context of HF.

Several works have used either TAC or AAC to induce LHD or LV pressure overload and to study pulmonary vascular changes.^{19,44} Although they might not replicate all the phenotypical characteristics of group 2 PH, it is clear that pulmonary vascular changes develop in response to a primary insult to the left ventricle, and have granted experimental evidence for PH-LHD treatment, including sildenafil.³ Inhaled iloprost showed a pulmonary-specific vasodilator effect in rats with AAC-induced PH,⁵ which was confirmed in patients with HFpEF.⁴⁷ In animal model of PH-HFpEF, treprostinil attenuated the development of PH.⁴⁸ Both of these prostanoids stimulate KCNK3 currents in human PASMC,¹³ providing that stimulation of KCNK3 activity might play a beneficial role in PH-LHD.

4.1 Limitations

Other works have shown that PH-LHD development of experimental PH-LHD, with measurement of LV filling pressures in both AAC with¹⁹ or without metabolic syndrome associated LV disease.⁴⁴ Unfortunately we were unable to measure LV pressures, using our right heart catheterization, where LA or pulmonary capillary wedge pressure measurement is not possible.⁴⁹ Despite that, we show a clear concentric LV hypertrophy phenotype, characteristic of LV pressure overload in the rat.³⁴

Our echocardiographic analysis lacks *E/A* wave analysis. Due to the elevated heart rate of small animals, and equipment capabilities, *E* and *A* wave separation was not observed in all animals. Indeed, we were only able to measure the *A* wave in around 60% of animals, and without differences between groups, due to the small number of measurements (data not shown). Despite that, in the animals where peak separation occurs, *E* > *A*, allowing us to measure *E* peak velocity with a significant degree of confidence in all animals, we showed that homozygous AAC-*Kcnk3*-mutated rats had an increased *E/e'* ratio.

The access to lung from patients with PH-LHD is extremely difficult, since most of these do not undergo to surgical procedures, we are not able to measure KCNK3 function and expression in human PH-LHD patients. Furthermore, the genetic variations of PH-LHD patients have been barely studied,⁵⁰ consequently no information is available regarding the presence of mutation in KCNK3.

We acknowledge that our model does not represent a pure phenotype of passive PH-LHD. Despite that, we undoubtedly demonstrate that the presence of a *Kcnk3*-LOF-mutation aggravates pulmonary vascular remodelling and PH, generating new hypothesis for the understanding of the pathophysiology.

5. Conclusion

PH secondary to LHD is a common disease, but without specific therapies and a poorly understood pathophysiology. With this work, we established KCNK3 as an important mediator of PH in the context of LV pressure overload. Using *Kcnk3*-mutated rats and a model of AAC-induced LV concentric hypertrophy, we revealed a compromised left heart compliance, and disproportionate pulmonary remodelling, associated with pro-inflammatory and proliferative processes, ultimately leading to exaggerated PH and RV dysfunction. Either by sensitizing the pulmonary circulation directly, or by worsening LV/LA function, we established the importance of KCNK3 in this pathology, and have opened the way for the development of new therapeutic options for this condition.

Supplementary material

Supplementary material is available at *Cardiovascular Research* online.

Authors' contributions

M.L., P.M.-F., M.-R.G., and F.A. participated in the research design. M.L., P.M.-F., M.-R.G., H.L.R., R.A., A.B., V.C., C.R.-M., V.D., and F.A. conducted the experiments and performed the data analysis. All authors draft the manuscript for important intellectual content the manuscript. All authors reviewed and revised the final version and approved manuscript submission.

Acknowledgements

This work benefited from the facilities and expertise of TEFOR—Investissement d'avenir—ANR-II-INSBS-0014. We thank Dr Ignacio Anegón, Laurent Tesson, and Séverine Ménoret from Transgenic Rats and Immunophenomics Core Facility (TRIP), platform TRIP—(Immunology—Nantes) for generating *Kcnk3*-mutant rats. The authors thank Dr J. Sabourin from INSERM UMR S1180, Faculté de Pharmacie, Université Paris-Saclay for critical reading of the manuscript. We wish to thank the staff at the ANIMEX platform for caring for the rat lines.

Conflict of interest

M.H. has relationships with drug companies, including Actelion, Bayer, GSK, Novartis, and Pfizer. In addition to being investigators in trials involving these companies, other relationships include consultancy services and memberships to scientific advisory boards. The other authors have no conflicts of interest.

Funding

This study was supported by grants from the French National Institute for Health and Medical Research (INSERM), the Université Paris-Saclay, the Marie Lannelongue Hospital, the French National Agency for Research (ANR) [grant no. ANR-18-CE14-0023 (KAPAH)]. F.P. received funding from

Fondation maladies rares in the frame of the 'Small animal models and rare diseases' programme to generate the *Kcnk3*-mutated rats. M.L. is supported by Therapeutic Innovation Doctoral School (ED569). P.M.-F., R.A., and C.B.-S. were funded by Portuguese Foundation for Science and Technology through Grant UID/IC/00051/2013 (COMPETE_2020, POCI) and projects IMPAcT (Ref. PTDC/MED-FSL/31719/2017; POCI-01-0145-FEDER-031719) and NETDIAMOND (Ref. POCI-01-0145-FEDER-016385).

Data availability

The authors declare that all supporting data are available within the article and its Online Data Supplement.

References

- Simonneau G, Montani D, Celermajer DS, Denton CP, Gatzoulis MA, Krowka M, Williams PG, Souza R. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* 2019;**53**:1801913.
- Vachiéry J-L, Tedford RJ, Rosenkranz S, Palazzini M, Lang I, Guazzi M, Coghan G, Chazova I, De Marco T. Pulmonary hypertension due to left heart disease. *Eur Respir J* 2019;**53**:1801897.
- Yin J, Kukucka M, Hoffmann J, Sterner-Kock A, Burhenne J, Haefeli WE, Kuppe H, Kuebler WM. Sildenafil preserves lung endothelial function and prevents pulmonary vascular remodeling in a rat model of diastolic heart failure. *Circ Heart Fail* 2011;**4**:198–206.
- Zhuang R, Wu J, Lin F, Han L, Liang X, Meng Q, Jiang Y, Wang Z, Yue A, Gu Y, Fan H, Zhou X, Liu Z. Fasudil preserves lung endothelial function and reduces pulmonary vascular remodeling in a rat model of end-stage pulmonary hypertension with left heart disease. *Int J Mol Med* 2018;**42**:1341–1352.
- Yin N, Kaestle S, Yin J, Hentschel T, Pries AR, Kuppe H, Kuebler WM. Inhaled nitric oxide versus aerosolized iloprost for the treatment of pulmonary hypertension with left heart disease. *Crit Care Med* 2009;**37**:980–986.
- Wood P, Besterman EM, Towers MK, McIlroy MB. The effect of acetylcholine on pulmonary vascular resistance and left atrial pressure in mitral stenosis. *Br Heart J* 1957;**19**:279–286.
- Ontkean M, Gay R, Greenberg B. Diminished endothelium-derived relaxing factor activity in an experimental model of chronic heart failure. *Circ Res* 1991;**69**:1088–1096.
- Fayyaz AU, Edwards WD, Maleszewski JJ, Konik EA, DuBrock HM, Borlaug BA, Frantz RP, Jenkins SM, Redfield MM. Global pulmonary vascular remodeling in pulmonary hypertension associated with heart failure and preserved or reduced ejection fraction. *Circulation* 2018;**137**:1796–1810.
- Guazzi M, Naeije R. Pulmonary hypertension in heart failure. *J Am Coll Cardiol* 2017;**69**:1718–1734.
- Dupuis J, Guazzi M. Pathophysiology and clinical relevance of pulmonary remodelling in pulmonary hypertension due to left heart diseases. *Can J Cardiol* 2015;**31**:416–429.
- Fernández AI, Yotti R, González-Mansilla A, Mombiola T, Gutiérrez-Ibanes E, Pérez del Villar C, Navas-Tejedor P, Chazo C, Martínez-Legazpi P, Fernández-Avilés F, Bermejo J. The biological bases of group 2 pulmonary hypertension. *Int J Mol Sci* 2019;**20**:5884.
- Chowdhury MA, Moukarbel GV, Gupta R, Frank SM, Anderson AM, Liu LC, Khouri SJ. Endothelin 1 is associated with heart failure hospitalization and long-term mortality in patients with heart failure with preserved ejection fraction and pulmonary hypertension. *Cardiology* 2019;**143**:124–133.
- Olschewski A, Veale EL, Nagy BM, Nagaraj C, Kwapiszewska G, Antigny F, Lambert M, Humbert M, Czirkák G, Enyedi P, Mathie A. TASK-1 (KCNK3) channels in the lung: from cell biology to clinical implications. *Eur Respir J* 2017;**50**:1700754.
- Lambert M, Capuano V, Olschewski A, Sabourin J, Nagaraj C, Girerd B, Weatherald J, Humbert M, Antigny F. Ion channels in pulmonary hypertension: a therapeutic interest? *Int J Mol Sci* 2018;**19**:3162.
- Lambert M, Boet A, Rucker-Martin C, Mendes-Ferreira P, Capuano V, Hatem S, Adão R, Brás-Silva C, Hautefort A, Michel J-B, Dorfmueller P, Fadel E, Kotsimbos T, Price L, Jourdon P, Montani D, Humbert M, Perros F, Antigny F. Loss of KCNK3 is a hallmark of RV hypertrophy/dysfunction associated with pulmonary hypertension. *Cardiovasc Res* 2018;**114**:880–893.
- Navas Tejedor P, Tenorio Castaño J, Palomino Doza J, Arias Lajara P, Gordo Trujillo G, López Meseguer M, Román Broto A, Lapunzina Abadía P, Escibano Subia P. An homozygous mutation in KCNK3 is associated with an aggressive form of hereditary pulmonary arterial hypertension. *Clin Genet* 2017;**91**:453–457.
- Southgate L, Machado RD, Gräf S, Morrell NW. Molecular genetic framework underlying pulmonary arterial hypertension. *Nat Rev Cardiol* 2020;**17**:85–95.
- Lambert M, Capuano V, Boet A, Tesson L, Bertero T, Nakhleh MK, Remy S, Anegón I, Pechoux C, Hautefort A, Rucker-Martin C, Manoury B, Domergue V, Mercier O, Girerd B, Montani D, Perros F, Humbert M, Antigny F. Characterization of *Kcnk3*-mutated rat, a novel model of pulmonary hypertension. *Circ Res* 2019;**125**:678–695.

19. Xiong PY, Tian L, Dunham-Snary KJ, Chen K-H, Mewburn JD, Neuber-Hess M, Martin A, Dasgupta A, Potus F, Archer SL. Biventricular increases in mitochondrial fission mediator (MifD51) and proglycolytic pyruvate kinase (PKM2) isoform in experimental group 2 pulmonary hypertension-novel mitochondrial abnormalities. *Front Cardiovasc Med* 2019;**5**:195.
20. Antigny F, Mercier O, Humbert M, Sabourin J. Excitation-contraction coupling and relaxation alteration in right ventricular remodelling caused by pulmonary arterial hypertension. *Arch Cardiovasc Dis* 2020;**113**:70–84.
21. Hohendanner F, Messroghli D, Bode D, Blaschke F, Parwani A, Boldt L-H, Heinzel FR. Atrial remodelling in heart failure: recent developments and relevance for heart failure with preserved ejection fraction. *ESC Heart Fail* 2018;**5**:211–221.
22. West JB, Dollery CT, Heard BE. Increased pulmonary vascular resistance in the dependent zone of the isolated dog lung caused by perivascular edema. *Circ Res* 1965;**17**:191–206.
23. Johnson DE, O'Keefe RA, Grandis JR. Targeting the IL-6/JAK/STAT3 signalling axis in cancer. *Nat Rev Clin Oncol* 2018;**15**:234–248.
24. Ellinghaus P, Scheubel RJ, Dobrev D, Ravens U, Holtz J, Huettner J, Nielsch U, Morawietz H. Comparing the global mRNA expression profile of human atrial and ventricular myocardium with high-density oligonucleotide arrays. *J Thorac Cardiovasc Surg* 2005;**129**:1383–1390.
25. Hancox JC, James AF, Marrion NV, Zhang H, Thomas D. Novel ion channel targets in atrial fibrillation. *Expert Opin Ther Targets* 2016;**20**:947–958.
26. Liang B, Soka M, Christensen AH, Olesen MS, Larsen AP, Knop FK, Wang F, Nielsen JB, Andersen MN, Humphreys D, Mann SA, Huttner IG, Vandenberg JJ, Svendsen JH, Haunsø S, Preiss T, Seeböhm G, Olesen S-P, Schmitt N, Fatkin D. Genetic variation in the two-pore domain potassium channel, TASK-1, may contribute to an atrial substrate for arrhythmogenesis. *J Mol Cell Cardiol* 2014;**67**:69–76.
27. Reiser PJ, Portman MA, Ning X-H, Moravec CS. Human cardiac myosin heavy chain isoforms in fetal and failing adult atria and ventricles. *Am J Physiol Heart Circ Physiol* 2001;**280**:H1814–H1820.
28. Sergeeva IA, Christoffels VM. Regulation of expression of atrial and brain natriuretic peptide, biomarkers for heart development and disease. *Biochim Biophys Acta* 2013;**1832**:2403–2413.
29. Zhao S, Wu H, Xia W, Chen X, Zhu S, Zhang S, Shao Y, Ma W, Yang D, Zhang J. Periostin expression is upregulated and associated with myocardial fibrosis in human failing hearts. *J Cardiol* 2014;**63**:373–378.
30. Bertero T, Handen AL, Chan SY. Factors associated with heritable pulmonary arterial hypertension exert convergent actions on the mir-130/301-vascular matrix feedback loop. *Int J Mol Sci* 2018;**19**:2289.
31. González-González L, Alonso J. Periostin: a matricellular protein with multiple functions in cancer development and progression. *Front Oncol* 2018;**8**:225.
32. Bruns DR, Tatman PD, Kalkur RS, Brown RD, Stenmark KR, Buttrick PM, Walker LA. The right ventricular fibroblast secretome drives cardiomyocyte dedifferentiation. *PLoS ONE* 2019;**14**:e0220573.
33. Duan W, Hicks J, Makara MA, Ilkayeva O, Abraham DM. TASK-1 and TASK-3 channels modulate pressure overload-induced cardiac remodeling and dysfunction. *Am J Physiol Heart Circ Physiol* 2020;**318**:H566–H580.
34. Nishimura K, Oydanic M, Zhang J, Babici D, Fraidenraich D, Vatner DE, Vatner SF. Rats are protected from the stress of chronic pressure overload compared with mice. *Am J Physiol Regul Integr Comp Physiol* 2020;**318**:R894–R900.
35. Teramoto K, Sengelov M, West E, Santos M, Nadruz W, Skali H, Shah AM. Association of pulmonary hypertension and right ventricular function with exercise capacity in heart failure. *ESC Heart Fail* 2020;**7**:1635–1644.
36. Harada D, Asanoi H, Noto T, Takagawa J. The impact of right ventricular dysfunction on the effectiveness of beta-blockers in heart failure with preserved ejection fraction. *J Cardiol* 2020;**76**:325–334.
37. Lejeune S, Roy C, Ciocea V, Slimani A, C de M, Amzulescu M, Pasquet A, Vancraeynest D, Beauloye C, Vanoverschelde J-L, Gerber BL, Pouleur A-C. Right ventricular global longitudinal strain and outcomes in heart failure with preserved ejection fraction. *J Am Soc Echocardiogr* 2020;**33**:973–984.e2.
38. Tian L, Xiong PY, Alizadeh E, Lima PDA, Potus F, Mewburn J, Martin A, Chen K-H, Archer SL. Supra-coronary aortic banding improves right ventricular function in experimental pulmonary arterial hypertension in rats by increasing systolic right coronary artery perfusion. *Acta Physiol* 2020;**229**:e13483.
39. Herron CR, Lowery AM, Hollister PR, Reynolds AB, Vincent PA. p120 regulates endothelial permeability independently of its NH2 terminus and Rho binding. *Am J Physiol Heart Circ Physiol* 2011;**300**:H36–H48.
40. Sidney LE, Branch MJ, Dunphy SE, Dua HS, Hopkinson A. Concise review: evidence for CD34 as a common marker for diverse progenitors. *Stem Cells* 2014;**32**:1380–1389.
41. Dierick F, Héry T, Hoareau-Coudert B, Mougenot N, Monceau V, Claude C, Crisan M, Besson V, Dorfmueller P, Marodon G, Fadel E, Humbert M, Yaniz-Galende E, Hulot J-S, Marazzi G, Sassoon D, Soubrier F, Nadaud S. Resident PW1⁺ progenitor cells participate in vascular remodeling during pulmonary arterial hypertension. *Circ Res* 2016;**118**:822–833.
42. Antigny F, Hautefort A, Meloche J, Belacel-Ouari M, Manoury B, Rucker-Martin C, Péchoux C, Potus F, Nadeau V, Tremblay E, Ruffenach G, Bourgeois A, Dorfmueller P, Breuils-Bonnet S, Fadel E, Ranchoux B, Jourdon P, Girerd B, Montani D, Provencher S, Bonnet S, Simonneau G, Humbert M, Perros F. Potassium channel subfamily K member 3 (KCNK3) contributes to the development of pulmonary arterial hypertension. *Circulation* 2016;**133**:1371–1385.
43. Anandharajan R, Eric A, Harikrishna T, Ling Y, Gladson M, Christy M, Sheila S, Tom B, Santhi G, Taylor S, Rizwan H, Susan M, James W. Abstract 16143: inflammation triggers the onset of hereditary pulmonary arterial hypertension in *kcnk3*^{-/-} animals. *Circ Am Heart Assoc* 2019;**10**:A16143.
44. Ranchoux B, Nadeau V, Bourgeois A, Provencher S, Tremblay É, Omura J, Coté N, Abu-Alhayja'a R, Dumais V, Nachbar RT, Tastet L, Dahou A, Breuils-Bonnet S, Marette A, Pibarot P, Dupuis J, Paulin R, Bouché O, Archer SL, Bonnet S, Potus F. Metabolic syndrome exacerbates pulmonary hypertension due to left heart disease. *Circ Res* 2019;**125**:449–466.
45. Borlaug BA, Obokata M. Is it time to recognize a new phenotype? Heart failure with preserved ejection fraction with pulmonary vascular disease. *Eur Heart J* 2017;**38**:2874–2878.
46. Miranda-Silva D, Gonçalves-Rodrigues P, Almeida-Coelho J, Hamdani N, Lima T, Conceição G, Sousa-Mendes C, Moura C, González A, Díez J, Linke WA, Leite-Moreira A, Falcão-Pires I. Characterization of biventricular alterations in myocardial (reverse) remodelling in aortic banding-induced chronic pressure overload. *Sci Rep* 2019;**9**:2956.
47. Grossman NL, Fiack CA, Weinberg JM, Rybin DV, Farber HW. Pulmonary hypertension associated with heart failure with preserved ejection fraction: acute hemodynamic effects of inhaled iloprost. *Pulm Circ* 2015;**5**:198–203.
48. Lai Y-C, Tabima DM, Dube JJ, Hughan KS, Vanderpool RR, Goncharov DA, St. Croix CM, Garcia-Ocaña A, Goncharova EA, Tofovic SP, Mora AL, Gladwin MT. SIRT3–AMP-activated protein kinase activation by nitrite and metformin improves hyperglycemia and normalizes pulmonary hypertension associated with heart failure with preserved ejection fraction. *Circulation* 2016;**133**:717–731.
49. Konecny F. Left atrial pressure measurement in a rat is currently impossible due to size limitations of balloon occlusion catheter. *J Thorac Cardiovasc Surg* 2018;**156**:1160–1161.
50. Mehra P, Mehta V, Sukhija R, Sinha AK, Gupta M, Girish MP, Aronow WS. Pulmonary hypertension in left heart disease. *Arch Med Sci* 2019;**15**:262–273.

Translational perspectives

Pulmonary hypertension (PH) due to left heart disease (PH-LHD) has higher incidence and prevalence than other forms of PH. However, no pulmonary-targeted drugs are available in clinical practice. The difficulties to have access to PH-LHD patient tissues, and the small number of preclinical models available compromise our understanding of the molecular mechanisms underlying PH-LHD. Using unique *Kcnk3*-loss-of-function-mutated rats, the present results demonstrate for the first time that *KCNK3* loss-of-function is a key event in the pathobiology of experimental PH due to AAC, suggesting that the *Kcnk3* channel could play a potential key role in the development of PH due to LHD.