
Brasier, M.J., Constable, Andrew John, Melbourne-Thomas, Jessica, Trebilco, Rowan, Griffiths, H.J., Van De Putte, Anton, & Sumner, Michael

Observations and models to support the first Marine Ecosystem Assessment for the Southern Ocean (MEASO)



Article. Version publiée - Published version.

© ELSEVIER

Citation APA:

Brasier, M., Constable, A. J., Melbourne-Thomas, J., Trebilco, R., Griffiths, H., Van De Putte, A., & Sumner, M. (2019). Observations and models to support the first Marine Ecosystem Assessment for the Southern Ocean (MEASO). *Journal of marine systems*, 197, 103182. doi:10.1016/j.jmarsys.2019.05.008

DOI: 10.1016/j.jmarsys.2019.05.008

Also available at: <http://hdl.handle.net/2013/ULB-DIPOT:oai:dipot.ulb.ac.be:2013/337820>

Cet article publié par ELSEVIER provient du Dépôt institutionnel de l'Université Libre de Bruxelles, DI-fusion <http://difusion.ulb.ac.be>. Il n'est accessible qu'aux membres de la communauté universitaire de l'ULB sur le réseau sécurisé de l'ULB.

Tout utilisateur autorisé peut lire, télécharger ou reproduire cet article à des fins d'usage privé ou à des fins non commerciales d'enseignement ou de recherche scientifique. Il ne peut être atteint à l'intégrité de l'article, et les noms des auteurs et de l'éditeur doivent être conservés. Tout téléchargement systématique des articles publiés par ELSEVIER mis à disposition dans DI-fusion est interdit.

This article published by ELSEVIER comes from the Institutional repository of Université Libre de Bruxelles, DI-fusion <http://difusion.ulb.ac.be>. It is accessible only to the members of the university community of ULB on the ULB secure network.

Any authorized user may read, download or reproduce this article for private usage, or for non commercial research or educational purposes. The integrity of the article and identification of the author and copyright owner must be preserved. Systematic downloading of articles published by ELSEVIER that are available in DI-fusion is not permitted.



ELSEVIER

Contents lists available at ScienceDirect

Journal of Marine Systems

journal homepage: www.elsevier.com/locate/jmarsys

Observations and models to support the first Marine Ecosystem Assessment for the Southern Ocean (MEASO)



M.J. Brasier^{a,*}, A. Constable^{a,b}, J. Melbourne-Thomas^{a,b}, R. Trebilco^a, H. Griffiths^c,
A. Van de Putte^{d,e}, M. Sumner^b

^a Antarctic Climate and Ecosystems Cooperative Research Centre, 20 Castray Esplanade, Hobart, Australia

^b Australian Antarctic Division, 203 Channel Highway, Kingston, Australia

^c British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 0ET, United Kingdom of Great Britain and Northern Ireland

^d Department of Biology, KU Leuven, Charles Deberiotstraat 32, Leuven, Belgium

^e OD Nature, Royal Belgian institute for Natural Sciences, Rue Vautierstraat 29, B-1000 Brussels

ARTICLE INFO

Keywords:

Antarctica
Conservation
Ecosystem-based management
CCAMLR

ABSTRACT

Assessments of the status and trends of habitats, species and ecosystems are needed for effective ecosystem-based management in marine ecosystems. Knowledge on imminent ecosystem changes (climate change impacts) set in train by existing climate forcings are needed for adapting management practices to achieve conservation and sustainability targets into the future. Here, we describe a process for enabling a marine ecosystem assessment (MEA) by the broader scientific community to support managers in this way, using a MEA for the Southern Ocean (MEASO) as an example. We develop a framework and undertake an audit to support a MEASO, involving three parts. First, we review available syntheses and assessments of the Southern Ocean ecosystem and its parts, paying special attention to building on the SCAR Antarctic Climate Change and Environment report and the SCAR Biogeographic Atlas of the Southern Ocean. Second, we audit available field observations of habitats and densities and/or abundances of taxa, using the literature as well as a survey of scientists as to their current and recent activities. Third, we audit available system models that can form a nested ensemble for making, with available data, circumpolar assessments of habitats, species and food webs. We conclude that there is sufficient data and models to undertake, at least, a circumpolar assessment of the krill-based system. The auditing framework provides the basis for the first MEASO but also provides a repository (www.SOKI.aq/display/MEASO) for easily amending the audit for future MEASOs. We note that an important outcome of the first MEASO will not only be the assessment but also to advise on priorities in observations and models for improving subsequent MEASOs.

1. Introduction

Assessments of the state of marine ecosystems and the causes of change in these systems are becoming very important for marine nations and internationally (e.g. Nymand-Larson et al., 2014; Constable et al., 2017). They will enable managers to understand how change in habitats, species, communities, and foodwebs (hereafter referred to collectively as ‘ecosystem changes’) may give rise to change in marine ecosystem services. Moreover, managers need to consider the potential for multiple causes of change from different societal uses (sectors) of marine ecosystems; there is an increasing need to develop multi-sectoral management systems that can appropriately adjust those sectors causing change. An imminent and pressing challenge is to develop management systems that will facilitate the adaptation of sectors to

expected future changes, such as those caused by climate change and ocean acidification.

At present, marine ecosystem assessments are mostly undertaken on a case-by-case basis when managing individual, or a small set of, ‘activities’ such as fisheries, pollution, and coastal engineering. They are generally based on empirical assessments from field observations. Typically, the combined effects across all activities are not directly assessed and managed, nor are the potential effects of climate change and ocean acidification included in these assessments individually or collectively. The latter effects are more often considered separately and heuristically based on reviews of disparate results in the existing peer-reviewed scientific literature, which we term ‘derivative assessments’. The most comprehensive derivative assessments for the marine environment are those by the Intergovernmental Panel on Climate Change

* Corresponding author.

E-mail address: madeleine.brasier@utas.edu.au (M.J. Brasier).

<https://doi.org/10.1016/j.jmarsys.2019.05.008>

Received 17 January 2019; Received in revised form 30 April 2019; Accepted 21 May 2019

Available online 27 May 2019

0924-7963/ © 2019 Elsevier B.V. All rights reserved.

(IPCC) and the United Nations World Ocean Assessment.

A scientific process is needed that directly assesses the potential for combined and cumulative effects and, particularly, can examine how those effects might continue in the short-to medium term future. The reason for assessing the future is that many effects may not be evident at the time of the assessment, although the drivers may have set them in train and made them unavoidable in the future. In the case of climate change and ocean acidification and based on the Earth system changes wrought by the ozone hole and its recovery, these changes may be two to three decades hence (IPCC, 2014).

Several important issues arise when attempting to consider and manage multiple drivers of effects, and the resulting cascading changes in ecosystems. Firstly, climate change and ocean acidification are not solely 'bottom up' drivers impacting on productivity of marine ecosystems. Species other than primary producers may be affected by changes in the physical and chemical systems. As a result, there may be 'top down' drivers that cause shifts in the structure and function of ecosystems as well (e.g. Johnson et al., 2011). Secondly, some parts of the ecosystem will be better studied than others because science tends to be more focussed on species or processes of direct interest to specific activities, such as fisheries. Lastly, future trajectories of the ecosystem may be difficult to foresee due to short time series of data or insufficient data to make empirical projections under climate change scenarios.

Ecosystems models provide a means to overcome these issues, they can be used in conjunction with time series data to validate and improve future predictions. In addition, they enable the integration of disparate datasets and knowledge of processes in order to examine the interactions of effects from different activities and from climate change and ocean acidification (Melbourne-Thomas et al., 2017). While there is an ongoing need to reduce uncertainty in our understanding of ecosystem function and to better incorporate this understanding into management-oriented models, these models will be central to developing realistic scenarios for the future (Constable et al., 2017).

1.1. What is a marine ecosystem assessment?

A marine ecosystem assessment (MEA) aims to bring together available data and knowledge from the scientific literature and different management bodies to, with the aid of models where possible, assess ecosystem status and change. Where possible, the relative importance of different stressors in causing that change will be assessed. Assessments of change will include historical change to the present, as well as providing realistic projections of change into the short-to-medium term future. The results are envisaged to directly support end-users, particularly policy makers, in adapting their work to ecosystem changes that may not be readily apparent in their jurisdiction but could impact on their objectives. Thus, a MEA aims to provide an overarching and integrated assessment, which has the flexibility and coverage enabling it to be adapted and useful to the needs of individual end-users, as well as providing context for derivative assessments and fisheries stock assessments (Fig. 1).

Here, we use the Southern Ocean as a case study to illustrate the information that would be compiled and the methods used to develop an integrated, overarching assessment – a Marine Ecosystem Assessment for the Southern Ocean (MEASO). The need for regular assessments of change of marine ecosystems around Antarctica and in the Southern Ocean (ASO ecosystems) has been identified by the Antarctic Treaty Consultative Meeting (ATCM) (ATCM, 2015), the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) (CCAMLR, 2015; SC-CCAMLR, 2011), the Scientific Committee on Antarctic Research (SCAR) (Kennicutt II et al., 2014; Turner et al., 2009; Turner et al., 2014) and in the work of the IPCC (IPCC, 2014; Nymand-Larson et al., 2014). A MEASO is intended to be a consensus report on status and trends in Southern Ocean habitats, species and foodwebs, drawing on the experience, results and methods

of the broader ASO research community. Fig. 2 illustrates how a regular MEASO process is intended to interact with policy makers. The MEASO cycle would operate similar to an IPCC cycle, over 5–10 years. The work would be expected to benefit from published syntheses and collaborations amongst researchers across the spectrum of existing research activities, as well as from observations from long-term monitoring programmes, e.g. the Southern Ocean Observing System (SOOS). Where possible, statistical and dynamic modelling would be used to assess the status and trends of habitats, species and food webs. The synthesis report would then provide summaries for use by policy-makers and other end-users. In addition, the synthesis would identify important priorities for advancing future assessments.

The first MEASO begins half way through the cycle and aims, through implementation of a first assessment, to establish processes and priorities for more comprehensive MEASOs in future. In this paper, we provide an 'audit' of the materials and methods available for the first MEASO. While this audit is comprehensive it is not intended to be exhaustive; our focus is on establishing a framework (using the Southern Ocean Knowledge and Information Wiki, www.soki.aq, as a repository) that can easily be amended and updated for future assessments.

Specifically, we start by summarising existing syntheses and derivative assessments on status and trends of habitats, species and food webs, including establishing spatial and temporal scales of reporting. Secondly, we document the types of observations available for assessing status and analysing trends, as well as the types of information that can be assembled for better understanding the pressures on different taxa in the ecosystem. In this section, we consulted researchers to better understand the data available and the scope of species-specific and assemblage-level assessments that may be undertaken. Thirdly, we summarise the status of models that could be used in a MEASO. Lastly, we identify the scope of the analyses that might be undertaken now, without substantially more research effort, to assess the status and trends in Southern Ocean ecosystems in a MEASO. Thus the primary purpose of this manuscript is to assess the information and data we have available to produce the first MEASO. In this instance we do not aim to undertake a detailed gap analysis or to provide a comprehensive set of recommendations for improving data collection and coverage for future MEASOs, as these will be key activities later in the MEASO process, outlined in the future directions section of this manuscript.

2. Syntheses of ecosystem status and trends

2.1. SCAR Antarctic Climate Change and the Environment report

The Scientific Committee on Antarctic Research (SCAR) has supported a number of efforts to provide syntheses on Antarctic and Southern Ocean science for policy makers. The Antarctic Environments Portal is one such initiative (<https://www.environments.aq>), where smaller syntheses may be found. A substantial synthesis on the effects of climate change on ASO systems – Antarctic Climate Change and the Environment (ACCE) - was undertaken by SCAR as part of the 2007–2009 International Polar Year (IPY) (Turner et al., 2009). The aim of ACCE was to describe how the physical climate system of Antarctica has varied over geological time and how environmental change during the instrumental period may affect biota (Convey et al., 2009). The 2009 report was largely focused on the West Antarctic Peninsula and how changes in sea-ice and primary production may affect ice-dependent species such as penguins and krill, as well the sensitivity of biota in other habitats, including benthos. Each year SCAR prepares updates of the ACCE report that highlight the new advances in our knowledge relevant to different sections of the report and provide some direction on priorities for future research; updates are presented at the Antarctic Treaty Consultative Meeting and published online (see <https://www.scar.org/policy/acce-updates/> for all ACCE updates).




Fig. 1. Comparison of the types of assessments undertaken for marine ecosystems, including the roles of observations and models within the workflows of the different assessments. Marine Ecosystem Assessments provide context for derivative and stock assessments (sideways arrows). In this example we use Antarctica as a region of interest.

2.2. Scientific Committee for the Conservation of Antarctic Marine Living Resources

The Scientific Committee for the Conservation of Antarctic Marine Living Resources (SC-CAMLR) has undertaken ecosystem syntheses on two occasions. In 2004, it held a Workshop on Plausible Ecosystem

Models for testing approaches to krill management (SC-CAMLR, 2004). In 2008, it held a joint workshop with the Scientific Committee of the International Whaling Commission on Input Data for Antarctic Marine Ecosystem Models (SC-CAMLR, 2008). Expert groups provided syntheses on different taxa for publication in CCAMLR Science in 2012, including on phytoplankton (Strutton et al., 2012), zooplankton




Fig. 2. The processes and work flow in a Marine Ecosystem Assessment (MEA) using the Marine Ecosystem Assessment for the Southern Ocean (MEASO) regarding the management region around Antarctica as an example. The dashed box in the lower right corner indicates the starting point of the first MEASO, to which this paper contributes by providing an audit of available knowledge, data, syntheses and models. Future audits might include identifying new or advanced data sets, assessment methods and models, and any assessments that may have been undertaken since the previous MEASO. The dashed box in top left corner demonstrates the potential interaction of MEAs with policy-makers. MEASO is envisaged to be an ongoing process, where each MEASO will advise on priorities for future research and monitoring to improve subsequent MEASOs.

(Atkinson et al., 2012a), krill (Atkinson et al., 2012b), fish as predators of krill (Kock et al., 2012), ice-breeding seals (Southwell et al., 2012), and penguins (Ratcliffe and Trathan, 2012).

2.3. SCAR Biogeographic Atlas of the Southern Ocean

The SCAR Biogeographic Atlas of the Southern Ocean (De Broyer et al., 2014) was produced using data collected during the Census of Antarctic Marine Life (CAML) voyages in the IPY 2007–2009 (content and data available at <http://data.biodiversity.aq/>). It is one of the major contributors to our current knowledge of the biodiversity and biogeography of Southern Ocean biota. The Atlas collates 1.07 million occurrence records of 9064 validated species from ~434,000 distinct sampling stations; these are fundamental data in providing the necessary geospatial framework for marine biodiversity knowledge and understanding, and for assessing its gaps (De Broyer et al., 2014). The Atlas is now regarded as a milestone product of 21st Century Antarctic Science (De Broyer and Koubii, 2015), it has been cited in 95 publications between 2015 and 2018 according to Google Scholar. It has contributed to major publications reviewing knowledge of climate change impacts on Antarctic ecosystems (e.g. Constable et al., 2014a; Chown et al., 2015) and potential ecological change under future conditions (Griffiths et al., 2017). In addition it has been used to advise future monitoring, management and conservation of ASO ecosystems (e.g. Gutt et al., 2017; Koubbi et al., 2017; Cavanagh et al., 2016; Constable et al., 2016; Xavier et al., 2016) including contributing to supporting scientific information for spatial management measures in CCAMLR (Teschke et al., 2014).

Table 1 summarises how the Atlas may be updated with more recent information. Trends in new research across taxa within the ASO include taxonomic, biogeographic, ecological and physiological studies. Information provided by the Atlas editorial team, many of the original lead authors of the chapters of the Atlas, along with literature reviews that we undertook was summarised into 7 sub-topics in the table. Not surprisingly, the advances in genetic technologies have resulted in many of the taxonomic groups undergoing revision, as well as enabling better understanding of spatial populations structures and food web linkages in the region. The Atlas team are in the process of creating an online version of the Atlas, which will display the original content of the Atlas but with integrated R code (in Bookdown, <https://bookdown.org/yihui/bookdown/>) to map the most recent records.

2.4. Other reports

A number of bodies under the auspices of the United Nations have developed syntheses on ASO ecosystems. These include the IPCC and Regular Process for Global Reporting and Assessment of the State of the Marine Environment (World Ocean Assessment). The most recent regional reports from each are those by IPCC Working Group II on Polar Regions in 2014 (Nyman-Larson et al., 2014) and by the World Ocean Assessment in 2016 on high-latitude ice and the biodiversity dependent upon it (Rice and Marschoff, 2016). An IPCC Special Report on the Oceans and Cryosphere in a Changing Climate has a Polar Regions chapter examining the effects of climate change on polar regions, including ecosystems due for release in 2020. Similar works have been conducted for the Arctic; the Conservation of Arctic Flora and Fauna State of the Arctic Marine Biodiversity Report has also investigated detectable changes and gaps in our ability to assess status and trends in Arctic marine ecosystems under changing conditions.

Several major research projects have also resulted in dedicated journal issues with specific publications on different taxa and processes (e.g. Brandt and Ebbe, 2007; Hofmann et al., 2011) while others have focused on the physical environmental changes and how these may affect biota now (e.g. Rogers et al., 2012; Murphy et al., 2013; Constable et al., 2014a; Chown et al., 2015) and under future conditions (Griffiths et al., 2017). Some papers have developed syntheses

along with advice for future monitoring, management and conservation of ASO ecosystems (e.g. Xavier et al., 2016; Cavanagh et al., 2016; Constable et al., 2016; Gutt et al., 2017; Koubbi et al., 2017).

3. Observations to support MEASO

3.1. Field programmes

Scientific observations within the ASO commenced in the late 1800s with the first Challenger expedition, shortly followed by the initiatives of the first International Polar Year and the Belgica and Discovery expeditions (Fig. 3). These early expeditions formed the foundation for benthic and pelagic species records in the Southern Ocean (Griffiths, 2010). In 1981 the first large-scale international research project, BIOMASS (Biological Investigations of Marine Antarctic Systems and Stocks) took place (El-Sayed, 1994). In 1978 satellite technologies were implemented to observe the variability and trends in Antarctic sea ice (Cavalieri and Parkinson, 2008). Since then satellites have also been used to monitor sea surface temperature, ocean topography and ocean colour (a proxy for chlorophyll concentration, Johnson et al., 2013). These combined with ongoing oceanographic research assist in characterising the changing pelagic habitats of the ASO.

Many ASO research programmes have targeted Antarctic krill and krill-dependent predators, especially whales. For example, the BROKE and BROKE-West expeditions were designed to improve our understanding of krill dynamics within east Antarctica (Nicol et al., 2000a, 2010) while the CCAMLR-2000 Survey (also CCAMLR-2000 Krill Synoptic Survey) was initiated to improved estimates of krill biomass in the Atlantic sector of the Southern Ocean (Trathan et al., 2001). The outcomes of these programmes were used by CCAMLR to set precautionary catch limits for the krill fishery (Hewitt et al., 2004). CCAMLR has established an ecosystem monitoring programme (CEMP) for monitoring krill-dependent species, which at present focusses on land-based predators (Agnew, 1997). It also provides for regular assessments of the status of fish stocks based on tagging and groundfish surveys (see Fishery Reports - <https://www.ccamlr.org/en/publications/fishery-reports>).

The CEMP was established in 1987 with national research agencies contributing data, as available, to the CCAMLR Secretariat. CEMP uses standardised methods to monitor 8 indicator species considered dependent on Antarctic krill. These species include the adélie penguin (*Pygoscelis adeliae*), chinstrap penguin (*P. antarctica*), gentoo penguin (*P. papua*), macaroni penguin (*Eudyptes chrysolophus*), black-browed albatross (*Thalassarche melanophrys*), Antarctic petrel (*Thalassoica antarctica*), cape petrel (*Daption capense*) and the Antarctic fur seal (*Arctocephalus gazella*). CEMP is developed from national contributions to individual programmes. General coordination is provided through the CCAMLR Working Group on Ecosystem Monitoring and Management.

Ecosystem-oriented research programmes and monitoring have been of increasing importance over the last few decades. Many are focussed on biogeochemistry or krill-based food webs. Long term observation programmes such as the Palmer Long Term Ecological Research (LTER, <http://pal.lternet.edu/>) and the Rothera Time Series (RaTS, <https://www.bas.ac.uk/project/rats/>) collect sustained observations within the vicinity of national research stations. These programmes collect oceanographic, biochemical and biological data to investigate inter-annual variation and climate change impacts on the Antarctic ecosystem. Other programmes are more fisheries oriented such as for the UK at South Georgia (krill, toothfish, icefish), Australia at Heard Island and McDonald Islands (toothfish, icefish) and Macquarie Island (toothfish), France at Kerguelen Islands (toothfish, icefish) and Crozet Islands (toothfish), South Africa at Prince Edward & Marion Islands (toothfish), and New Zealand, UK and Norway in the Ross Sea (toothfish). Land-based predators are extensively monitored on many subantarctic islands, including South Georgia, Crozet, and Kerguelen.

Table 1

Expected research findings by taxa achieved since the publication of the SCAR Biogeographic Atlas of the Southern Ocean (De Broyer et al., 2014). Shaded cells indicate new research/data available for certain taxa within the following categories: Taxonomic re-evaluation (previous taxonomic classifications have been altered); species discovery (previously undescribed morphological or cryptic species have been discovered); invasive species (species previously considered non-Antarctic have now been recorded within the Southern Ocean); species shift (evidence of species shifts within the ASO e.g. the poleward movement); sample coverage (additional samples are now available from previously un-sampled locations or un-sorted material increasing spatial coverage within the ASO); ecological (improved understanding of ecological traits, e.g. diet, habitat, reproduction, which may change distribution of taxa now or in the future); physiological (insights into physiological traits, e.g. acclimation or adaptation to changing temperature or acidity, which might influence distributions). Taxonomic experts who provided information for this table are acknowledged at the end of this manuscript.

Taxa	Taxonomic re-evaluation	Species discovery	Species shift	Sample Coverage	Ecological	Physiological
Polychaetes						
Bryozoa						
Ascidian						
Benthic hydroids						
Stylasteridae						
Antarctic hexacorals						
Harpacticoid copepods						
Pycnogonida						
Benthic Ostracoda						
Benthic amphipods						
Isopoda						
Cumacea						
Crabs and lobsters						
Shrimps						
Pelagic copepods						
Halocyprid ostracods						
Hyperiid amphipods						
Euphausiids						
Lysianassoidea amphipods						
Tanaidacea						
Asteroidea						
Crionoids						
Echinoids						
Fish						
Benthic foraminifera						
Gastropoda						
Bivalvia						
Octopuses						
Pteropods						
Squid						
Marine nematodes						
Porifera						
Gelatinous zooplankton						
Near surface zooplankton						
Tintinnid ciliates						
Macroalgae						
Sea-ice metazoans						

Many of the field programmes shown in Fig. 3 were international research efforts consisting of multiple expeditions within different regions of the ASO. The CAML, which ran from 2005 to 2010, coordinated 18 major research voyages to the Antarctic and the Southern Ocean during the 2007–2009 International Polar Year (Schiaparelli et al., 2013), many of which targeted unsampled regions, for example the deep-sea benthos within the Amundsen and Bellingshausen Seas (Linse et al., 2013). Overall the CAML voyages sampled about 350 sites within the ASO collecting pelagic, demersal and benthic fauna using a variety of sampling gears (Stoddart, 2010). The species data collected during CAML voyages were deposited in SCAR-MarBIN (Scientific Committee on Antarctic Research Marine Biodiversity Network, now <http://www.biodiversity.aq/>) data portal, containing data for over 14,000 species.

Since BIOMASS, Southern Ocean GLOBEC provided the impetus to develop internationally co-ordinated, integrated studies of the krill-based food web (Hofmann et al., 2011). In 2008, it morphed into the IMBER and SCAR programme, Integrating Climate and Ecosystem Dynamics in the Southern Ocean (ICED) (Murphy et al., 2008), with a continued focus on process studies, as well as a new emphasis in

developing ecosystem models (Murphy et al., 2012).


The SOOS was established as a partnership between SCAR and the Scientific Committee on Oceanic Research to develop sustained observing of essential physical, chemical and biological variables to underpin research and monitoring of the region (Rintoul et al., 2011; Meredith et al., 2013; Constable et al., 2016; Newman et al., accepted). Although in its infancy, SOOS is beginning to provide mechanisms for retrieving data for the purposes of a MEASO. Its development of regional working groups (Newman et al., accepted) will enable further implementation of co-ordinated field observations identified as important to MEASO in the future.

3.2. Taxon-level assessments

In its simplest form, an assessment of the status and trends of a species can be derived from abundance data of taxa over time (Constable et al., 2014b). Using the defined assessment components in Table 2, we review the relative spatial coverage of and, in the case of pelagic taxa, the observed trends in taxa-specific assessments within the

published literature (Table 3, Table 4). This is an important part of the MEASO process in order to establish an understanding of studies and data available to assess status and trends across the ASO and over time.

At this stage, we did not review the utility of the assessments for the purposes of MEASO; while our review here is not exhaustive, the number of assessments indicated for each taxonomic group indicate the



* Programmes have a circumpolar approach and outlook but individual surveys may have targeted specific regions or sites.
 ** Programmes not associated with the listed international monitoring schemes

(caption on next page)

Fig. 3. Timeline including often-cited examples of the major scientific observations within the Antarctic Southern Ocean ecosystem from early scientific observations and modern survey and sampling programmes. Including Scientific Committee for Antarctic Research (SCAR) lead programmes: BIOMASS - Biological Investigations of Marine Antarctic Systems and Stocks (1981–1991), SO-CPR – Southern Ocean Continuous Plankton Recorder (1991 onwards), EASIZ - Ecology of the Antarctic Sea Ice Zone (1994–2004), APIS - The International Antarctic Pack Ice Seals Programme. Commission for the Conservation of Antarctic Living resources (CCAMLR) programmes: CEMP - CCAMLR Ecosystem Monitoring Programme, BROKE - Baseline Research on Oceanography, Krill and the Environment. International Polar Year (IPY) programmes and the Census of Antarctic Marine Life (CAML) (2005–2010). International Whaling Commission (IWC) programmes: SOWER - Southern Ocean Whale and Ecosystems Research Programme (1978–2009), SORP - Southern Ocean Research Partnership. Regional working groups: SO-GLOBEC - Southern Ocean Global Ocean Ecosystem Dynamics, ICED - Integrating Climate and Ecosystem Dynamics in the Southern Ocean, SOOS - Southern Ocean Observing System. National long-term observation examples including: LTER - Long-term Ecological Research Programme, AMLR - Antarctic Marine Living Resources, RATS - Rothera Antarctic Time Series, SOCCOM - Southern Ocean Carbon and Climate Observations and Modelling project. Details of additional CEMP sites and ongoing monitoring see <https://www.ccamlr.org/en/science/cemp-sites>.

relative attention given to each group. The published assessments varied in the amount of data included. Some were derived from long-term data sets that may help identify trends in species abundance/density over time. Others were “snapshots” of a species that may be used as an indicator of status but not trends. Our results highlight the real differences in coverage between taxa and between sectors. Ideally a MEASO would be based on circumpolar assessments of abundance or density but, at present, this only exists for a limited number of species (Klekociuk and Wienecke, 2017). Only few locations/regions are well sampled across the spectrum of taxa. Some types of areas, such as the deep-sea benthos, are only poorly sampled (Brandt et al., 2014).

Some of the earliest species-level studies within the ASO focused on krill and marine mammals, dating back to the early 20th Century during the sealing and whaling eras. This was followed by a rise in scientific estimates of krill abundance between 1930 and 1980 (Pauly et al., 2000; Nicol et al., 2000b) and an interest in their potential relationship between krill abundance and large-scale oceanographic processes (for a review of early works see Priddle et al. (1988) and for the earliest fisheries data see Fedulov et al. (1996)). For birds and marine mammal's quantitative abundance data was scarce until the 1970s and close to non-existent prior to the 1950s (Croxall, 1992). Crude estimates of seal and whale populations are suggested in Laws

(1977). In many early works the methodologies are unpublished or are unreliable.

Species-level abundance or density data have been compiled in recent decades for a number of species, including zooplankton (from SO-CPR surveys; <https://www.scar.org/science/cpr/home/>), Antarctic krill (raw data in KRILLBASE; <https://www.bas.ac.uk/project/krillbase/#data>), Adelie penguins (colony counts; <http://www.penguinmap.com/>), whales (assessments; <https://iwc.int/status>), albatross (assessments; <https://acap.aq/acap-species?lang=en>), species with conservation assessments (IUCN red list; <https://www.iucn.org/resources/conservation-tools/iucn-red-list-threatened-species>), and in CCAMLR fishery reports by CCAMLR management area (<https://www.ccamlr.org/en/publications/fishery-reports> and supplementary information). Additional individual assessments have been submitted to the CCAMLR Working Group for Ecosystems Monitoring and Management. These can be found online and available on request; for example reports, submitted for the Predator Survey Workshop in 2008 include reports of fur seal, flying bird and penguins abundance (<https://www.ccamlr.org/en/wg-emm-psw-08>).

These sources have varying degrees of quality-control. For some datasets, limited repeat observations may make trends difficult to estimate. Attention may need to be given to interannual variation


Table 2
Definitions and categories of the assessed components in Tables 3 and 4.

Assessed component	Definition	Categories/criteria
Status assessments	The number of assessments of status (abundance, density) that we found publicly available per taxa within the Antarctic Southern Ocean. ^a	
Relative spatial coverage	The relative number of published assessments of status within each sector (or circumpolar) per taxa. ^a	Atlantic Indian West Pacific East Pacific Circumpolar
Relative depth coverage (benthic only)	The relative number of published assessments of status across depth categories. ^a	Shelf < 1000 m Shelf-slope = 0–3000 m Shelf-slope-basin = 0–> 3000 m Basin > 3000 m
Trend assessments (pelagic only)	The total number of status assessments over time within the Antarctic Southern Ocean. ^a	
Earliest data	An indication of the earliest abundance, density or biomass data.	
Observed trends (pelagic only)	An indication of the observed trends in the abundance, density or biomass of a taxa over time from the literature ^a	Increase (↑) = all published trends indicate an increase in abundance/density Decrease (↓) = all published trends indicate a decrease in abundance/density No change (–) = all published trends indicate no change in abundance/density Interannual variation (–) = all published trends indicate interannual variation in abundance/density Contrasting trends (?) = published trends vary within species, between species or with location No published trends (x) = no trends within the published literature for that taxa.

^a For exact number and links to references see Supplementary information.

Table 3

Review of the spatial coverage and observed trends of taxon-specific assessments for plankton, krill, fish and air breathing species in the Antarctic Southern Ocean. The definitions of each assessed component is outlined in Table 2. For relative spatial coverage; dark blue = Atlantic sector, light blue = Indian Sector, dark green = West Pacific sector, light green = East Pacific and yellow = circumpolar. A full list of references used to assess the relative spatial coverage and determine observed trends is available in the supplementary information.

Taxon	Status assessments	Relative spatial coverage	Trend assessments	Earliest data	Observed trends
Plankton 	155		-	1900s	Not assessed*
Krill 	40		20	1920s	~ ~ ~ ~
Mackerel Icefish 	5		3	1970s	~ ~ x x
Toothfish 	10		7	1980s	↓ ? ~ x
Baleen whales 	18		4	1970s	x ↑ ↑ x
Toothed whales 	5		1	1970s	x x ↓ x
Ice-breeding Seals 	28		7	1970s	? ? ↓ x
Elephant Seals 	24		21	1950s	↑ ? x ?
Fur Seals 	15		14	1950s	↑ ↑ x ↑
Adelie Penguin 	23		12	1940s	? ↑ ↑ ↓
Chinstrap Penguin 	16		10	1950s	↓ ~ x ↓
Emperor penguin 	17		6	1940s	x ↓ ~ ↓
Gentoo penguin 	10		6	1970s	↑ x x ?
Macaroni penguin 	6		3	1950s	↓ ? x x
Antarctic and Cape Petrels 	9		3	1960s	↓ ↓ x x
Black-browed Albatross 	18		5	1960s	x ? x x



Observed trends

- ↑ Increase
- ↓ Decrease
- No change
- ~ Interannual-variation
- ? Contrasting trends, local variation or variation between species
- X No published trends

* Sectoral observed trends in plankton not assessed due to high variation on local scales

Table 4

Review of the spatial and depth coverage of taxon-specific assessments for major benthic invertebrate taxonomic groups in the Antarctic Southern Ocean. The definitions of each assessed component is outlined in Table 2. For relative spatial coverage; dark blue = Atlantic sector, light blue = Indian Sector, dark green = West Pacific sector, light green = East Pacific and yellow = circumpolar. A full list of references used to assess the relative spatial and depth coverage is available in the supplementary information.

Taxon	Status assessments	Relative spatial coverage	Relative depth coverage	Earliest data
All benthos	 108			1960
Meiofauna	 3			1980
Macrofauna	 13			1970
Megafauna	 10			1980
Annelida	 9			1970
Cnidaria	 2			1980
Crustacea	 14			1970
Echinodermata	 4			1980
Foraminifera	 5			1970
Mollusca	 8			1970
Porifera	 5			1960



Depth Distribution

- Shelf (<1000 m)
- Shelf-slope (0 - 3000 m)
- Shelf-slope-basin (0 - >3000 m)
- Basin (>3000 m)

associated with El Niño–Southern Oscillation (Trathan et al., 2003; Meredith et al., 2005; Fielding et al., 2014). Importantly, inconsistencies between surveys and/or sampling biases of different survey methods may need to be accounted for through standardisation procedures. These issues have been important to resolve in existing datasets, including standardisation across sampling methods and spatial and temporal coverage for Antarctic krill (in KRILLBASE; Loeb and Santora, 2015; Cox et al., 2018), Adelie penguins (Adelie penguin census repositories; Southwell et al., 2013) and ice-breeding seals (APIS repositories; Southwell et al., 2012).

Data on Antarctic benthic communities has been assembled since the 1960s, recorded mostly from trawl, dredge, corer and camera data (Gutt et al., 2013). However, the density of taxa has often not been recorded (Downey et al., 2012); the difficulties in collecting quantitative benthic samples mean that many studies are only semi-quantitative (Clarke, 2008). Some equipment including the epibenthic sledge and camera technologies are able to generate quantitative abundance data for macro and megafauna species respectively (Gutt and Starmans, 2003; Brandt et al., 2007a; Post et al., 2017). Studies that assess trends

over time are rare, usually in shallow water habitats close to research stations (e.g. Conlan et al., 2004; Stark and Kim, 2014). Table 3 summarises the coverage of benthic assessments by depth and sector. To date the greatest number of benthic studies have been conducted in the Weddell Sea, around the West Antarctic Peninsula and Ross Sea (Gutt et al., 2013).

3.3. Consultation on research activities on density or abundance

In addition to the review of the literature and online sources, we consulted 92 scientists from 18 different countries for information on assessments of density/abundance of ASO taxa. The aim of this consultation was to determine the spatio-temporal coverage of research programmes estimating abundance (or relative density) of taxa within the Southern Ocean in each decade from 1980 to the present. A total of 14 broad taxonomic groups sub-divided into 49 monitoring groups were listed within the consultative document over 13 different sites within the ASO (full instructions, taxonomic groups and data are provided in supplementary material).




Fig. 4. Compiled community survey responses by taxa, location and region over time. Colour scheme for the relative number of national research programmes measuring species abundance: yellow (low 1–5 surveys/programmes), orange (medium 6–15 surveys/programmes); green (high > 15 surveys/programmes). Grey shading indicates additional data available from other circumpolar studies and databases including SO-GLOBEC (phytoplankton and pelagos) and IWC (toothed and baleen whales). Blue shading indicates data within CCAMLR sources including fishery assessments (bathypelagic and ground-fish) and from CEMP monitoring sites (penguins and fur seals). The four cells within each taxon indicate time by decade from 1980 to 2010. Further details of the survey are available in the supplementary information. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Completed responses were received from 30 individuals from 13 of the targeted countries including (number of responses): Argentina (1), Australia (5), Canada (1), Chile (2), France (1), Germany (5), India (1), Italy (4), Japan (1), Russia (1), South Africa (2), United Kingdom (3) and the USA (3). Additional information was also provided by New Zealand, Australia and the USA. Others indicated that they were not able to contribute or had already contributed to previous responses for their nation. These data are available on SOKI and contributors acknowledged at the end of this manuscript.

The greatest survey coverage, indicated by the highest number of research surveys or programmes, over time and taxa was recorded for the West Antarctic Peninsula, one of the most accessible regions of the ASO, while the main spatial gaps (fewer surveys or programmes) appear to be the Amundsen Sea, Bellingshausen Sea and Macquarie Ridge (Fig. 4). The number of surveys generally increased with time in most regions reflecting the increase in Antarctic research capacity with time. Across taxa, flying birds were mostly covered at locations near to coastal research stations, and benthic taxa were not well represented. Such spatial biases, inherent when studying the Southern Ocean, are


discussed in Griffiths et al. (2014). In previous research, the most intense sampling tends to be at the more easily accessible locations, e.g. close to research stations, or the WAP, and depths <1000 m. Taxonomic biases are somewhat easier to overcome; there was a surge in species data recorded in online databases during the SCAR Biogeographic Atlas project (Griffiths et al., 2011, 2014). However, we still lack abundance data for many groups. These differences could reflect both the nature of scientific programmes or, the relative success of the community-based survey approach.

4. Models to support MEASO

Models underpin the scientific method (Peters, 1991). The term ‘model’ is used in many ways (see Melbourne-Thomas et al., 2017), ranging from (i) heuristic discussions on a system and/or hypotheses of various complexities, to (ii) statistical models aimed at predicting the magnitude of one or more dependent variables based on a series of independent and related variables, to (iii) formal system-level structures linking objects (nodes – physical and chemical variables, species, human

Table 5

Different modelling approaches used within Antarctic and Southern Ocean (ASO) ecosystems from physical and biogeochemical to whole ecosystem models, anticipated utility within the MEASO project and current ASO coverage. For examples of model implementation in Antarctica see, for Qualitative models: [Melbourne-Thomas et al. \(2013\)](#), [Goedegebuure et al. \(2017\)](#). Earth System models: Reviewed in [Cavanagh et al. \(2017\)](#). Regional Physical models: [Dinniman et al. \(2011\)](#), [Corney et al. \(2019\)](#), [Mazloff et al. \(2010\)](#). Regional Biogeochemical models: [Pasquer et al. \(2005\)](#), [Saenz and Arrigo \(2014\)](#), [Vancoppenolle et al. \(2010\)](#), [Melbourne-Thomas et al. \(2015\)](#), [Priester et al. \(2017\)](#). Single species models: [Hofmann et al. \(1998\)](#), [Murphy et al. \(2004\)](#), [Thorpe et al. \(2007\)](#), [Wiedenmann et al. \(2008\)](#), [Jenouvrier et al. \(2014\)](#). Food web models: [Mori and Butterworth \(2004, 2005, 2006\)](#), [Pinkerton et al. \(2010\)](#), [Hill et al. \(2012\)](#), [Ballerini et al. \(2014\)](#), [Gurney et al. \(2014\)](#), [McCormack et al. \(in prep\)](#), [Subramaniam et al. \(in prep\)](#). Benthic models; [Johst et al. \(2006\)](#) [Potthoff et al. \(2006a, 2006b\)](#). Specific interaction models: [Constable \(2005, 2006, 2008\)](#), [Watters et al. \(2013\)](#), [Plagányi and Butterworth \(2007\)](#), [Klein et al. \(2018\)](#), [Tulloch et al. \(2017\)](#). End to end models: [Melbourne-Thomas et al. \(in prep\)](#).

Model type	Description	Examples	Anticipated utility	Implementation	ASO Coverage
Qualitative models 	Framework to examine ecosystem responses to press perturbation.	Qualitative network models	Understand linkages and feedback mechanisms.	West Antarctic Peninsula and aspatial	
Earth system 	Simulation of physical, chemical and biological processes within the earth system. Can incorporate global climate models.	CMIP5 models	Provides forcings for regional models.	Global with circum-Antarctic detail. Note, southern boundary may not be to the coast.	
Regional physical 	Simulation of physical conditions within the Southern Ocean such as temperature, salinity and currents.	Ocean General Circulation Models (OGCM) Regional Ocean Modelling System (ROMS) Southern Ocean State Estimate (SOSE)	Provides regional physical forcing for ecosystem models.	Ross Sea, West Antarctic Peninsula Indian sector	
Regional biogeochemical 	Simulation of biogeochemistry in the Southern Ocean e.g. nutrient cycling, carbon uptake, productivity	Nutrient, phytoplankton, zooplankton and detritus (NPZD)	Understand different drivers that control the base ASO productivity.	All sectors, including pelagic and in sea ice	
Single species 	Simulation to understand the ecology of a single species based on current biological knowledge and environmental setting.	Krill examples: Advection, recruitment, relationship with physical drivers e.g. sea ice and climatic variation etc	Filling gaps in space and time, where we have patchy abundance data.	Mostly commercially exploited species (krill, seals, whales); Scotia Arc, South Georgia, West Antarctic Peninsula, Ross Sea, Indian sector	
Foodweb models 	Simulation of the trophic interactions within an ecosystem from primary producers to higher predators. Used to investigate the impacts of changes in primary production, fishing effort and species loss.	Mass balance Ecopath with Ecosim. Size spectrum models	Representation of food entire food web to explore relative importance of trophic linkages and the relative impact of different climate and fishing scenarios.	Ross Sea, Scotia Arc, South Georgia, West Antarctic Peninsula, Indian sector	
Benthic models 	Simulation of habitat complexity that shapes biological communities in benthic ecosystems, and roles in benthic-pelagic coupling		Explore dynamics of benthic assemblages in relation to iceberg scour, environmental change and fisheries.	Weddell Sea, Scotia Sea Indian sector	
Specific interaction models 	Dynamic models of the interactions between selected species within the ecosystem. Can provide quantitative information on ecosystem performance for use in management of human activities including fishing.	FOOSA SMOM EPOC	Subset a food web to specific primary interactions for exploring effects of environmental change or fisheries scenarios.	Mostly krill and krill predators (penguins, whales). Scotia Sea, circum-Antarctic	
Socio-ecological 	Framework used to understand the interactions between societal, environmental and governmental factors.		To investigate policy-relevant scenarios that may contribute to adaptive conservation and management of marine social-ecological systems.	Yet to be implemented in Antarctica.	
End to end models 	These models include submodels on physics, chemistry, biology, human uses, economics and management. They are spatially structured and can resolve small time-steps (minutes) if needed. They enable exploration of direct and indirect effects of change in one or more components on the other elements of the system. Components can be modelled at different levels of complexity from pools to complex populations with behaviours.	Atlantis	Enable exploration of system-level scenarios of change as well as having methods for evaluating how well management and adaptation measures may work under various scenarios.	Currently being implemented for East Antarctica.	

uses) by processes (edges – trophic interactions, physiological responses, competitive interactions), the behaviour of which are forced by system drivers (variables – seasonality, ENSO, climate change, fisheries).

Hereafter, the latter system-level models are termed ‘system models’. In this section, we focus on the system models, regarding that statistical models, which include species distribution models, underpin the species-

specific analyses. System models are those that help identify causes and effects and consequent changes when the forcing variables change.

System models can be used to test outstanding hypotheses on the effects of change, develop plausible scenarios of current and future change given the data, and for undertaking more precise assessments of the status and trends of the ecosystem (and its likelihood) using estimation procedures (Murphy et al., 2012; Melbourne-Thomas et al., 2017). Ranging in complexity from single species to whole ecosystems (Table 5), system models provide scientists with a method for linking disparate studies on status of some important species with many other studies on processes and ecosystem interactions, thereby enabling complex system studies even though not all components of the system have been observed simultaneously. Thus, system models, couched in observations, can be used to explore the outcomes from multiple ecosystem interactions and perturbations and reporting the consequences to decision makers (Watters et al., 2013; Klein et al., 2018). With the rise of ecosystem-based management practices, which are supported by CCAMLR (Constable, 2004, 2011; Kock et al., 2007), the development of ecosystem models to investigate future climate, fishing and conservation scenarios are increasingly important (Gurney et al., 2014).

The main ecological and modelling challenges in the development of system models is summarised in Murphy et al. (2012). Some of the first ecological modelling applications within the Southern Ocean were based on Antarctic krill because of its importance to whales as well as its emerging importance as a target commercial species (see references in Hill et al., 2006). Antarctic krill is a relatively well studied species, with much information on its growth rate, transport, and population dynamics which can be incorporated into models (Siegel, 2016), however it will be important to ensure that future modelling approaches are flexible enough to allow representation of potential shifts to non-krill dominated ecosystem states (McCormack et al., in review, accepted; Trebilco et al., in review-a).

Early modelling studies investigated the interaction between krill aggregations and harvesting operations in attempt to utilise the krill catch rate as a proxy for abundance (Mangel, 1988; Butterworth, 1988) while conceptual models provided qualitative descriptions of the food-web and model multi-species interactions (for references see Hill et al., 2006). Qualitative network models have since been used to examine directional responses of ecosystem components to perturbations, including the mechanisms behind observed changes and the impacts of model complexity on results (Melbourne-Thomas et al., 2012, 2013). This approach provides a quick yet substantial insight into system functioning (Levins, 1996).

Quantitative food web and ecosystem models have also been developed to simulate responses to ecosystem perturbations including fishing (Fulton, 2010). These include the widely used Ecopath with Ecosim, a mass balance model with a time dynamic simulation based on the functional groups within an ecosystem (Christensen and Walters, 2004, see <https://ecopath.org> for model and software details). Pinkerton et al. (2010) used a similar framework to Ecopath but included key non-trophic transfers (e.g. seasonal release of material from sea-ice, vertical detrital flux) to investigate the ecosystem impacts of fishing in the Ross Sea.

More recently, end-to-end, or whole-ecosystem models, attempt to include all major relevant processes within the ecosystem, such as nutrient cycling, climate forcing, environmental variability and harvesting as well as representations of biological species/functional groups that include ecological processes such as feeding, growth, reproduction and dispersal (Fulton, 2010; Murphy et al., 2012). An end-to-end modelling framework, Atlantis (Fulton, 2010; Fulton et al., 2011) is currently under development for implementation in East Antarctica. This model will enable development of climate change scenarios for the regional ecosystem as well as evaluating different management and adaptation options for fisheries and other activities.

Fig. 5 illustrates how the different system models described in Table 5 might fit together in a nested, ensemble of models. While not all

the available models described in Table 5 will be used in the initial MEASO, the aim will be to utilise scenarios of environmental change from Earth System models (Cavanagh et al., 2017), along with time-series of observations of physics, chemistry and biology, to drive regional food web and/or species models. These latter models can then be used to investigate the consequences, and their likelihoods, of the different scenarios on different parts of the ecosystem (see, for example, Klein et al., 2018).

How might this work in practice?

Qualitative models are a useful means for developing a suitable, plausible network of interactions expected in an ecosystem model, linking physical, chemical and biological components. Once formed, a qualitative model can then be used to generate possible directions of change in different species/functional groups arising from press perturbations in different parts of the network, particularly in the physical and chemical components. For example, possible changes in the krill-based food web have been explored for the West Antarctic Peninsula (Melbourne-Thomas et al., 2013; Trebilco et al., in review-a, in review-b). Overall, this process can be used to simplify food web models in order to achieve computational efficiencies, in preparation for using the nested ensemble of models.

Earth System models can provide the state of habitat variables and primary producers across the Southern Ocean, although sea ice may not be well described at present (Cavanagh et al., 2017). The ability for these models to represent the actual state of the Southern Ocean can be assessed as to their fit to time series of ocean observations; the relative ability for representing reality is termed 'model skill'. Environmental scenarios from models with high skill will establish the base conditions for driving the regional food web and/or species models. The results for the different scenarios can be immediately used for looking at potential shifts in suitable habitats for different species under the different scenarios (e.g. krill eggs - Kawaguchi et al., 2013; krill larvae in sea ice - Melbourne-Thomas et al., 2016; krill growth potential - Hill et al., 2013).

Time series of observations of physics, chemistry and biology can be used to establish the starting conditions for model assessments of projected changes under different scenarios. While end-to-end models take account of the interactions between physics, chemistry and biology when undertaking projections, single species and food web assessments can be undertaken using a hierarchical approach to the models. For example, biogeochemical models can help bound the production in a region based on time-series (observations or model data) of the physical environment. The time-series of production can then be used as inputs to species-specific models or to underpin the productivity in a food web. Models such as Ecopath with Ecosim, can help ensure the starting conditions of the relative biomasses of species or functional groups are appropriate given the observed relative abundances amongst a subset of taxa. Thus, projections into the future will have plausibility given these initial calculations. For some regions and species, time-series of observations will enable species and food web models to be fit to the data, enabling a test of the plausibility of the models given the precision in the estimation of parameters.

Given the development of models to date, it will be possible to at least examine biological scenarios under different future environmental scenarios from Earth System models for Antarctic krill and krill-based food webs (e.g. some recent models available are Constable and Kawaguchi, 2017; Murphy et al., 2017).

Uncertainties in the outcomes of the projections arise from parametric uncertainty, natural variation and the role of extreme events in altering trajectories of different taxa. In addition, uncertainties can arise from different views of how the ecosystems work - structural uncertainty. Estimating the uncertainty in the consequences of the different scenarios will be an important part of the assessment (Constable, 2004; Fulton, 2010; Link et al., 2012). A further step in reducing uncertainties using this hierarchical, ensemble of models, combined with existing time-series of data, will be to evaluate how an observing system for Southern




Fig. 5. Single and integrated model approaches that could be used in marine ecosystem assessments as described in Table 5. The generic Southern Ocean food web shown here represents different energy pathways and the most commonly studied species and interactions in model analyses.

Ocean ecosystems might be improved to better contribute in the future to a subsequent MEASO (Constable et al., 2016).

5. Summary and future directions

Southern Ocean ecosystems cover a range of different physical and chemical environments with four different meridional sectors (ocean scale) and subantarctic and polar zonal divisions within sectors. The best studied sector is the East Pacific (West Antarctic Peninsula) followed by the Atlantic sector, both of which have had emphases on the Antarctic krill-based pelagic systems. Nevertheless, a nested ensemble of models with sufficient time series of observations are available to undertake circumpolar assessments of, at least, the Antarctic krill-based system. This can be achieved by applying available knowledge and general principles of interactions between physical, chemical and biological components of food webs.

We provide here a framework for auditing available data, syntheses and system models (incorporating knowledge of autecological and ecosystem processes) for a MEASO. This framework provides a means of easily collating works and information not yet included in our audit in order to make them available for future assessments. While we have had an emphasis on the scientific literature, it will be possible to use the auditing process in future to collate and make available data and models not yet or not able to be established in the literature.

An important task for MEASO will be to evaluate the degree to which future assessments may benefit from programmes to fill in taxonomic gaps in data within each of the main sectors. Here, it will be important to consider how advice to end users, such as different management bodies, may be improved by filling in those gaps. As described in the use of the system models, it will be possible to evaluate how the ecosystem parts of the SOOS could be improved by increasing spatial and/or temporal coverage of observations of particular taxa (Meredith et al., 2013; Constable et al., 2016) and important components of their habitats (Trebilco et al., in review-b). A major gap that can be identified by our audit here is the need to have greater coverage of observations and modelling of benthic systems, particularly as they may pertain to

managing the interactions of fisheries with benthic habitats as well as the role of benthic habitats in the carbon cycle (e.g. Barnes et al., 2018).

Technological advances have greatly increased our efficiency to obtain ecological data in the Southern Ocean. These advances include the use of genetics to identify species, study diversity, population connectivity and phylogeography (e.g. Grant et al., 2011; Clucas et al., 2014), stable isotopes analysing diet for foodweb studies (Raymond et al., 2011); acoustics, automated cameras and satellites in locating species and monitoring populations and habitats (e.g. Fretwell et al., 2012; Southwell et al., 2013; Trebilco et al., in review-b) and autonomous and remotely operated vehicles to survey the most remote and ice-covered regions (Gutt et al., 2017). Our temporal coverage has also increased with a number of moored and remote observing systems, providing continuous and sustained data collection. The development of a network of long-term biological monitoring stations and survey transects within ASO ecosystems has been suggested and may be feasible with these technological advances (Griffiths, 2010; Constable et al., 2014a, 2014b, 2016). Importantly, the development of improved observing in the region is coordinated by the SOOS (www.soos.aq) (Newman et al., accepted).

In addition to advancing technologies, capacity building and knowledge sharing has been highlighted as a strategic goal in preserving Antarctica's biodiversity (Chown et al., 2017). International research committees and networks such as SCAR and SOOS help to coordinate activities between their member countries with Antarctic programmes at different stages of development (Summerhayes, 2008; Newman et al., accepted). In its lifetime SCAR has expanded from 12 to 44 member countries including 14 initial stage programmes and 12 associate members. Colombia, an associate member of SCAR, is an example of a developing nation, which in the last 40 years has progressed from sending their first scientist to Antarctica on an international programme to the development of 37 research projects and leading their third international science expedition in 2016–2017 (Diaz, 2017). From the start MEASO has reflected SCAR's capacity building ethos, encouraging collaboration between Antarctic nations and involvement of early career scientists as demonstrated at the MEASO conference in 2018 with 23 participating

countries and 57 early career scientists out of the 173 attendees.

Biological assessments in the ASO began with the BIOMASS programme of the Scientific Committee on Antarctic Research in the 1980s, followed by a series of assessments and the CAML project in the International Polar Year leading to the SCAR Biogeographic Atlas of the Southern Ocean and the report on the Antarctic Climate Change and Environment. The Marine Ecosystem Assessment for the Southern Ocean is a further step in these assessments aiming to provide needed scientific advice to support the sustainable management and conservation of the region long in to the future.

Acknowledgements

We thank the MEASO community for the contributions to and expressions of interest in the MEASO project during the 2018 MEASO meeting in Hobart, Tasmania (www.measo2018.aq).

We would like to acknowledge the following international researchers who provided information regarding taxa-specific abundance surveys undertaken in Antarctica and the Southern Ocean by their national research agencies (the responses were used in the production of Fig. 4) - by country: Irene Schloss (Argentina); Philippe Ziegler, John Kitchener, Karen Westwood, John van de Hoff, Colin Southwell, Nicole Hill, Andrea Walters (Australia); Evgeny Pakhomov (Canada); Cesar Cardenas, Humberto Gonzalez (Chile); Yan Ropert-Coudert (France); Julian Gutt, Santiago Pineda Metz, Bettina Meyer, Angelika Brandt, Helen Herr (Germany); Parli Bhaskar (India); Lillo Guglielmo, Iole Leonori, Marino Vacchi, Silvia Olmastroni (India); Tsuneo Odate (Japan); Matt Pinkerton (New Zealand); Azwianewi Makhado (South Africa); Martin Edwards, Phil Trathan, Sophie Fielding, Angus Atkinson (United Kingdom); Eileen Hofman, Christian Reiss, Oscar Schofield, Jefferson Hinke (United States of America).

Additional thanks are due to lead and co-authors of the SCAR Biogeographic Atlas of the Southern Ocean who provide information regarding recent research in the field of research since the Atlas publication. With expertise: Séverine Alvain (Phytoplankton); Charles Amsler (Macroalgae); David Barnes (Bryozoa); Narissa Bax (Stylasteridae); Simone Brandão (Benthic Ostracoda); Alistar Crame (Antarctic Evolution); Bruno Danis (Asteroidea); Dhugal Lindsay (gelatinous zooplankton); John Dolan (Tintinnid ciliates); Rachel Downey (Porifera); Guy Duhamel (Fish); Kai H George (harpacticoid copepods); Andrew Gooday (Foraminifera); Huw Griffiths (Crabs and lobsters); Charlene Guillaumot (Asteroidea); Charlotte Havermans (Lysianassoidea amphipods); Graham Hosie (zooplankton); Falk Huettman (pelagic communities); Stefanie Kaiser (Isopoda); Juliana H. M. Kouwenberg (pelagic copepods); Sophie Mormede (distribution modelling); Camille Moreau (Asteroidea); Ute Mühlenthal-Siegel (Cumacea); Alexandra Post (environmental setting); Ben Raymond (pelagic regionalisation); Estafanía Rodríguez (Antarctic Hexacorals); Yan Ropert-Coudert (birds and marine mammals); Thomas Saucède (echinoids); Stefano Schiaparelli (Gastropoda); Keri Swadling (sea-ice metazoans); José Xavier (squid) and Wolfgang Zeidler (Hyperidea amphipods).

We are grateful to Australian Coastal and Oceans Modelling and Observations (ACOMO) for the opportunity to present our MEASO research at the 2018 ACOMO workshop and contribute to their special issue. Finally we would like to thank the two reviewers and editorial team that provide very useful comments during the review stages.

Funding

Funding for the MEASO conference and the development of materials for this paper were provided by: Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia; Pew Charitable Trusts, Washington D.C., USA; World Wildlife Fund, Sydney, Australia, Austral Fisheries, Perth, Australia; Australian Longline, Hobart, Australia; National Institute of Polar Research, Tokyo, Japan; Coalition of Legal

Toothfish Operators, Perth, Australia; Tasmanian Polar Network, Hobart, Australia; EU Lifewatch.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jmarsys.2019.05.008>.

References

- Agnew, D.J., 1997. The CCAMLR ecosystem monitoring programme. *Antarct. Sci.* 9, 235–242. <https://doi.org/10.1017/S095410209700031X>.
- ATCM, 2015. Report of the Thirty-Eighth Antarctic Treaty Consultative Meeting, Volume II. Secretariat of the Antarctic Treaty, Buenos Aires, Argentina.
- Atkinson, A., Ward, P., Hunt, B.P.V., Pakhomov, E.A., Hosie, G.W., 2012a. An overview of Southern Ocean zooplankton data: abundance, biomass, feeding and functional relationships. *CCAMLR Sci.* 19, 171–218.
- Atkinson, A., Nicol, S., Kawaguchi, S., Pakhomov, E., Quetin, L., Ross, R., Hill, S., Reiss, C., Siegel, V., Tarling, G., 2012b. Fitting *Euphausia superba* into Southern Ocean food-web models: a review of data sources and their limitations. *CCAMLR Sci.* 19, 219–245.
- Ballerini, T., Hofmann, E.E., Ainley, D.G., Daly, K., Marrari, M., Ribic, C.A., Smith Jr., W.O., Steele, J.H., 2014. Productivity and linkages of the food web of the southern region of the western Antarctic Peninsula continental shelf. *Prog. Oceanogr.* 122, 10–29. <https://doi.org/10.1016/j.pocean.2013.11.007>.
- Barnes, David K.A., Fleming, A., Sands Chester, J., Quartino Maria, L., Deregibus, D., 2018. Icebergs, sea ice, blue carbon and Antarctic climate feedbacks. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376, 20170176. <https://doi.org/10.1098/rsta.2017.0176>.
- Brandt, A., Ebbe, B., 2007. ANDEEP III ANtartic benthic DEEP-sea biodiversity: colonisation history and recent community patterns. *Deep-Sea Res. Part II. Top. Stud. Oceanogr.* 54, 1645–1904. <https://doi.org/10.1016/j.dsr2.2007.07.001>.
- Brandt, A., Gooday, A.J., Brandao, S.N., Brix, S., Brökeland, W., Cedhagen, T., Choudhury, M., Cornelius, N., Danis, B., De Mesel, I., Diaz, R.J., 2007a. First insights into the biodiversity and biogeography of the Southern Ocean deep sea. *Nature* 447, 307.
- Brandt, A., Griffiths, H., Gutt, J., Linse, K., Schiaparelli, S., Ballerini, T., Danis, B., Pfannkuche, O., 2014. Challenges of deep-sea biodiversity assessments in the Southern Ocean. *Adv. Polar. Res.* 25, 204–212. <https://doi.org/10.13679/j.advps.2014.3.00204>.
- Butterworth, D.S., 1988. Some aspects of the relation between Antarctic krill abundance and CPUE measures in the Japanese krill fishery. In: *Selected Scientific Papers, 1988. SC-CAMLR-SSP/5*, pp. 109–125.
- Cavalieri, D.J., Parkinson, C.L., 2008. Antarctic sea ice variability and trends, 1979–2006. *J. Geophys. Res. Oceans.* 113, C07004. <https://doi.org/10.1029/2007JC004558>.
- Cavanagh, R.D., Broszeit, S., Pilling, G.M., Grant, S.M., Murphy, E.J., Austen, M.C., 2016. Valuing biodiversity and ecosystem services: a useful way to manage and conserve marine resources? *Proc. R. Soc. Lon. B.* 283, 20161635. <https://doi.org/10.1098/rspb.2016.1635>.
- Cavanagh, R.D., Murphy, E.J., Bracegirdle, T.J., Turner, J., Knowland, C.A., Corney, S.P., Smith Jr., W.O., Waluda, C.M., Johnston, N.M., Bellerby, R.G., Constable, A.J., 2017. A synergistic approach for evaluating climate model output for ecological applications. *Front. Mar. Sci.* 4, 308. <https://doi.org/10.3389/fmars.2017.00308>.
- CCAMLR, 2015. Report of the Thirty-Fourth Meeting of the Commission for the Conservation of Antarctic Marine Living Resources. CCAMLR, Hobart, Australia.
- Chown, S.L., Clarke, A., Fraser, C.I., Cary, S.C., Moon, K.L., McGeoch, M.A., 2015. The changing form of Antarctic biodiversity. *Nature* 522, 431–438. <https://doi.org/10.1038/nature14505>.
- Chown, S.L., Brooks, C.M., Terauds, A., Le Bohec, C., van Klaveren-Impagliazzo, C., Whittington, J.D., Butchart, S.H., Coetzee, B.W., Collen, B., Convey, P., Gaston, K.J., 2017. Antarctica and the strategic plan for biodiversity. *PLoS Biol.* 15, e2001656. <https://doi.org/10.1371/journal.pbio.2001656>.
- Christensen, V., Walters, C.J., 2004. Ecopath with Ecosim: methods, capabilities and limitations. *Ecol. Model.* 172, 109–139. <https://doi.org/10.1016/j.ecolmodel.2003.09.003>.
- Clarke, A., 2008. Antarctic marine benthic diversity: patterns and processes. *J. Exp. Mar. Biol. Ecol.* 366, 48–55. <https://doi.org/10.1016/j.jembe.2008.07.008>.
- Clucas, G.V., Dunn, M.J., Dyke, G., Emslie, S.D., Levy, H., Naveen, R., Polito, M.J., Pybus, O.G., Rogers, A.D., Hart, T., 2014. A reversal of fortunes: climate change 'winners' and 'losers' in Antarctic Peninsula penguins. *Sci. Rep.* 4, 5024. <https://doi.org/10.1038/srep05024>.
- Conlan, K.E., Kim, S.L., Lenihan, H.S., Oliver, J.S., 2004. Benthic changes during 10 years of organic enrichment by McMurdo Station, Antarctica. *Mar. Poll. Bull.* 49, 43–60. <https://doi.org/10.1016/j.marpolbul.2004.01.007>.
- Constable, A.J., 2004. Managing fisheries effects on marine food webs in Antarctica: trade-offs among harvest strategies, monitoring, and assessment in achieving conservation objectives. *Bull. Mar. Sci.* 74, 583–605.
- Constable, A.J., 2005. Implementing plausible ecosystem models for the Southern Ocean: an ecosystem, productivity, ocean, climate (EPOC) model. In: *Workshop Document Presented to WG-EMM Subgroup of CCAMLR, WG-EMM-05/33*.
- Constable, A.J., 2006. Using the EPOC modelling framework to assess management procedures for Antarctic krill in Statistical Area 48: evaluating spatial differences in productivity of Antarctic krill. In: *Workshop Document Presented to WG-EMM*

- Subgroup of CCAMLR, WG-EMM-06/38.
- Constable, A.J., 2008. Implementation of FOOSA (KPFM) in the EPOC modelling framework to facilitate validation and possible extension of models used in evaluating krill fishery harvest strategies that will minimise risk of localised impacts on krill predators. In: Workshop Document Presented to WG-SAM Subgroup of CCAMLR, WG-SAM-08/15.
- Constable, A.J., 2011. Lessons from CCAMLR on the implementation of the ecosystem approach to managing fisheries. *Fish Fish.* <https://doi.org/10.1111/j.1467-2979.2011.00410.x>.
- Constable, A.J., Kawaguchi, S., 2017. Modelling growth and reproduction of Antarctic krill, *Euphausia superba*, based on temperature, food and resource allocation amongst life history functions. *ICES J. Mar. Sci.* 75, 738–750. <https://doi.org/10.1093/icesjms/fsx190>.
- Constable, A.J., Melbourne-Thomas, J., Corney, S.P., Arrigo, K.R., Barbraud, C., Barnes, D.K., Bindoff, N.L., Boyd, P.W., Brandt, A., Costa, D.P., Davidson, A.T., 2014a. Climate change and Southern Ocean ecosystems I: how changes in physical habitats directly affect marine biota. *Glob. Chang. Biol.* 20, 3004–3025. <https://doi.org/10.1111/gcb.12623>.
- Constable, A.J., Costa, D., Murphy, E., Hofmann, E., Schofield, O., Press, A., Johnston, N.M., Newman, L., 2014b. Chapter 9.3. Assessing status and change in Southern Ocean ecosystems. In: De Broyer, C., Koubbi, P., Griffiths, H.J., Raymond, B., Udekem d'Acoz, C. d' (Eds.), *Biogeographic Atlas of the Southern Ocean*. Scientific Committee on Antarctic Research, Cambridge, pp. 404–407.
- Constable, A.J., Costa, D.P., Schofield, O., Newman, L., Urban Jr., E.R., Fulton, E.A., Melbourne-Thomas, J., Ballerini, T., Boyd, P.W., Brandt, A., Willaim, K., 2016. Developing priority variables (“ecosystem Essential Ocean Variables”—eEOVs) for observing dynamics and change in Southern Ocean ecosystems. *J. Marine Syst.* 161, 26–41. <https://doi.org/10.1016/j.jmarsys.2016.05.003>.
- Constable, A.J., Melbourne-Thomas, J., Trebilco, R., Press, A.J., Haward, M., 2017. ACE CRC position analysis: managing change in Southern Ocean ecosystems. Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia Available online at: http://accrcr.org.au/publication_categories/position-analyses/.
- Convey, P., Bindschadler, R., Di Prisco, G., Fahrbach, E., Gutt, J., Hodgson, D.A., Mayewski, P.A., Summerhayes, C.P., Turner, J., ACCE Consortium, 2009. Antarctic climate change and the environment. *Ant. Sci.* 21, 541–563. <https://doi.org/10.1017/S0954102009990642>.
- Corney, S.P., Gwyther, D., Melbourne-Thomas, J., Galton-Fenzi, B.K., Mori, M., Bestley, S., Constable, A.J., 2019. Building the physics into end-to-end models: development and assessment of a circumpolar ROMS model for use as forcing of ocean ecosystem models. *JGR-Oceans* (in review).
- Cox, M.J., Candy, S., de la Mare, W.K., Nicol, S., Kawaguchi, S., Gales, N., 2018. No evidence for a decline in the density of Antarctic krill *Euphausia superba* Dana, 1850, in the Southwest Atlantic sector between 1976 and 2016. *J. Crustacean Biol.* 38, 656–661. <https://doi.org/10.1093/jcblol/ruy072>.
- Croxall, J.P., 1992. Southern Ocean environmental changes: effects on seabird, seal and whale populations. *Phil. Trans. R. Soc. B.* 388, 319–328. <https://doi.org/10.1098/rstb.1992.0152>.
- De Broyer, C., Koubii, P., 2015. Biogeographic Atlas of the Southern Ocean. *Ant. Sci.* <https://doi.org/10.1017/S0954102015000140>.
- De Broyer, C., Koubbi, P., Griffiths, H.J., Raymond, B., Udekem d'Acoz, C. d', Van de Putte, A.P., Danis, B., David, B., Grant, S., Gutt, J., Held, C., Hosie, G., Huettmann, F., Post, A., Ropert-Coudert, Y., 2014. *Biogeographic Atlas of the Southern Ocean*. Scientific Committee on Antarctic Research, Cambridge.
- Diaz, C., 2017. Colombian Antarctic Program Brochure 2017. www.scar.org/members/national-reports/colombia/3626-colombian-antarctic-programme-cap-brochure-2017/file/, Accessed date: 26 April 2019.
- Dinniman, M.S., Klinck, J.M., Smith Jr., W.O., 2011. A model study of Circumpolar Deep Water on the West Antarctic Peninsula and Ross Sea continental shelves. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 58, 1508–1523. <https://doi.org/10.1016/j.dsr2.2010.11.013>.
- Downey, R.V., Griffiths, H.J., Linse, K., Janussen, D., 2012. Diversity and distribution patterns in high southern latitude sponges. *PLoS One* 7, e41672. <https://doi.org/10.1371/journal.pone.0041672>.
- El-Sayed, S.Z., 1994. *Southern Ocean Ecology: The BIOMASS Perspective*. Cambridge University Press, Cambridge.
- Fedulov, P.P., Murphy, E.J., Shulgovsky, K.E., 1996. Environment-krill relations in the South Georgia marine ecosystem. *CCAMLR Sci.* 3, 13–30.
- Fielding, S., Watkins, J.L., Trathan, P.N., Enderlein, P., Waluda, C.M., Stowasser, G., Tarling, G.A., Murphy, E.J., 2014. Interannual variability in Antarctic krill (*Euphausia superba*) density at South Georgia, Southern Ocean: 1997–2013. *ICES J. Mar. Sci.* 71, 2578–2588. <https://doi.org/10.1093/icesjms/fsu104>.
- Fretwell, P.T., LaRue, M.A., Morin, P., Kooymann, G.L., Wienecke, B., Ratcliffe, N., Fox, A.J., Fleming, A.H., Porter, C., Trathan, P.N., 2012. An emperor penguin population estimate: the first global, synoptic survey of a species from space. *PLoS One* 7, e33751. <https://doi.org/10.1371/journal.pone.0033751>.
- Fulton, E.A., 2010. Approaches to end-to-end ecosystem models. *J. Marine Syst.* 81, 171–183. <https://doi.org/10.1016/j.jmarsys.2009.12.012>.
- Fulton, E.A., Link, J.S., Kaplan, I.C., Savina-Rolland, M., Johnson, P., Ainsworth, C., Horne, P., Gorton, R., Gamble, R.J., Smith, A.D., Smith, D.C., 2011. Lessons in modelling and management of marine ecosystems: the Atlantis experience. *Fish Fish.* 12, 171–188. <https://doi.org/10.1111/j.1467-2979.2011.00412.x>.
- Goedegebuure, M., Melbourne-Thomas, J., Corney, S.P., Hindell, M.A., Constable, A.J., 2017. Beyond big fish: the case for more detailed representations of top predators in marine ecosystem models. *Ecol. Model.* 359, 182–192. <https://doi.org/10.1016/j.ecolmodel.2017.04.004>.
- Grant, R.A., Griffiths, H.J., Steinke, D., Wadley, V., Linse, K., 2011. Antarctic DNA barcoding: a drop in the ocean? *Polar Biol.* 34, 775–780. <https://doi.org/10.1007/s00300-010-0932-7>.
- Griffiths, H.J., 2010. Antarctic marine biodiversity—what do we know about the distribution of life in the Southern Ocean? *PLoS One* 5, e11683. <https://doi.org/10.1371/journal.pone.0011683>.
- Griffiths, H.J., Danis, B., Clarke, A., 2011. Quantifying Antarctic marine biodiversity: the SCAR-MarBIN data portal. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 58, 18–29. <https://doi.org/10.1016/j.dsr2.2010.10.008>.
- Griffiths, H.J., Van de Putte, A., Danis, B., 2014. Chapter 2.2. Data analysis: patterns and implications. In: De Broyer, C., Koubbi, P., Griffiths, H.J., Raymond, B., Udekem d'Acoz, C. d' (Eds.), *Biogeographic Atlas of the Southern Ocean*. Scientific Committee on Antarctic Research, Cambridge, pp. 16–26.
- Griffiths, H.J., Meijers, A.J., Bracegirdle, T.J., 2017. More losers than winners in a century of future Southern Ocean seafloor warming. *Nat. Clim. Chang.* 7, 749–754.
- Gurney, L.J., Pakhomov, E.A., Christensen, V., 2014. An ecosystem model of the Prince Edward Island archipelago. *Ecol. Model.* 294, 117–136. <https://doi.org/10.1016/j.ecolmodel.2014.09.008>.
- Gutt, J., Starman, A., 2003. Patchiness of the megabenthos at small scales: ecological conclusions by examples from polar shelves. *Polar Biol.* 26, 276–278.
- Gutt, J., Barnes, D.K., Lockhart, S.J., Van de Putte, A., 2013. Antarctic macrobenthic communities: A compilation of circumpolar information. *Nature Conservation* 4, 1–13. <https://doi.org/10.3897/natureconservation.4.4499>.
- Gutt, J., Isla, E., Bertler, A.N., Bodeker, G.E., Bracegirdle, T.J., Cavanagh, R.D., Comiso, J.C., Convey, P., Cummings, V., De Conto, R., De Master, D., 2017. Cross-disciplinarity in the advance of Antarctic ecosystem research. *Mar. Genomics.* <https://doi.org/10.1016/j.margen.2017.09.006>.
- Hewitt, R.P., Watkins, J., Naganobu, M., Sushin, V., Brierley, A.S., Demer, D., Kasatkina, S., Takao, Y., Goss, C., Malysenko, A., Brandon, M., 2004. Biomass of Antarctic krill in the Scotia Sea in January/February 2000 and its use in revising an estimate of precautionary yield. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 51, 1215–1236. <https://doi.org/10.1016/j.dsr2.2004.06.011>.
- Hill, S.L., Murphy, E.J., Reid, K., Trathan, P.N., Constable, A.J., 2006. Modelling Southern Ocean ecosystems: krill, the food-web, and the impacts of harvesting. *Biol. Rev.* 81, 581–608. <https://doi.org/10.1017/S1464793106007123>.
- Hill, S.L., Keeble, K., Atkinson, A., Murphy, E.J., 2012. A foodweb model to explore uncertainties in the South Georgia shelf pelagic ecosystem. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 59, 237–252. <https://doi.org/10.1016/j.dsr2.2011.09.001>.
- Hill, S.L., Phillips, T., Atkinson, A., 2013. Potential climate change effects on the habitat of Antarctic krill in the Weddell quadrant of the Southern Ocean. *PLoS One* 8, e72246. <https://doi.org/10.1371/journal.pone.0072246>.
- Hofmann, E.E., Klinck, J.M., Locarnini, R.A., Fach, B., Murphy, E., 1998. Krill transport in the Scotia Sea and environs. *Ant. Sci.* 10, 406–415. <https://doi.org/10.1017/S0954102098000492>.
- Hofmann, E.E., Wiebe, P.H., Costa, D.P., Torres, J.J., 2011. Understanding the linkages between Antarctic food webs and the environment: a synthesis of Southern Ocean GLOBEC studies. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 58, 1505–1740. <https://doi.org/10.1016/j.dsr2.2011.02.001>.
- IPCC, 2014. *Climate Change 2014: Synthesis Report, Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Pachauri, R.K., Meyer, L.A. (Eds.), (Geneva, Switzerland).
- Jenouvrier, S., Holland, M., Stroeve, J., Serreze, M., Barbraud, C., Weimerskirch, H., Caswell, H., 2014. Projected continent-wide declines of the emperor penguin under climate change. *Nat. Clim. Chang.* 4, 715–718. <https://doi.org/10.1038/NCLIMATE2280>.
- Johnson, C.R., Banks, S.C., Barrett, N.S., Cazassus, F., Dunstan, P.K., Edgar, G.J., Frusher, S.D., Gardner, C., Haddon, M., Helidoniotis, F., Hill, K.L., 2011. Climate change cascades: shifts in oceanography, species' ranges and subtidal marine community dynamics in eastern Tasmania. *J. Exp. Mar. Biol. Ecol.* 400, 17–32. <https://doi.org/10.1016/j.jembe.2011.02.032>.
- Johnson, R., Stratton, P.G., Wright, S.W., McMinn, A., Meiners, K.M., 2013. Three improved satellite chlorophyll algorithms for the Southern Ocean. *J. Geophys. Res. Oceans.* 118, 3694–3703. <https://doi.org/10.1002/jgrc.20270>.
- Johst, K., Gutt, J., Wissel, C., Grimm, V., 2006. Diversity and disturbances in the Antarctic megabenthos: feasible versus theoretical disturbance ranges. *Ecosystems* 9, 1145–1155. <https://doi.org/10.1007/s10021-006-0054-9>.
- Kawaguchi, S., Ishida, A., King, R., Raymond, B., Waller, N., Constable, A., Nicol, S., Wakita, M., Ishimatsu, A., 2013. Risk maps for Antarctic krill under projected Southern Ocean acidification. *Nat. Clim. Chang.* 3, 843–847. <https://doi.org/10.1038/nclimate1937>.
- Kennicutt II, M.C., Chown, S., Cassano, J., Liggett, D., Massom, R., Peck, L., Rintoul, S., Storey, J., Vaughan, D., Wilson, T., Sutherland, W., 2014. Six priorities for Antarctic science. *Nature* 512, 23–25. <https://doi.org/10.1038/512023a>.
- Klein, E.S., Hill, S.L., Hinke, J.T., Phillips, T., Watters, G.M., 2018. Impacts of rising sea temperature on krill increase risks for predators in the Scotia Sea. *PLoS One* 13, e0191011. <https://doi.org/10.1371/journal.pone.0191011>.
- Klekociuk, A., Wienecke, B., 2017. Australia State of the Environment 2016: Antarctic Environment, Independent Report to the Australian Government Minister for the Environment and Energy. Australian Government Department of the Environment and Energy, Canberra. <https://doi.org/10.4226/94/58b65b2b307c0>.
- Kock, K.-H., Reid, K., Croxall, J., Nicol, S., 2007. Fisheries in the Southern Ocean and ecosystem approach. *Phil. Trans. R. Soc. B.* 362, 2333–2349. <https://doi.org/10.1098/rstb.2006.1954>.
- Kock, K.H., Barrera-Oro, E., Belchier, M., Collins, M.A., Duhamel, G., Hanchet, S., Pshenichnov, L., Welsford, D., Williams, R., 2012. The role of fish as predators of krill (*Euphausia superba*) and other pelagic resources in the Southern Ocean. *CCAMLR Sci* 19, 115–169.

- Koubbi, P., Grant, S., Ramm, D., Vacchi, M., Ghigliotti, L., Pisano, E., 2017. Conservation and management of Antarctic silverfish *Pleuragramma antarctica* populations and habitats. In: Vacchi, M., Pisano, E., Ghigliotti, L. (Eds.), *The Antarctic Silverfish: A Keystone Species in a Changing Ecosystem*. vol. 3. Springer, Cham, pp. 287–305.
- Laws, R.M., 1977. Seals and whales of the Southern Ocean. *Phil. Trans. R. Soc. B* 279, 81–96.
- Levins, R., 1996. The strategy of model building in population biology. *Am. Sci.* 54, 421–431.
- Link, J.S., Ihde, T.F., Harvey, C.J., Gaichas, S.K., Field, J.C., Brodziak, J.K.T., Townsend, H.M., Peterman, R.M., 2012. Dealing with uncertainty in ecosystem models: the paradox of use for living marine resource management. *Prog. Oceanogr.* 102, 102–114. <https://doi.org/10.1016/j.pocan.2012.03.00>.
- Linse, K., Griffiths, H.J., Barnes, D.K., Brandt, A., Davey, N., David, B., De Grave, S., Eléaume, M., Glover, A.G., Hemery, L.G., Mah, C., 2013. The macro-and megabenthic fauna on the continental shelf of the eastern Amundsen Sea, Antarctica. *Cont. Shelf Res.* 68, 80–90. <https://doi.org/10.1016/j.csr.2013.08.012>.
- Loeb, V.J., Santora, J.A., 2015. Climate variability and spatiotemporal dynamics of five Southern Ocean krill species. *Prog. Oceanogr.* 134, 93–122. <https://doi.org/10.1016/j.pocan.2015.01.002>.
- Mangel, P., 1988. Analysis and modelling of the Soviet Southern Ocean krill fleet. In: *Selected Scientific Papers, 1988. SC-CAMLR-SSP/5*, pp. 127–235.
- Mazloff, M.R., Heimbach, P., Wunsch, C., 2010. An eddy-permitting Southern Ocean state estimate. *J. Phys. Oceanogr.* 40, 880–899. <https://doi.org/10.1175/2009JPO4236.1>.
- McCormack, S.A., Melbourne-Thomas, J., Trebilco, R., Blanchard, J.L., Raymond, B., Constable, A., 2019;al., in review. It's not all about Antarctic krill - food web structures are fundamentally different across the Southern Ocean. *Ecography* (in review).
- McCormack, S., Melbourne-Thomas, J., Trebilco, R., Constable, A.J., Blanchard, J., 2019;al., accepted. Alternative Energy Pathways in Southern Ocean Food Webs: Insights From a Balanced Model of Prydz Bay, Antarctica. (accepted).
- Melbourne-Thomas, J., Wotherspoon, S., Raymond, B., Constable, A., 2012. Comprehensive evaluation of model uncertainty in qualitative network analyses. *Ecol. Monogr.* 82, 505–519. <https://doi.org/10.1890/12-0207.1>.
- Melbourne-Thomas, J., Constable, A., Wotherspoon, S., Raymond, B., 2013. Testing paradigms of ecosystem change under climate warming in Antarctica. *PLoS One* 8, e55093. <https://doi.org/10.1371/journal.pone.0055093>.
- Melbourne-Thomas, J., Wotherspoon, S., Corney, S., Molina-Balari, E., Marini, O., Constable, A., 2015. Optimal control and system limitation in a Southern Ocean ecosystem model. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 114, 64–73. <https://doi.org/10.1016/j.dsr2.2013.02.017>.
- Melbourne-Thomas, J., Corney, S.P., Trebilco, R., Meiners, K.M., Stevens, R.P., Kawaguchi, S., Sumner, M.D., Constable, A.J., 2016. Under ice habitats for Antarctic krill larvae: could less mean more under climate warming? *Geophys. Res. Lett.* 43, 10322–10327. <https://doi.org/10.1002/2016GL070846>.
- Melbourne-Thomas, J., Constable, A.J., Fulton, E.A., Corney, S.P., Trebilco, R., Hobday, A.J., Blanchard, J.L., Boschetti, F., Bustamante, R.H., Cropp, R., Everett, J.D., 2017. Integrated modelling to support decision-making for marine social-ecological systems in Australia. *ICES J. Mar. Sci.* 74, 2298–2308. <https://doi.org/10.1093/icesjms/lsx078>.
- Melbourne-Thomas J., Mori M., Fulton E.A., Sumner M., Corney S., Beeton N., Porobic J., Trebilco R., McCormack S. and Constable A., In prep. Implementation and calibration of the Atlantis end-to-end ecosystem model for East Antarctica.
- Meredith, Michael P., Brandon, Mark A., Murphy, Eugene J., Trathan, Philip N., Thorpe, Sally E., Bone, Douglas G., Chernyshkov, Pavel P., Sushin, Viacheslav A., 2005. Variability in hydrographic conditions to the east and northwest of South Georgia, 1996–2001. *J. of Mar. Syst.* 53, 143–167. <https://doi.org/10.1016/j.jmarsys.2004.05.005>.
- Meredith, M.P., Schofield, O., Newman, L., Urban, E., Sparrow, M., 2013. The vision for a Southern Ocean Observing System. *Curr. Opin. Environ. Sustain.* 5, 06–313. <https://doi.org/10.1016/j.cosust.2013.03.002>.
- Mori, M., Butterworth, D.S., 2004. Consideration of multi-species interaction in the Antarctic: an initial model of the minke whale-blue whale-krill interaction. *Afr. J. Mar. Sci.* 26, 245–259. <https://doi.org/10.2989/18142320409504060>.
- Mori, M., Butterworth, D.S., 2005. Modelling the predator-prey interactions of krill, baleen whales and seals in the Antarctic ecosystem. In: *Workshop Document Presented to WG-EMM Subgroup of CCAMLR, WG-EMM-05/34*.
- Mori, M., Butterworth, D.S., 2006. A first step towards modelling the krill - predator dynamics of the Antarctic ecosystem. *CCAMLR Sci* 13, 217–277.
- Murphy, E.J., Thorpe, S.E., Watkins, J.L., Hewitt, R., 2004. Modelling the krill transport pathways in the Scotia Sea: spatial and environmental connections generating the seasonal distribution of krill. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 51, 1435–1456. <https://doi.org/10.1016/j.dsr2.2004.06.019>.
- Murphy, E.J., Cavanagh, R.C., Johnston, N.M., Reid, K., Hofmann, E., 2008. Integrating Circumpolar Climate and Ecosystem Dynamics: ICED. Science and Implementation Plan. IGBP: GLOBEC and IMBER.
- Murphy, E.J., Cavanagh, R.D., Hofmann, E.E., Hill, S.L., Constable, A.J., Costa, D.P., Pinkerton, M.H., Johnston, N.M., Trathan, P.N., Klinck, J.M., Wolf-Gladrow, D.A., 2012. Developing integrated models of Southern Ocean food webs: including ecological complexity, accounting for uncertainty and the importance of scale. *Prog. Oceanogr.* 102, 74–92. <https://doi.org/10.1016/j.pocan.2012.03.006>.
- Murphy, E.J., Hofmann, E.E., Watkins, J.L., Johnston, N.M., Pinones, A., Ballerini, T., Hill, S.L., Trathan, P.N., Tarling, G.A., Cavanagh, R.A., Young, E.F., 2013. Comparison of the structure and function of Southern Ocean regional ecosystems: the Antarctic Peninsula and South Georgia. *J. Mar. Syst.* 109, 22–42. <https://doi.org/10.1016/j.jmarsys.2012.03.011>.
- Murphy, E.J., Thorpe, S.E., Tarling, G.A., Watkins, J.L., Fielding, S., Underwood, P., 2017. Restricted regions of enhanced growth of Antarctic krill in the circumpolar Southern Ocean. *Nat. Sci. Rep.* 7, 6963. <https://doi.org/10.1038/s41598-017-07205-9>.
- Newman, L., Heil, P., Trebilco, R., Katsumata, K., Constable, A.J., van Wijk, E., Assman, K., Beja, J., Bricher, P., Coleman, R., Costa, D., Diggs, S., Farneti, R., Fawcett, S., Gille, S.T., Hendry, K.R., Henley, S.F., Hofmann, E., Maksym, T., Mazloff, M., Meijers, A.J., Meredith, M.P., Moreau, S., Ozsoy, B., Robertson, R., Schloss, I., Schofield, O., Shi, J., Sikes, E.L., Smith, L.J., Swart, S., Wahlin, A., Williams, G., Williams, M.J., Herraiz-Borreguero, L., Kern, S., Lieser, J., Massom, R., Melbourne-Thomas, J., Miloslavich, P., Spreen, G., 2019;al., accepted. Delivering sustained, coordinated and integrated observations of the Southern Ocean for global impact. *Front. Mar. Sci* (accepted).
- Nicol, S., Pauly, T., Bindoff, N.L., Strutton, P.G., 2000a. “BROKE” a biological/oceanographic survey off the coast of East Antarctica (80–150°E) carried out in January–March 1996. *Deep-sea Res. Part II Top. Stud. Oceanogr.* 47, 2281–2297. [https://doi.org/10.1016/S0967-0645\(00\)00026-6](https://doi.org/10.1016/S0967-0645(00)00026-6).
- Nicol, S., Constable, A.J., Pauly, T., 2000b. Estimates of circumpolar abundance of Antarctic krill based on recent acoustic density measurements. *CCAMLR Sci* 7, 87–99.
- Nicol, S., Meiners, K., Raymond, B., 2010. BROKE-West, a large ecosystem survey of the South West Indian Ocean sector of the Southern Ocean, 30°E–80°E (CCAMLR Division 58.4.2). *Deep-sea Res. Part II Top. Stud. Oceanogr.* 9, 693–700. <https://doi.org/10.1016/j.dsr2.2009.11.002>.
- Nymand-Larson, J., Anisimov, O., Constable, A.J., Hollowed, A., Maynard, N., Prestrud, P., Prowse, T., Stone, J., 2014. Chapter 28: Polar regions. In: *Field, C.B., Barros, R.B. (Eds.), Climate Change 2014: Impacts, Adaptation, and Vulnerability. Report of Working Group II. Intergovernmental Panel on Climate Change, San Francisco*, pp. 1567–1612.
- Pasquer, B., Laruelle, G., Becquevort, S., Schoemann, W., Gooose, H., Lancelot, C., 2005. Linking ocean biogeochemical cycles and ecosystem structure and function: results of the complex SWAMCO-4 model. *J. Sea Res.* 53, 93–108. <https://doi.org/10.1016/j.seares.2004.07.001>.
- Pauly, T., Nicol, S., Higginbottom, I., Hosie, G., Kitchener, J., 2000. Distribution and abundance of Antarctic krill (*Euphausia superba*) off East Antarctica (80–150°E) during the austral summer of 1995/1996. *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 47, 2465–2488. [https://doi.org/10.1016/S0967-0645\(00\)00032-1](https://doi.org/10.1016/S0967-0645(00)00032-1).
- Peters, R.H., 1991. *A Critique for Ecology*. Cambridge University Press, Cambridge, UK.
- Pinkerton, M.H., Bradford-Grieve, J.M., Hanchet, S.M., 2010. A balanced model of the food web of the Ross Sea, Antarctica. *CCAMLR Sci.* 17, 1–32.
- Plagányi, É.E., Butterworth, D.S., 2007. A spatial multi-species operating model of the Antarctic Peninsula krill fishery and its impacts on land-breeding predators. In: *Workshop Document Presented to WG-SAM Subgroup of CCAMLR, WG-SAM-07-12*.
- Post, A.L., Lavoie, C., Domack, E.W., Leventer, A., Shevenell, A., Fraser, A.D., 2017. Environmental drivers of benthic communities and habitat heterogeneity on an East Antarctic Shelf. *Ant. Sci.* 29, 17–32. <https://doi.org/10.1017/S0954102016000468>.
- Potthoff, M., Johst, K., Gutt, J., 2006a. How to survive as a pioneer species in the Antarctic benthos: minimum dispersal distance as a function of lifetime and disturbance. *Polar Biol.* 29, 543–551. <https://doi.org/10.1007/s00300-005-0086-1>.
- Potthoff, M., Johst, K., Gutt, J., Wissel, C., 2006b. Clumped dispersal and species coexistence. *Ecol. Model.* 198, 247–254. <https://doi.org/10.1016/j.ecolmodel.2006.04.003>.
- Priddle, J., Croxall, J.P., Everson, I., Heywood, R.B., Murphy, E.J., Prince, P.A., Sear, C.B., 1988. Large-scale fluctuations in distribution and abundance of krill—a discussion of possible causes. In: *Sahrhage, D. (Ed.), Antarctic Ocean and Resources Variability*. Springer, Berlin, Heidelberg, pp. 169–182.
- Priester, C.R., Melbourne-Thomas, J., Klocker, A., Corney, S., 2017. Abrupt transitions in dynamics of a NPZD model across Southern Ocean fronts. *Ecol. Model.* 359, 372–382. <https://doi.org/10.1016/j.ecolmodel.2017.05.030>.
- Ratcliffe, N., Trathan, P., 2012. A review of the diet and at-sea distribution of penguins breeding within the CAMLR Convention Area. *CCAMLR Sci* 19, 75–114.
- Raymond, B., Marshall, M., Nevitt, G., Gillies, C.L., Van Den Hoff, J., Stark, J.S., Losekoot, M., Woehler, E.J., Constable, A.J., 2011. A Southern Ocean dietary database. *Ecology* 92, 1188. <https://doi.org/10.1890/10.1907.1>.
- Rice, J., Marschoff, E., 2016. High-latitude ice and the biodiversity dependent on it. In: *Inniss, L., Simcock, A., Ajawin, A.Y., Alcalá, A.C., Bernal, P., Calumpang, H.P., Araghi, P.E., Green, S.O., Harris, P., Kamara, O.K., Kohata, K. (Eds.), The First Global Integrated Marine Assessment*. United Nations, pp. 1–10 Chapter 46.
- Rintoul, S., Sparrow, M., Meredith, M.P., Wadley, V., Speer, K., Hofmann, E., Summerhayes, C., Urban, E., Bellerby, R., 2011. The Southern Ocean Observing System: Initial Science and Implementation Strategy. SCAR-SCOR, Cambridge, UK.
- Rogers, A.D., Johnston, N.M., Murphy, E.J., Clarke, A., 2012. Antarctic Ecosystems: An Extreme Environment in a Changing World. John Wiley & Sons, Sussex.
- Saenz, B.T., Arrigo, K.R., 2014. Annual primary production in Antarctic sea ice during 2005–2006 from a sea ice state estimate. *J. Geophys. Res. Oceans.* 119, 3645–3678. <https://doi.org/10.1002/2013JC009677>.
- SC-CCAMLR, 2004. Report of the Twenty-Third Meeting of the Scientific Committee (SC-CCAMLR-XXIII), Annex 4. In: *Report of the Workshop on Plausible Ecosystem Models for Testing Approaches to Krill Management*. CCAMLR, Hobart, Australia.
- SC-CCAMLR, 2008. Report of the Twenty-Seventh Meeting of the Scientific Committee (SC-CCAMLR-XXVII) Report of the Joint CCAMLR-IWC Workshop. CCAMLR, Hobart, Australia.
- SC-CCAMLR, 2011. Report of the Thirtieth Meeting of the Scientific Committee (SC-CCAMLR-XXX), Annex 4. In: *Report of the Working Group on Ecosystem Monitoring and Management*. CCAMLR, Hobart, Australia.
- Schiaparelli, S., Danis, B., Wadley, V., Stoddart, D.M., 2013. The census of Antarctic marine life: The first available baseline for Antarctic marine biodiversity. In: *Verde, C., di Prisco, G. (Eds.), Adaptation and Evolution in Marine Environments*. vol. 2. Springer, Berlin, Heidelberg, pp. 3–19. https://doi.org/10.1007/978-3-642-27349-0_1.

- Siegel, V., 2016. Biology and ecology of Antarctic krill. In: *Advances in Polar Ecology*. vol. 1 Springer International Publishing, Switzerland.
- SOOS <http://www.soos.aq/news/current-news/170-new-southern-ocean-knowledge-and-information-wiki-soki>.
- Southwell, C., Bengtson, J., Bester, M.N., Schytte-Blix, A., Bornemann, H., Boveng, P., Cameron, M., Forcada, J., Laake, J., Nordøy, E., Plötz, J., 2012. A review of data on abundance, trends in abundance, habitat utilisation and diet for Southern Ocean ice-breeding seals. *CCAMLR Sci* 19, 1–49.
- Southwell, C., McKinlay, J., Low, M., Wilson, D., Newbery, K., Lieser, J.L., Emmerson, L., 2013. New methods and technologies for regional-scale abundance estimation of land-breeding marine animals: application to Adélie penguin populations in East Antarctica. *Polar Biol.* 36, 843–856.
- Stark, J.S., Kim, S.L., Oliver, J.S., 2014. Anthropogenic disturbance and biodiversity of marine benthic communities in Antarctica: a regional comparison. *PLoS One* 9, e98802. <https://doi.org/10.1371/journal.pone.0098802>.
- Stoddart, M., 2010. Antarctic biology in the 21st century—advances in, and beyond the international polar year 2007–2008. *Polar Sci* 4, 97–101. <https://doi.org/10.1016/j.polar.2010.04.004>.
- Strutton, P.G., Lovenduski, N.S., Mongin, M., Matear, R., 2012. Quantification of Southern Ocean phytoplankton biomass and primary productivity via satellite observations and biogeochemical models. *CCAMLR Sc.* 19, 247–265.
- Subramaniam, R., Corney, S., Melbourne-Thomas, J., Swadling, K., 2019;al., in prep. A Foodweb Model to Evaluate Climate Change Impacts and Fisheries Management for a Large Oceanic Plateau in the Southern Ocean. (in prep).
- Summerhayes, C.P., 2008. International collaboration in Antarctica: the international polar years, the international geophysical year, and the scientific committee on Antarctic research. *Polar Rec* 44, 321–334. <https://doi.org/10.1017/S0032247408007468>.
- Teschke, K., Bester, M.N., Bornemann, H., Brandt, A., Brtnik, P., De Broyer, C., Burkhardt, E., Dieckmann, G., Flores, H., Gerdes, D., Griffiths, H.J., 2014. Scientific Background Document in Support of the Development of a CCAMLR MPA in the Weddell Sea (Antarctica) SC-CCAMLR-XXXIII/BG/02. pp. 1–92.
- Thorpe, S.E., Murphy, E.J., Watkins, J.L., 2007. Circumpolar connections between Antarctic krill (*Euphausia superba* Dana) populations: investigating the roles of ocean and sea ice transport. *Deep-sea Res. Part I Oceanogr. Res. Pap.* 54, 792–810. <https://doi.org/10.1016/j.dsr.2007.01.008>.
- Trathan, P.N., Watkins, J.L., Murrar, A.W.A., Brierly, A.S., Everson, I., Goss, C., Priddle, J., Reid, K., Ward, P., 2001. The CCAMLR-2000 Krill Synoptic Survey: a description of the rationale and design. *CCAMLR Sci.* 8, 1–23.
- Trathan, P.N., Brierley, A.S., Brandon, M.A., Bone, D.G., Goss, C., Grant, S.A., Murphy, E.J., Watkins, J.L., 2003. Oceanographic variability and changes in Antarctic krill (*Euphausia superba*) abundance at South Georgia. *Fish. Oceanogr.* 12, 569–583. <https://doi.org/10.1046/j.1365-2419.2003.00268.x>.
- Trebilco, R., Melbourne-Thomas, J., Constable, A.J., 2019;al., in review-a. The policy relevance of Southern Ocean food web structure: implications of food web change for fisheries, conservation and carbon export. *Mar. Policy* (in review).
- Trebilco, R., Melbourne-Thomas, J., Sumner, M., Wotherspoon, S., Constable, A.J., 2019;al., in review-b. Assessing status and trends of open-ocean habitats: a regionally resolved approach and Southern Ocean application. *Ecol. Indic* (in review).
- Tulloch, V.J.D., Plagányi, É.E., Matear, R., Brown, C.J., Richardson, A.J., 2017. Ecosystem modelling to quantify the impact of historical whaling on Southern Hemisphere baleen whales. *Fish. Fish.* 19, 117–137. <https://doi.org/10.1111/faf.12241>.
- Turner, J., Bindschadler, R., Convey, P., Di Prisco, G., Fahrbach, E., Gutt, J., Hodgson, D., Mayewski, P., Summerhayes, C., 2009. Antarctic Climate Change and the Environment. Scientific Committee on Antarctic Research, Cambridge.
- Turner, J., Barrand, N.E., Bracegirdle, T.J., Convey, P., Hodgson, D.A., Jarvis, M., Jenkins, A., Marshall, G., Meredith, M.P., Roscoe, H., Shanklin, J., 2014. Antarctic climate change and the environment: an update. *Polar Rec* 50, 237–259. <https://doi.org/10.1017/S0032247413000296>.
- Vancoppenolle M., Goosse H., de Montety A., Fichefet, T., Tremblay, B., Tison, J.-L., 2010. Modeling brine and nutrient dynamics in Antarctic sea ice: the case of dissolved silica. *J. Geophys. Res.* 115, C02005. doi:<https://doi.org/10.1029/2009JC005369>.
- Watters, G.M., Hill, S.L., Hinke, J.T., Matthews, J., Reid, K., 2013. Decision-making for ecosystem-based management: evaluating options for a krill fishery with an ecosystem dynamics model. *Ecol. Appl.* 23, 710–725. <https://doi.org/10.1890/12-1371.1>.
- Wiedenmann, J., Cresswell, K., Mangel, M., 2008. Temperature-dependent growth of Antarctic krill: predictions for a changing climate from a cohort model. *Mar. Ecol. Prog. Ser.* 358, 191–202. <https://doi.org/10.3354/meps07350>.
- Xavier, J.C., Brandt, A., Ropert-Coudert, Y., Badhe, R., Gutt, J., Havermans, C., Jones, C., Costa, E.S., Lochte, K., Schloss, I.R., Kennicutt, M.C., 2016. Future challenges in Southern Ocean ecology research. *Front. Mar. Sci.* 3, 94. <https://doi.org/10.3389/fmars.2016.00094>.