

Tatiana Takiishi,<sup>1</sup> Lei Ding,<sup>1</sup> Femke Baeke,<sup>1</sup> Isabella Spagnuolo,<sup>2</sup> Guido Sebastiani,<sup>2</sup> Jos Laureys,<sup>1</sup> Annemieke Verstuyf,<sup>1</sup> Geert Carmeliet,<sup>1</sup> Francesco Dotta,<sup>2</sup> Tom L. Van Belle,<sup>1</sup> Conny A. Gysemans,<sup>1</sup> and Chantal Mathieu<sup>1</sup>



# Dietary Supplementation With High Doses of Regular Vitamin D<sub>3</sub> Safely Reduces Diabetes Incidence in NOD Mice When Given Early and Long Term

*Diabetes* 2014;63:2026–2036 | DOI: 10.2337/db13-1559

**High doses of the active form of vitamin D<sub>3</sub>, 1,25-dihydroxyvitamin D<sub>3</sub> [1,25(OH)<sub>2</sub>D<sub>3</sub>], prevent diabetes in the NOD mouse but also elicit unwanted calcemic side effects. Because immune cells themselves can convert vitamin D<sub>3</sub> into 1,25(OH)<sub>2</sub>D<sub>3</sub> locally, we hypothesized that dietary vitamin D<sub>3</sub> can also prevent disease. Thus, we evaluated whether dietary administration of high doses of regular vitamin D<sub>3</sub> (800 IU/day) during different periods of life (pregnancy and lactation, early life [3–14 weeks of age], or lifelong [3–35 weeks of age]) safely prevents diabetes in NOD mice. We found that only lifelong treatment raised serum 25-hydroxyvitamin D<sub>3</sub> from 173 nmol/L in controls to 290 nmol/L, without inducing signs of calcemic or bone toxicity, and significantly reduced diabetes development in both male and female NOD mice. This diabetes protection by vitamin D<sub>3</sub> correlated with preserved pancreatic insulin content and improved insulinitis scores. Moreover, vitamin D<sub>3</sub> treatment decreased interferon- $\gamma$ -positive CD8<sup>+</sup> T cells and increased CD4<sup>+</sup>(CD25<sup>+</sup>)FoxP3<sup>+</sup> T cells in pancreatic draining lymph nodes. In conclusion, this study shows for the first time that high doses of regular dietary vitamin D<sub>3</sub> can safely prevent diabetes in NOD mice when administered lifelong, although caution is warranted with regards to administering equivalently high doses in humans.**

Type 1 diabetes is recognized as an autoimmune-mediated disorder with a variable prodromal phase characterized by

the progressive loss of the insulin-producing  $\beta$ -cells in the pancreatic islets in genetically at-risk individuals (1,2). Several facts support a critical role for environmental factors that trigger the development of type 1 diabetes. Leading environmental candidates include exposure to enteroviruses, early introduction of wheat, and insufficient vitamin D levels. The prevalence of type 1 diabetes increases with latitude of residence and decreased sunlight exposure, whereas exposure to ultraviolet light, known to induce vitamin D production in the skin, is associated with the lower incidence of type 1 diabetes in countries closer to the equator (3,4). These observations strengthen the hypothesis that an inadequate vitamin D status due to insufficient sun exposure, dietary uptake, and/or abnormalities in its metabolism may increase the risk of type 1 diabetes (5). Vitamin D deficiency indeed increases the onset and severity of autoimmune type 1 diabetes in at-risk children (6) and also in the NOD mouse (7,8). Moreover, most epidemiological data, based on dietary questionnaires, suggested that vitamin D supplementation during pregnancy, infancy, or early adulthood might be associated with a reduced risk of type 1 diabetes later in life (6,9,10). It has been suggested that vitamin D in early life is important for gut maturation, thereby reducing permeability for agents/proteins that can act as potent antigenic stimuli (11,12).

Despite these exciting results, several areas of controversy remain in this domain, and randomized, double-blinded,

<sup>1</sup>Clinical and Experimental Endocrinology (CEE), Department of Clinical and Experimental Medicine, Katholieke Universiteit Leuven, Campus Gasthuisberg, Leuven, Belgium

<sup>2</sup>Diabetes Unit, Department of Medicine, Surgery and Neurosciences, University of Siena and Fondazione Umberto Di Mario Organizzazione Non Lucrativa di Utilità Sociale, Siena, Italy

Corresponding author: Chantal Mathieu, chantal.mathieu@uzleuven.be.

Received 11 October 2013 and accepted 6 February 2014.

This article contains Supplementary Data online at <http://diabetes.diabetesjournals.org/lookup/suppl/doi:10.2337/db13-1559/-/DC1>.

C.A.G. and C.M. share senior authorship.

© 2014 by the American Diabetes Association. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

placebo-controlled trials with extended observation periods in humans are limited. Moreover, most of the human studies have several shortcomings, including recall bias, absence of repeated 25(OH)D measurements, and lack of a quantifiable evaluation of intake of vitamin D from food or supplements or data on sunlight exposure. So far, convincing evidence is missing on whether dietary vitamin D as D<sub>2</sub> (ergocalciferol, from plants) or D<sub>3</sub> (cholecalciferol, from animals) can reduce or prevent type 1 diabetes. Moreover, what formulation, dose, route, or duration of treatment is ideal to intervene in the diabetes process remains unclear.

In a small pilot study from our group, NOD mice treated with cholecalciferol at a daily dose of 1,000 IU (25 µg), given intraperitoneally (i.p.) from 3 until 70 days of age, were unaffected in their diabetes presentation (13), whereas the active form of vitamin D<sub>3</sub>, 1,25-dihydroxyvitamin D<sub>3</sub> [1,25(OH)<sub>2</sub>D<sub>3</sub>, 5 µg/kg given i.p. on alternative days] from weaning until 200 days of age clearly reduced the development of autoimmune diabetes, although this inhibition occurred only at doses that also caused mild hypercalcemia and bone demineralization (7,14). In another study, oral administration of 50 ng 1,25(OH)<sub>2</sub>D<sub>3</sub> per day reduced diabetes incidence compared with vitamin D-deficient NOD mice (8).

In the current study, we evaluated the effects of high-dose oral vitamin D<sub>3</sub> supplementation administered at various intervals in life—during pregnancy and neonatal life as well as during infancy (early life) and adulthood (lifelong)—in safety, immune modulation, and the development of diabetes in already vitamin D-sufficient NOD mice.

## RESEARCH DESIGN AND METHODS

### Animals

NOD mice, originally obtained from Professor Wu (Department of Endocrinology, Peking Union Medical College Hospital, Beijing, China), were housed and bred in animal facility of the Katholieke Universiteit (KU) Leuven since 1989. Housing of NOD mice occurred under semibarrier conditions, and animals were fed sterile food and water ad libitum. NOD mice were screened for the onset of diabetes by evaluating glucose levels in urine (Clinistix; Bayer Diagnostics, Tarrytown, NY) and venous blood (Accu-Chek Aviva; Roche Diagnostics Belgium, Vilvoorde, Belgium). Mice were diagnosed as diabetic when having positive glycosuria and two consecutive blood glucose measurements above 200 mg/dL. At the time of the experiments, the breeding stock had a diabetes incidence of 84% in female and 38% in male mice. Animals were maintained in accordance with the National Institutes of Health *Guide for the Care and Use of Laboratory Animals*.

### Experimental Design

NOD mice were randomly assigned to be fed a vitamin D<sub>3</sub>-sufficient control diet or a vitamin D<sub>3</sub>-supplemented

diet during three periods of life. The control diet consisted of the Ssniff R/M-H diet (BioServices BV, Uden, the Netherlands) plus 1% calcium, 0.7% phosphorus, and 1,000 IU vitamin D<sub>3</sub>/kg diet (4 IU/day based on a consumption of 4 g chow/20 g body wt [BW] daily). The vitamin D<sub>3</sub>-supplemented diet consisted of the Ssniff R/M-H diet plus 1% calcium, 0.7% phosphorus and 200,000 IU vitamin D<sub>3</sub>/kg diet (800 IU/day). The different feeding periods were early life (age 3–14 weeks), lifelong (age 3–35 weeks), and during pregnancy and neonatal life (until the end of lactation). All mice were monitored until 35 weeks of age. The research protocol of this study was approved and performed in accordance with the KU Leuven Ethics Committees (Leuven, Belgium).

### Calcium and Bone Parameters

At 35 weeks of age, blood was collected by heart puncture and a femur was removed. Serum and the femur were stored at –20°C until biochemical determinations were performed. Calcium content of the femur, measured on HCl-dissolved bone ashes, and of the serum was analyzed by SYNCHRON Clinical Systems (Beckman Coulter, Analis SA, Suarlée, Belgium) and corrected for dry weight (bone). Phosphate levels in the serum were analyzed by the same system. Levels of osteocalcin and 25(OH)D<sub>3</sub> were measured using an in-house (15) and 25(OH)D<sub>3</sub> RIA kit (DiaSorin NV, Anderlecht, Belgium), respectively.

### Microcomputed Tomography

Microcomputed tomography (µCT) analysis of the femur was performed ex vivo using the high resolution SkyScan 1172 system (settings: 50 kV, 200 µA, 0.5-mm Al filter, 5-µm pixel size; Bruker-microCT, Kontich, Belgium) as previously described (16). In brief, serial tomographs, reconstructed from raw data using the cone-beam reconstruction software (NRecon v.1.4.4.0; Bruker-microCT), were used to calculate trabecular and cortical parameters, respectively, from the metaphyseal and middiaphyseal area. A bone standard was used for the calibration of bone-density measurements. Measurements were performed in the Laboratory of Clinical and Experimental Endocrinology, KU Leuven, according to the guidelines of the American Society for Bone and Mineral Research.

### Glucose Tolerance Test

Glycemia was measured in tail vein blood using a glucometer before and after (5, 15, 30, 60, and 120 min) an intraperitoneal injection of 2 g/kg BW D-glucose monohydrate dissolved in sterile PBS.

### Histology, Immunohistochemistry, and Insulin Determination

Severity of insulinitis was assessed by histological screening of pancreatic sections of at least four animals per group imbedded in paraffin taken from experimental NOD mice at 35 weeks of age. Tissue sections (6 µm) from formalin-fixed paraffin-embedded pancreata of each animal were cut and collected 100 µm apart, then stained with hematoxylin and eosin. Islets were observed under light microscopy

at original magnification  $\times 20$  or  $\times 40$ , enumerated, and graded by an independent investigator in blinded fashion. At least 25 islets per pancreatic sample were scored for islet infiltration as follows: 0, no infiltration; 1, peri-insulinitis; 2, lymphocyte infiltration in less than 50% of the islet; 3, lymphocyte infiltration in more than 50% of the islet; or 4, completely destroyed islets.

Immunofluorescence detection of FoxP3-expressing cells was performed on 6- $\mu\text{m}$ -thick paraffin tissue sections. Briefly, after antigen retrieval (10 mmol/L citrate buffer [pH = 6], 10 min), sections were incubated with fluorescein isothiocyanate (FITC)-conjugated anti-FoxP3 primary antibody (5  $\mu\text{g}/\text{mL}$ , clone FJK-16a; eBioscience) in Tris-buffered saline/proteinase I with 5% BSA (overnight, 4°C), followed by AlexaFluor (AF) 488-conjugated rabbit anti-FITC antibody (12.5  $\mu\text{g}/\text{mL}$ ; Molecular Probes, Invitrogen). For signal amplification, AF 488-conjugated anti-rabbit IgG was used (8  $\mu\text{g}/\text{mL}$ ; Molecular Probes). Next, insulin was revealed using guinea pig anti-swine antibody (0.3 mg/mL, 60 min; DakoCytomation) and AF 555-conjugated goat anti-guinea pig IgG (4  $\mu\text{g}/\text{mL}$ , 60 min; Molecular Probes). The other half of each pancreas was used for insulin content determination as described (17).

#### Flow Cytometric Analysis

Single-cell suspensions of spleen and pancreatic lymph nodes (PLN) were prepared from mice at 35 weeks of age. The following antibodies were used for staining: CD4 (GK1.5), CD8 (53-6.7), CD25 (PC61.5), CD11b (M1/70), CD11c (N418), I-A<sup>k</sup> (10-3.6, cross-reactive with I-A<sup>B7</sup> of NOD mice), the programmed death 1 ligand (MIH5), interferon- $\gamma$  (IFN- $\gamma$ ), FoxP3, and matching isotype controls (all eBioscience). Cells were analyzed in a Gallios flow cytometer with Kaluza software (Beckman Coulter).

#### Statistical Analysis

Graphs were plotted and statistics calculated with GraphPad Prism software (GraphPad Software, La Jolla, CA). Differences in the incidence of diabetes were assessed using the Mantel-Cox log-rank test. Results were expressed

as mean  $\pm$  SEM if normally distributed or as median and interquartile range if not normally distributed. The Student *t* test or Mann-Whitney test were used for comparison between groups, respectively. *P* values of  $< 0.05$  were considered statistically significant (\**P*  $< 0.05$ , \*\**P*  $< 0.01$ , \*\*\**P*  $< 0.001$ ).

## RESULTS

### Safety of Vitamin D<sub>3</sub> Supplementation

On the basis of dose-titration studies (data not shown), we selected 800 IU/day (40,000 IU/kg BW/day) as a dose raising the vitamin D levels into the “highly vitamin D-sufficient” zone [25(OH)D<sub>3</sub> concentration  $> 230$  nmol/L], without inducing hypercalcemia or bone decalcification. Only in mice that were treated lifelong (from 3 until 35 weeks of age), serum 25(OH)D<sub>3</sub> concentration increased to a mean concentration of 290 nmol/L (236–351 nmol/L) in male and female NOD mice at 35 weeks of age compared with 173 nmol/L (125–224 nmol/L) in the group fed the control diet (Table 1). This protocol did not perturb normal weight evolutions and did not cause significant alterations in serum phosphate or calcium levels (Table 1). However, the 25(OH)D<sub>3</sub> concentrations in mice supplemented with vitamin D<sub>3</sub> during pregnancy and lactation or during early life (from 3 until 14 weeks of age) were comparable to mice fed the control diet until 35 weeks of age (Supplementary Fig. 1).

Histological analysis of kidney and heart of the vitamin D<sub>3</sub> group revealed no calcifications in aorta and renal arteries (data not shown). To verify whether vitamin D<sub>3</sub> supplementation affected bone metabolism, serum osteocalcin and bone parameters were evaluated. Serum osteocalcin concentrations were not altered by lifelong vitamin D<sub>3</sub> supplementation (Table 1), indicating no alterations in bone turnover. Moreover, bone dry weight and the ratio of calcium-to-bone dry weight was similar to controls in the vitamin D<sub>3</sub> group at the end of the observation period (Table 1). The bone microstructure analysis by  $\mu\text{CT}$  showed that dietary vitamin D<sub>3</sub> did not affect bone-mineral density and did not impair cortical or trabecular bone architecture

**Table 1—Vitamin D and calcium metabolism after vitamin D<sub>3</sub> supplementation in NOD mice**

| Parameters                          | 35 weeks of age |                     |                 |                     |
|-------------------------------------|-----------------|---------------------|-----------------|---------------------|
|                                     | Females         |                     | Males           |                     |
|                                     | Controls        | 800 IU              | Controls        | 800 IU              |
| Weight (g)                          | 26.5 $\pm$ 0.5  | 26.0 $\pm$ 0.7      | 32.2 $\pm$ 0.4  | 33.6 $\pm$ 3.2      |
| Serum 25(OH)D <sub>3</sub> (nmol/L) | 184.5 $\pm$ 6.1 | 301.3 $\pm$ 20.9*** | 160.3 $\pm$ 5.1 | 280.0 $\pm$ 10.9*** |
| Serum calcium (mmol/L)              | 2.4 $\pm$ 0.1   | 2.6 $\pm$ 0.1       | 2.6 $\pm$ 0.2   | 2.6 $\pm$ 0.1       |
| Serum phosphate (mmol/L)            | 3.2 $\pm$ 0.3   | 3.7 $\pm$ 0.3       | 3.3 $\pm$ 0.2   | 3.0 $\pm$ 0.2       |
| Serum osteocalcin (ng/mL)           | 52.2 $\pm$ 5.8  | 49.5 $\pm$ 5.8      | 35.4 $\pm$ 4.1  | 39.3 $\pm$ 3.2      |
| Bone dry weight (mg)                | 51.1 $\pm$ 0.9  | 52.7 $\pm$ 0.7      | 51.5 $\pm$ 5.4  | 56.3 $\pm$ 0.9      |
| Calcium-to-bone dry weight          | 0.30 $\pm$ 0.21 | 0.36 $\pm$ 0.21     | 0.33 $\pm$ 0.01 | 0.31 $\pm$ 0.02     |

Summary data (mean  $\pm$  SEM) from female and male (*n* = 4–5 per experiment). NOD mice were fed normal chow (4 IU vitamin D, control) or chow supplemented with 800 IU/day vitamin D<sub>3</sub>. Statistical significance was calculated by the Student *t* test. \*\*\**P*  $< 0.001$  vs. controls.

(Fig. 1A and B). The vitamin D<sub>3</sub>-supplemented diet did not affect any of the analyzed bone parameters (Fig. 1B). Overall, long-term vitamin D<sub>3</sub> supplementation was tolerated well, and we found no indications of vitamin D toxicity or adverse metabolic side effects.

### Vitamin D<sub>3</sub> Supplementation Reduces Diabetes Incidence in NOD Mice When Given Early and Lifelong


We next assessed the effect of vitamin D<sub>3</sub> on diabetes development in the NOD mouse. Vitamin D<sub>3</sub> was given prophylactically (800 IU/day) during pregnancy and lactation (until 3 weeks of age), during early life (from 3 until 14 weeks of age), or lifelong from early life onwards (from 3 until 35 weeks of age). Only lifelong vitamin D<sub>3</sub> supplementation could significantly inhibit diabetes development in female and male NOD mice, resulting in a 52% reduction in diabetes incidence in females and a 58% reduction in males compared with mice fed the control diet (Fig. 2A). Only the long-term supplementation was effective in controlling diabetes progression, because shortening the period of vitamin D<sub>3</sub> treatment from 3 until 14 weeks of age caused a loss of diabetes protection by vitamin D<sub>3</sub>: in females, the diabetes incidence in vitamin D<sub>3</sub>-supplemented mice was comparable to the control group (Fig. 2B), whereas in males, vitamin D<sub>3</sub> supplementation demonstrated a trend to decreased diabetes development, yet this was not significant (Fig. 2B). When given during pregnancy and lactation, vitamin D<sub>3</sub> supplementation had no effect on diabetes development in female or male offspring (Fig. 2C).

### Lifelong Vitamin D<sub>3</sub> Supplementation Preserves $\beta$ -Cell Function and Reduces Severe Insulinitis


Prevention of diabetes by vitamin D<sub>3</sub> suggests preservation of sufficient  $\beta$ -cell mass and/or function. To test this, we first did an intraperitoneal glucose tolerance test and found that the disease-free vitamin D<sub>3</sub> group controlled a glucose bolus with similar efficiency as normoglycemic NOD mice fed the control diet (Fig. 3A). Second, the insulin content of pancreata harvested from disease-free vitamin D<sub>3</sub>-supplemented mice was comparable to amounts present in normoglycemic controls. Importantly, pancreata of disease-free vitamin D<sub>3</sub>-supplemented animals contained significantly more insulin than pancreata of diabetic NOD mice fed the control diet at 35 weeks of age (Fig. 3B). Third, scoring of the insulinitis revealed that immune infiltration in islets worsened in the control group, whereas less severe insulinitis, including a higher number of insulinitis-free islets, were observed in most of the vitamin D<sub>3</sub>-supplemented mice (Fig. 3C and D). Taken together, treatment efficiency was reflected not only in parameters measuring  $\beta$ -cell function and mass but also in the grade of insulinitis.

### Vitamin D<sub>3</sub> Supplementation Decreases Effector CD8<sup>+</sup>IFN- $\gamma$ and CD4<sup>+</sup>IFN- $\gamma$ T Cells and Increases CD4<sup>+</sup>(CD25<sup>+</sup>)FoxP3<sup>+</sup> Regulatory T Cells

We found that lifelong vitamin D<sub>3</sub> supplementation (from 3 until 35 weeks of age) did not cause immune depletion because the total number of lymphocytes in the spleen and PLN was similar to values found in sex-matched NOD



**Figure 1**—Dietary vitamin D<sub>3</sub> supplementation does not affect bone parameters. *A*:  $\mu$ CT-based visualization of trabecular bone in femur from control (*upper panels*) and vitamin D<sub>3</sub>-supplemented NOD mice (*lower panels*). *B*: From left to right: quantification of the trabecular bone volume (BV/TV), trabecular thickness (Tb.Th), trabecular number (Tb.N), and cortical thickness (Ct.Th) in control mice and long-term 800 IU/day vitamin D<sub>3</sub>-supplemented mice at 35 weeks of age. Statistical significance was calculated using Student *t* test. ns, not significant.




**Figure 2**—Lifelong vitamin D<sub>3</sub> supplementation reduces diabetes incidence in NOD mice. Female and male NOD mice were fed control chow until 35 weeks of age or fed chow supplemented with 800 IU/day vitamin D<sub>3</sub> per from 3 until 35 weeks of age (lifelong) (A), from 3 until 14 weeks of age (early life) (B), or during coupling until 3 weeks of age (pregnancy and lactation) (C). Dietary regimens are depicted graphically; in gray is the period when mice received the vitamin D<sub>3</sub>-supplemented diet. Mice with two consecutive measurements of blood glucose levels >200 mg/dL were considered diabetic. Statistical significance was calculated vs. NOD mice fed control chow by Mantel-Cox log-rank test. \**P* < 0.05; \*\**P* < 0.01.

mice fed the control diet (Fig. 4). Also, there were no significant differences in the frequencies of dendritic cell and macrophage subsets in spleen and PLN between the vitamin D<sub>3</sub> and the control group (data not shown). Likewise, expression of the antigen-presenting molecule MHC class II I-A<sup>g7</sup>

and the programmed death 1 ligand on antigen-presenting cells was unaltered in mice receiving dietary vitamin D<sub>3</sub> supplementation (data not shown).

We did not observe differences in the absolute number and proportions of CD4<sup>+</sup> and CD8<sup>+</sup> T cells in the spleen



**Figure 3**—Lifelong vitamin D<sub>3</sub> supplementation preserves insulin content and decreases severe insulinitis. **A**: Intraperitoneal glucose tolerance tests (IPGTT) were performed at the end of the observation period (at 35 weeks of age) in experimental groups, as indicated. **B**: Evaluation of insulin content in pancreatic extracts from experimental groups by ELISA, as indicated. Data are expressed as ng/mg of pancreas. Graphs show the median and interquartile range. Statistical significance between two groups was calculated using the Mann-Whitney test. ns, not significant; \*\**P* < 0.01. **C**: Pancreatic sections of indicated groups were stained with hematoxylin and eosin, and insulinitis was scored as indicated in RESEARCH DESIGN AND METHODS. From left to right graphs show the percentage of insulinitis, percentage of insulinitis-free islets, and percentage of islets with heavy insulinitis in vitamin D<sub>3</sub>-supplemented mice vs. control normoglycemic and diabetic mice. Statistical significance between two groups was calculated using the Mann-Whitney test; ns, not significant; \**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001. **D**: The following score was used for the evaluation of insulinitis: 0 (white, intact islets), I (vertical lines, peri-insulinitis), II (gray, <50% infiltration), III (horizontal lines, >50% infiltration), IV (black, complete destruction).

and PLN after vitamin D<sub>3</sub> supplementation (Fig. 4 and Supplementary Fig. 2). However, lifelong vitamin D<sub>3</sub> supplementation induced a shift in the balance between effector T cells (Teff) and regulatory T cells (Tregs). As such, vitamin D<sub>3</sub> supplementation significantly decreased the frequency as well as the total number of CD8<sup>+</sup> T cells producing IFN- $\gamma$  in the spleen and PLN of female and male NOD mice (Fig. 5A and Supplementary Fig. 3). Moreover, the proportions and the absolute numbers of CD4<sup>+</sup> T cells expressing IFN- $\gamma$  were lower in the spleen of vitamin D<sub>3</sub>-supplemented female mice and in the PLN of vitamin D<sub>3</sub>-supplemented male mice compared with sex-matched NOD mice fed the control diet (Fig. 5B and Supplementary Fig. 3).


Expression of FoxP3, a transcription factor associated with Treg differentiation and function (18), is presently the most reliable marker to identify Tregs in mice. Female and male NOD mice receiving lifelong vitamin D<sub>3</sub> supplementation showed a significantly higher frequency of CD4<sup>+</sup>FoxP3<sup>+</sup> Tregs in the PLN but not in the spleen compared with sex-matched mice fed the control diet (Fig. 6A). Analysis of CD25 expression on CD4<sup>+</sup>FoxP3<sup>+</sup> T cells

revealed that this increase concerned activated Tregs, which were increased not only in the spleen but also in the PLN of vitamin D<sub>3</sub>-treated mice (Fig. 6B) (19,20). Of note, vitamin D<sub>3</sub> supplementation during pregnancy and lactation or during early life (from 3 until 14 weeks of age) did not increase the percentages of CD4<sup>+</sup>(CD25<sup>+</sup>) FoxP3<sup>+</sup> T cells in the spleen or PLN (Supplementary Fig. 4).

Enumeration of FoxP3<sup>+</sup> cells in the pancreas revealed higher numbers of FoxP3<sup>+</sup> T cells in the islet infiltrates of vitamin D<sub>3</sub>-supplemented mice, especially in the females, compared with diabetic controls (Fig. 7), suggesting these Tregs suppress autoreactive responses not only in the PLN but also at the site of inflammation.

## DISCUSSION

Vitamin D is a prohormone that is well known for its role in calcium homeostasis and bone mineralization and metabolism, but accumulating data suggest additional roles in many important body functions, including the regulation of innate and adaptive immune responses. In humans, the daily vitamin D intake recommended by the




**Figure 4**—Unaltered lymphocyte number and subsets in spleen and PLN of mice after lifelong vitamin D<sub>3</sub> supplementation. Spleen and PLN were harvested from NOD mice ( $n = 5-7$ ) at 35 weeks of age. The absolute number of lymphocytes (A), CD4<sup>+</sup> (B), and CD8<sup>+</sup> T-cell subsets (C) were measured by flow cytometry in the spleen (left panels) and PLN (right panels) of control and vitamin D<sub>3</sub>-supplemented mice. Graphs show median and interquartile range. Statistical significance was calculated by the Mann-Whitney test.

Institute of Medicine is 400 IU/day for children younger than 1 year, up to 600 IU/day for all individuals between 1 and 70 years, assuming already some sunlight exposure, and 800 IU/day for individuals older than 70 years (21). However, many clinicians suggest that for a good health state and to allow extraskeletal benefits, the daily supply of vitamin D from all sources should be greater than 800 IU.

Until now, data are very limited on whether dietary vitamin D supplements can in fact modify the course of autoimmune diseases such as type 1 diabetes in humans. These studies often lack precise information on the dosage and regimen of vitamin D supplementation, which can lead to contradictory results. Meta-analysis of records from four case-control studies and one cohort study implies that the risk of type 1 diabetes development was significantly reduced in children treated with regular vitamin D between 7 and 12 months of age compared with those who were supplemented before 7 months of age or not supplemented (5,22). A recent meta-analysis of two cohort studies and six case-control studies on vitamin D intake during early life corroborated these findings (23). The importance of maternal vitamin D intake during

pregnancy is also not clear: a case-control study in Norway indicated that the risk of type 1 diabetes was higher in the offspring of women with the lowest 25(OH)D concentrations (10), whereas other studies did not find any correlation between maternal serum 25(OH)D values and type 1 diabetes risk in the offspring (24,25). Only properly designed, randomized, and long-term trials will bring the definitive answer.

Here, we provide the first evidence that only early and lifelong vitamin D<sub>3</sub> supplementation with 800 IU/day, which significantly increased serum 25(OH)D<sub>3</sub> levels above sufficiency and corrected the immunological defects in diabetes-prone mice that already had sufficient, even high, vitamin D<sub>3</sub> levels, safely reduced diabetes development. Vitamin D<sub>3</sub> supplementation prevented severe insulinitis and preserved  $\beta$ -cell mass best. In contrast, a short and early intervention (age 3–14 weeks) with regular vitamin D<sub>3</sub> was less protective, in line with our previous study showing that short-term intraperitoneal administration of 1,000 IU of vitamin D<sub>3</sub> to NOD mice early in life, corresponding to neonatal life and childhood in man, was insufficient to protect against diabetes (13).



**Figure 5**—Decrease of CD8 $^+$ IFN- $\gamma^+$  T cells in spleen and PLN after lifelong vitamin D $_3$  supplementation. Spleen and PLN were harvested from NOD mice ( $n = 5-7$ ) at 35 weeks of age. The frequencies of CD8 $^+$ IFN- $\gamma^+$  (A) and CD4 $^+$ IFN- $\gamma^+$  (B) T-cell subsets were measured by flow cytometry and are displayed as the percentage of IFN- $\gamma^+$  in CD8 $^+$  (A) or CD4 $^+$  (B) gate in the spleen (left panels) and PLN (right panels) of control and vitamin D $_3$ -supplemented mice. Graphs show median and interquartile range. Statistical significance was calculated by the Mann-Whitney test. \* $P < 0.05$ .

Of note, we did not evaluate the effects of vitamin D supplementation at a later stage in life (e.g., age 14–35 weeks) because we demonstrated previously that a late intervention with the bioactive vitamin D $_3$  metabolite (from 14 weeks of age, when insulinitis is present) failed to prevent disease (26).


Oral administration of vitamin D may favor regulation of tight junction proteins necessary to maintain mucosal integrity (27). Moreover, Ooi et al. (28) demonstrated that oral bioactive vitamin D $_3$  can regulate the composition of the gut bacterial microflora and that deficiency of vitamin D or its receptor can cause impaired epithelial integrity, dysbiosis, increased inflammation, and more severe experimental colitis. Active vitamin D also regulates the development and function of regulatory invariant natural killer T cells and intraepithelial CD8 $\alpha$  lymphocytes in the gut (28,29). These data imply that vitamin D supplementation might be a way to modulate the gut microflora and its microbiome and, consequently, regulates gastrointestinal innate and adaptive immune responses.

Recently, Roy et al. (30) suggested that not just the antagonistic interaction between Treg and Teff cells but

also their continuous regulation by vitamin D ultimately determines the outcome of an (auto)immune disease. This implies that optimal vitamin D concentrations are vital in keeping the balance between Treg and Teff cells in individuals (or animals) with an underlying immune dysfunction. In fact, female and male NOD mice fed a vitamin D $_3$ -sufficient control diet have normal concentrations of serum 25(OH)D $_3$  (mean, 173 nmol/L) and still present a high incidence of diabetes. Thus, our data suggest that 25(OH)D $_3$  concentrations of more than 230 nmol/L may be required to modulate immune dysregulation in NOD mice. Although further studies are needed, a recent report from the Institute of Medicine indicated that serum 25(OH)D $_3$  concentrations in humans above 125 nmol/L may result in potential adverse reactions and there should be caution against overtreatment (21).

Several studies in human subjects evaluated the influence of administering vitamin D to pregnant and/or breastfeeding women on the risk of developing type 1 diabetes in their children (31,32). Despite some early studies suggesting a protective effect of vitamin D, a meta-analysis by Zipitis and Akobeng (22) concluded that supplementing pregnant women with vitamin D






**Figure 6**—Lifelong vitamin D<sub>3</sub> supplementation increases Treg frequencies. At 35 weeks of age, spleen (*right panels*) and PLN (*left panels*) were harvested from NOD mice ( $n = 5$  to 6), and the frequency of Tregs was determined as the percentage of FoxP3<sup>+</sup> in the CD4<sup>+</sup> fraction (A) and activated Tregs as the percentage of CD25<sup>+</sup>FoxP3<sup>+</sup> in the CD4<sup>+</sup> gate (B) using flow cytometry. Graphs show median and interquartile range. Statistical significance was calculated using the Mann-Whitney test. \* $P < 0.05$ ; \*\* $P < 0.01$ .

had no implications on type 1 diabetes development in their infants. Our current observations in NOD mice are consistent with this. Moreover, our data showing that vitamin D<sub>3</sub> supplementation during lactation failed to prevent disease in NOD mice are in line with an earlier study showing that 16 IU/day vitamin D from conception until 10 weeks of age does not suffice to interfere with disease progression in NOD mice (33). It is suggested that during this early period, the adaptive immune system has not yet fully matured and thus escapes the immunomodulatory action of vitamin D. As such, supplementation with vitamin D<sub>3</sub> during pregnancy and lactation in already vitamin D<sub>3</sub>-sufficient mothers will not have immunoregulatory actions on the hyperactivity of effector immune cells against self-antigens. We also found that our vitamin D<sub>3</sub> protocol given from early life on and lifelong did not cause immune depletion because the total number of lymphocytes and T-cell subsets (e.g., CD4, CD8) in the spleen and PLN remained unchanged. However, vitamin D<sub>3</sub> supplementation dampened Th1 responses, which are often associated with autoimmune responses (34), and skewed the Teff-to-Treg balance in favor of Tregs in diabetes-prone mice. Interestingly, the increase of Tregs concerned activated Tregs according to CD25 expression (20,35). Our observations are supported by several *in vitro* and *in vivo* findings. The active form of vitamin D exerts a marked

inhibitory effect on the adaptive immune cells by inhibiting Th1 polarization (36) and triggering the emergence of Tregs with the functional capacity to suppress activation and proliferation of Teff cells (37,38). Here, we report that upon vitamin D<sub>3</sub> supplementation, these Treg (subsets) preferentially accumulate in draining PLN and pancreatic islets. Our group had already documented that a vitamin D analog had the capacity to imprint human T cells with a specific homing signature favoring migration to sites of inflammation (37).

Finally, caution is warranted when interpreting these data, because not only lifelong administration of vitamin D<sub>3</sub> was necessary for protection but also extremely high doses were needed, namely, more than 200 times the amount of vitamin D normally present in the daily amount of food of a mouse. Converting these doses to humans is difficult, and we advocate that rather the serum concentrations of 25(OH)D<sub>3</sub> reached after supplementation versus baseline in the mice should be considered. The values achieved in this study (mean, 290 nmol/L) are well above the present advocated targets of 50–100 nmol/L for humans (39), but it is important to keep in mind that control NOD mice are already very vitamin D-sufficient [serum 25(OH)D<sub>3</sub> concentrations >150 nmol/L]. If we apply the estimate of Heaney et al. (40) that a 40 IU/day increment in intake raises 25(OH)D<sub>3</sub> by 0.7 nmol/L in



**Figure 7**—Increased numbers of FoxP3<sup>+</sup> cells in the pancreatic infiltrates of lifelong vitamin D<sub>3</sub>-supplemented mice. Number of FoxP3<sup>+</sup>-expressing cells in pancreatic infiltrates as determined by manual counting of FoxP3<sup>+</sup> cells on immunostained paraffin sections ( $n = 6$ –14). Scatter plots show the median with interquartile range. Statistical significance was calculated using the Mann-Whitney test. ns, not significant; \*\*\* $P < 0.001$ .

healthy men with a mean baseline value of 70 nmol/L, then a vitamin D<sub>3</sub> dose of 12,500 IU/day or more will be needed to raise serum 25(OH)D<sub>3</sub> concentrations to 290 nmol/L. These doses are thus completely out of the range of the suggested supplements for bone health (39) and may lead to calcemic side effects when administered long-term. At present, it is thus not advised to systematically administer these high doses in humans, because randomized intervention trials will be needed to demonstrate or refute the potential of these mega doses in diabetes prevention in humans.

In conclusion, lifelong dietary supplementation with high doses of regular vitamin D<sub>3</sub> brings 25(OH)D<sub>3</sub> concentrations above 230 nmol/L, prevents diabetes in NOD mice in a safe way, and is accompanied by induction of Tregs. This preclinical study confirms the potential of exploiting the vitamin D system in the prevention of type 1 diabetes in humans, but the fact that only lifelong supplementation with high doses altered disease presentation warns for overly optimistic statements on duration of treatment.

**Acknowledgments.** The authors thank Erik Van Herck, Ivo Jans, Suzanne Marcelis, Jaak Billen, Karolien Ciotkowski, and Martine Gillis (CEE, KU Leuven, Belgium) for excellent technical assistance and also Riet Van Looveren, Karen Moermans, Ingrid Stockmans, Nick van Gastel, and Steve Stegen (CEE, KU Leuven, Belgium) for their assistance with the  $\mu$ CT analysis.

**Funding.** This work was supported by grants from the Fund for Scientific Research Flanders (FWO-Vlaanderen G.0734.10), the University of Leuven (KU Leuven GOA 2009/10 and 2014/10), and the European Community's Health Seventh Framework Programme (FP7/2009-2014 under grant agreement 241447 with acronym NAIMIT). T.T. is supported through a PDM postdoctoral fellowship of the KU Leuven. L.D. is supported through an IRO fellowship of the

KU Leuven. F.D. received support from the Italian Ministry of Research, the Italian Ministry of Health, and the Tuscany Region. C.A.G. is supported by the KU Leuven. C.M. is a clinical researcher of the FWO-Vlaanderen.

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** T.T., L.D., F.B., I.S., G.S., and J.L. acquired data. T.T., A.V., G.C., F.D., C.A.G., and C.M. interpreted the data. T.T., A.V., G.C., F.D., T.L.V.B., C.A.G., and C.M. drafted the manuscript. T.L.V.B., C.A.G., and C.M. revised the manuscript critically for important intellectual content. All authors gave final approval of the version to be published. T.T., C.A.G., and C.M. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- van Belle TL, Coppieters KT, von Herrath MG. Type 1 diabetes: etiology, immunology, and therapeutic strategies. *Physiol Rev* 2011;91:79–118
- Eisenbarth GS. Type 1 diabetes mellitus. A chronic autoimmune disease. *N Engl J Med* 1986;314:1360–1368
- EURODIAB ACE Study Group. Variation and trends in incidence of childhood diabetes in Europe. *Lancet* 2000;355:873–876
- Mohr SB, Garland CF, Gorham ED, Garland FC. The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide. *Diabetologia* 2008;51:1391–1398
- Vitamin D supplement in early childhood and risk for Type 1 (insulin-dependent) diabetes mellitus. The EURODIAB Substudy 2 Study Group. *Diabetologia* 1999;42:51–54
- Hyppönen E. Vitamin D and increasing incidence of type 1 diabetes—evidence for an association? *Diabetes Obes Metab* 2010;12:737–743
- Giulietti A, Gysemans C, Stoffels K, et al. Vitamin D deficiency in early life accelerates Type 1 diabetes in non-obese diabetic mice. *Diabetologia* 2004;47:451–462
- Zella JB, McCary LC, DeLuca HF. Oral administration of 1,25-dihydroxyvitamin D<sub>3</sub> completely protects NOD mice from insulin-dependent diabetes mellitus. *Arch Biochem Biophys* 2003;417:77–80
- Stene LC, Ulriksen J, Magnus P, Joner G. Use of cod liver oil during pregnancy associated with lower risk of Type 1 diabetes in the offspring. *Diabetologia* 2000;43:1093–1098
- Sørensen IM, Joner G, Jenum PA, Eskild A, Torjesen PA, Stene LC. Maternal serum levels of 25-hydroxy-vitamin D during pregnancy and risk of type 1 diabetes in the offspring. *Diabetes* 2012;61:175–178
- Bruns ME, Bruns DE, Avioli LV. Vitamin D-dependent calcium-binding protein of rat intestine: changes during postnatal development and sensitivity to 1,25-dihydroxycholecalciferol. *Endocrinology* 1979;105:934–938
- Vaarala O. Leaking gut in type 1 diabetes. *Curr Opin Gastroenterol* 2008;24:701–706
- Mathieu C, van Etten E, Gysemans C, Decallonne B, Bouillon R. Seasonality of birth in patients with type 1 diabetes. *Lancet* 2002;359:1248
- Mathieu C, Waer M, Laureys J, Rutgeerts O, Bouillon R. Prevention of autoimmune diabetes in NOD mice by 1,25 dihydroxyvitamin D<sub>3</sub>. *Diabetologia* 1994;37:552–558
- Verhaeghe J, Van Herck E, Van Bree R, Van Assche FA, Bouillon R. Osteocalcin during the reproductive cycle in normal and diabetic rats. *J Endocrinol* 1989;120:143–151
- Lieben L, Masuyama R, Torrekens S, et al. Normocalcemia is maintained in mice under conditions of calcium malabsorption by vitamin D-induced inhibition of bone mineralization. *J Clin Invest* 2012;122:1803–1815
- Gysemans C, Waer M, Laureys J, et al. Islet xenograft destruction in the hu-PBL-severe combined immunodeficient (SCID) mouse necessitates anti-CD3 preactivation of human immune cells. *Clin Exp Immunol* 2000;121:557–565
- Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* 2003;299:1057–1061

19. Setoguchi R, Hori S, Takahashi T, Sakaguchi S. Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)-2 and induction of autoimmune disease by IL-2 neutralization. *J Exp Med* 2005;201:723–735
20. Zelenay S, Lopes-Carvalho T, Caramalho I, Moraes-Fontes MF, Rebelo M, Demengeot J. Foxp3+ CD25- CD4 T cells constitute a reservoir of committed regulatory cells that regain CD25 expression upon homeostatic expansion. *Proc Natl Acad Sci U S A* 2005;102:4091–4096
21. Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011;96:53–58
22. Zipitis CS, Akobeng AK. Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child* 2008;93:512–517
23. Dong JY, Zhang WG, Chen JJ, Zhang ZL, Han SF, Qin LQ. Vitamin D intake and risk of type 1 diabetes: a meta-analysis of observational studies. *Nutrients* 2013;5:3551–3562
24. Simpson M, Brady H, Yin X, et al. No association of vitamin D intake or 25-hydroxyvitamin D levels in childhood with risk of islet autoimmunity and type 1 diabetes: the Diabetes Autoimmunity Study in the Young (DAISY). *Diabetologia* 2011;54:2779–2788
25. Miettinen ME, Reinert L, Kinnunen L, et al. Serum 25-hydroxyvitamin D level during early pregnancy and type 1 diabetes risk in the offspring. *Diabetologia* 2012;55:1291–1294
26. Gysemans CA, Cardozo AK, Callewaert H, et al. 1,25-Dihydroxyvitamin D<sub>3</sub> modulates expression of chemokines and cytokines in pancreatic islets: implications for prevention of diabetes in nonobese diabetic mice. *Endocrinology* 2005;146:1956–1964
27. Fujita H, Sugimoto K, Inatomi S, et al. Tight junction proteins claudin-2 and -12 are critical for vitamin D-dependent Ca<sup>2+</sup> absorption between enterocytes. *Mol Biol Cell* 2008;19:1912–1921
28. Ooi JH, Li Y, Rogers CJ, Cantorna MT. Vitamin D regulates the gut microbiome and protects mice from dextran sodium sulfate-induced colitis. *J Nutr* 2013;143:1679–1686
29. Ooi JH, Chen J, Cantorna MT. Vitamin D regulation of immune function in the gut: why do T cells have vitamin D receptors? *Mol Aspects Med* 2012;33:77–82
30. Roy S, Shrinivas K, Bagchi B. Vitamin D sensitivity to the immune responses and autoimmunity: A chemical network model study [Internet], 2013. Ithaca, NY. Cornell University Library. Available from <http://arxiv.org/abs/1304.7193>. Accessed 13 June 2013
31. Marjamäki L, Niinistö S, Kenward MG, et al. Maternal intake of vitamin D during pregnancy and risk of advanced beta cell autoimmunity and type 1 diabetes in offspring. *Diabetologia* 2010;53:1599–1607
32. Brekke HK, Ludvigsson J. Vitamin D supplementation and diabetes-related autoimmunity in the ABIS study. *Pediatr Diabetes* 2007;8:11–14
33. Hawa MI, Valorani MG, Buckley LR, et al. Lack of effect of vitamin D administration during pregnancy and early life on diabetes incidence in the nonobese diabetic mouse. *Horm Metab Res* 2004;36:620–624
34. Nicholson LB, Kuchroo VK. Manipulation of the Th1/Th2 balance in autoimmune disease. *Curr Opin Immunol* 1996;8:837–842
35. Weiss L, Letimier FA, Carriere M, et al. In vivo expansion of naive and activated CD4+CD25+FOXP3+ regulatory T cell populations in interleukin-2-treated HIV patients. *Proc Natl Acad Sci U S A* 2010;107:10632–10637
36. Casteels K, Waer M, Laureys J, et al. Prevention of autoimmune destruction of syngeneic islet grafts in spontaneously diabetic nonobese diabetic mice by a combination of a vitamin D<sub>3</sub> analog and cyclosporine. *Transplantation* 1998;65:1225–1232
37. Baeke F, Korf H, Overbergh L, et al. The vitamin D analog, TX527, promotes a human CD4+CD25highCD127low regulatory T cell profile and induces a migratory signature specific for homing to sites of inflammation. *J Immunol* 2011;186:132–142
38. Jeffery LE, Burke F, Mura M, et al. 1,25-Dihydroxyvitamin D<sub>3</sub> and IL-2 combine to inhibit T cell production of inflammatory cytokines and promote development of regulatory T cells expressing CTLA-4 and FoxP3. *J Immunol* 2009;183:5458–5467
39. Bouillon R, Van Schoor NM, Gielen E, et al. Optimal vitamin D status: a critical analysis on the basis of evidence-based medicine. *J Clin Endocrinol Metab* 2013;98:E1283–E1304
40. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr* 2003;77:204–210